

# Received Level Estimation, Behavioral Response, and Diel Behavior Analyses for Delphinids Tagged at the Pacific Missile Range Facility in 2021-2022

E. Elizabeth Henderson, *NIWC Pacific*

Cameron Martin, *NIWC Pacific*

Annette E. Harnish, *Cascadia Research Collective*

Michaela A. Kratofil, *Cascadia Research Collective*

Robin W. Baird, *Cascadia Research Collective*

Steve W. Martin *National Marine Mammal Foundation*

16 January 2024

**Naval Information  
Warfare Center**



**PACIFIC**

Cascadia  
Research  
Collective



non-profit research and education since 1979



NATIONAL  
MARINE MAMMAL  
FOUNDATION

## EXECUTIVE SUMMARY

This report summarizes the results of satellite-tagging efforts conducted prior to Submarine Command Course (SCC) training events from August 2021 and 2022 on the Pacific Missile Range Facility (PMRF) off the island of Kaua'i. Twenty-one delphinids were satellite tagged with the intent of observing their behavior before, during, and after the SCCs and estimating received levels of mid-frequency active sonar (MFAS). Of these, 16 individuals remained in the area or with direct paths between their location and sources of surface ship hull-mounted MFAS such that received levels could be estimated using propagation modeling. These included six short-finned pilot whales (*Globicephala macrorhynchus*), one false killer whale (*Pseudorca crassidens*), five melon-headed whales (*Peponocephala electra*), one rough-toothed dolphin (*Steno bredanensis*), and two common bottlenose dolphins (*Tursiops truncatus*). The analytical methods developed and described in Henderson et al. (2021) were applied to the resulting data; these included smoothing the animal tracks in 5-min steps using the R-package *crawl*, calculating the 95% confidence error ellipse around each step location, and then modeling transmission loss along multiple radials through the error ellipse from the sea surface to the seafloor. Dives were also modeled using behavior logs from the tags so that the depth of the animal at the time of each transmission per 5-min bin could also be considered when estimating the received levels (RLs). The median RL for each 5-min bin was calculated based on the aggregation of all transmissions and radials per bin, this plus two times the standard deviation ( $\pm 2*SD$ ) are reported as the range of possible exposure values. Dive behavior variables were also analyzed within diel period (dawn, day, dusk, night) and across SCC phase (before, Phase A, Interphase, Phase B, after) to determine if dive behavior changed relative to the presence of training activity or MFAS.

For all animals, the maximum median received sound pressure levels ranged from 72.6 to 147.7 dB re 1  $\mu$ Pa. In fact, as many of the animals had moved out of the area prior to the onset of MFAS, maximum median RLs were at or below 100 dB re 1  $\mu$ Pa for eight of the animals. The other eight were on or near the range during periods of MFAS and therefore had relatively higher estimated RLs, although no estimated RL exceeded 156.2 dB re 1  $\mu$ Pa (including  $+2*SD$ ). Some differences in dive behavior were noted as well, including changes in median dive depth and median dive duration between SCC phases, as well as changes in dive rates or the amount of time spent at the surface. However, there were no consistent changes in behavior either within or across species; these variables differed between the SCC phases in a variety of ways across individuals. In addition, no changes in horizontal movement were observed relative to periods of MFAS. All tagged animals were also compared to Cascadia Research Collective's photo-identification catalog; most were matched to existing animals in the catalog, with one new bottlenose dolphin and two new rough-toothed dolphins added. Of the pre-existing identifications, the false killer whale, three of the pilot whales, two of the bottlenose dolphins, and one of the rough-toothed dolphins all belong to known Hawaiian island populations, and the latter two belong to Kaua'i island-specific populations. These data, aggregated with previously tagged or photo-identified animals from Kaua'i and the other Hawaiian Islands, build the basis for understanding long-term consequences of exposure at the population level as well as the potential to identify and understand the impacts of repeated exposures. As most of the tagged animals in this study are island (or Islands) residents, it is possible they have been exposed to MFAS previously. Since none of the animals demonstrated consistent or overt responses, it is also possible they have habituated to (or at least tolerate) MFAS and other training-associated sounds. However, with additional future data and continued improvements in

analytical methods for satellite tag data, these aggregated data may be able to be used in finer-scale behavioral response analyses, and some of these uncertainties may be addressed.

# CONTENTS

EXECUTIVE SUMMARY .....	2
LIST OF FIGURES .....	5
LIST OF TABLES .....	11
INTRODUCTION .....	15
METHODS .....	16
Field Operations .....	16
Tag Programming .....	17
Data Processing .....	18
Dive Analysis .....	18
Acoustic Data, MFAS Detection, and Localization .....	19
Acoustic Propagation Modeling .....	19
Ship Exposures .....	23
Behavioral Response and Diel Analysis .....	24
RESULTS .....	26
Tagging and Photo-Identification .....	26
Behavioral Response, Diel Analysis, and Received Level Estimation .....	28
Short-finned pilot whales .....	29
False killer whales .....	50
Melon-headed whales .....	53
Rough-toothed dolphins .....	67
Common bottlenose dolphins .....	74
DISCUSSION .....	82
<b>LITERATURE CITED</b> .....	<b>87</b>
APPENDIX .....	92
Additional dive behavior results and movement narratives .....	92
Short-finned pilot whales .....	92
False killer whale .....	95
Melon-headed whales .....	100
Rough-toothed dolphins .....	102
Common bottlenose dolphins .....	107

## LIST OF FIGURES

Figure 1 – The Barking Sands Underwater Range Expansion (BSURE) is outlined in purple, the Barking Sands Tactical Underwater Range (BARSTUR) is outlined in yellow, and the Shallow Water Training Range (SWTR) is outlined in green. Ship activity primarily occurs on BSURE, but animals are considered to be on the range if they overlap with any portion of the range (Taken from Henderson et al. 2021). ...	16
Figure 2 – Example of radial slices from ship through the error ellipse of a tag location. Blue diamonds approximate location of hydrophones for MFAS detection. Lower right shows a portion of the Kaua'i coastline. ....	20
Figure 3 – Example of single slice from ship MFAS source full depth maximum distance estimated receive level in units of dB re 1 $\mu$ Pa. Dark color indicates the seafloor, highlighting the steep angle limitation close to source. The red insert box illustrates potential location in this slice of the animal location using the 95% CI ellipse and depth information. ....	21
Figure 4 – 3D concept utilizing multiple radials from MFAS source through an animal's estimated location in 3D space. ....	22
Figure 5 – Example of all estimated receive levels for a single MFAS transmission via 25 slices from the MFAS ship to one estimated possible 95% CI location in three dimensions. ....	22
Figure 6 - Plan view of exposure geometry for tag GmTag235 on 8/20/2022 22:23 GMT illustrating the situation where the animal may, or may not, have been exposed to MFAS depending on where it actually was within its 95% confidence interval location uncertainty. To account for this, an estimated probability of exposure metric is included for all estimated exposures to MFAS. ....	23
Figure 7 - Median RLs for GmTag232 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the $\pm 2$ SD values. ....	33
Figure 8 - Median RLs for GmTag233 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the $\pm 2$ SD values. ....	34
Figure 9 - Median RLs for GmTag234 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the $\pm 2$ SD values. ....	35
Figure 10 - Movements of GmTag233 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level, and time is given in GMT. Additionally, the RL circles are colored by "intensity" which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygon represents the area of ship activity during each of the three MFAS bouts that GmTag233 was exposed to and the corresponding diamond point represents the mean ship location during the bouts. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary. ....	36

Figure 11 - Top. Boxplot showing dive depths of GmTag232 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag232 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 80.4 dB re 1  $\mu$ Pa. .... 39

Figure 12 - Top. Boxplot showing dive depths of GmTag233 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag233 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 81.2 dB re 1  $\mu$ Pa. .... 40

Figure 13 - Top. Boxplot showing dive depths of GmTag234 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag234 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 72.6 dB re 1  $\mu$ Pa. .... 41

Figure 14 - Median RLs for GmTag235 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values. .... 43

Figure 15 - Median RLs for GmTag236 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values. .... 44

Figure 16 - Median RLs for GmTag237 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values. .... 45

Figure 17 - Top: Movements of GmTag237 during the August 2022 SCC event, including extensive movements away from the range after exposure phases. Bottom: zoomed-in panel of the track to highlight movements that occurred before, during, and after the exposure, prior to the animal’s extensive movements away from the area. Note that for this particular event, ship-based MFAS activity occurred during both A and B phases. The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level, and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the MFAS bouts that GmTag237 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary. .... 46

Figure 18 - Top. Boxplot showing dive depths of GmTag235 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag235 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 129.2 dB re 1  $\mu$ Pa. .... 48

Figure 19 - Top. Boxplot showing dive depths of GmTag237 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag237 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 128.8 dB re 1  $\mu$ Pa. .... 49

Figure 20 - Median RLs for PcTag074 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values. .... 51

Figure 21 - Movements of PcTag074 during and after the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the three MFAS bouts that PcTag074 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary. .... 52

Figure 22 - Median RLs for PeTag029 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values. .... 55

Figure 23 - Top: Movements of PeTag029 prior to the August 2021 SCC event; this tag stopped transmitting before Phase B of the SCC event, but was exposed to MFAS during a ULT period that occurred prior to Phase A of the SCC (see text for description of phases). Bottom: Movements of PeTag029 during the MFAS exposures that occurred during the ULT event. The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The gray shaded rectangular polygon represents the area of ship activity during the ULT and the corresponding diamond point represents the mean ship location during the ULT. The dashed black line represents the PMRF boundary. .... 56

Figure 24 - Median RLs for Pe31 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values. .... 58

Figure 25 - Median RLs for PeTag032 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values. .... 59

Figure 26 - Movements of PeTag031 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the three MFAS bouts that PeTag031 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary. .... 60

Figure 27 - Movements of PeTag032 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular

polygons represent the area of ship activity during each of the three MFAS bouts that PeTag032 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary. .... 61

Figure 28 - Top. Boxplot showing dive depths of PeTag031 by SCC Phase and time of day. Bottom. Barplot showing dive rates of PeTag031 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 136.6 dB..... 62

Figure 29 - Median RLs for PeTag033 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values..... 63

Figure 30 - Median RLs for PeTag034 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values..... 64

Figure 31 - Movements of PeTag033 during the August 2022 SCC event, including extensive movements away from the range after exposure phases. Note that for this particular event, ship-based MFAS activity occurred during both A and B phases. The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the MFAS bouts that PeTag033 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary. .... 65

Figure 32 - Movements of PeTag034 during the August 2022 SCC event, including extensive movements away from the range after exposure phases. Note that for this particular event, ship-based MFAS activity occurred during both A and B phases. The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the MFAS bouts that PeTag034 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary. .... 66

Figure 33 - Median RLs for SbTag023 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values..... 70

Figure 34 - Movements of SbTag023 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times are given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular



polygons represent the area of ship activity during each of the three MFAS bouts that SbTag023 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary. .... 71

Figure 35 - Top. Boxplot showing dive depths of SbTag023 by SCC Phase and time of day. Bottom. Barplot showing dive rates of SbTag023 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 141.9 dB..... 73

Figure 36 - Median RLs for TtTag039 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values..... 77

Figure 37 - Movements of TtTag039 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the three MFAS bouts that TtTag039 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary. .... 78

Figure 38 - Top. Boxplot showing dive depths of TtTag039 by SCC Phase and time of day. Bottom. Barplot showing dive rates of TtTag039 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 100.0 dB..... 79

Figure 39 - Median RLs for Tt040 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values..... 80

Figure 40 - Movements of TtTag040 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the three MFAS bouts that TtTag040 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary. .... 81

Figure 41 - Top. Boxplot showing dive depths of GmTag236 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag236 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 128.8 dB..... 94

Figure 42 - Top. Boxplot showing dive depths of PcTag074 by SCC Phase and time of day. Bottom. Barplot showing dive rates of PcTag074 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 90.5 dB..... 99

Figure 43 - Top. Map showing crawl model trackline of PeTag035 during the 2022 August SCC. Bottom. Map showing crawl model trackline of PeTag036 during the 2022 August SCC. .... 101

Figure 44 - Top. Boxplot showing dive depths of SbTag025 by SCC Phase and time of day. Bottom. Barplot showing dive rates of SbTag025 by SCC Phase and time of day. .... 104

Figure 45 - Map showing crawl model trackline of SbTag024 during the 2021 August SCC. .... 105

Figure 46 - Map showing crawl model trackline of SbTag025 during the 2022 August SCC. This individual was not exposed to Navy activities..... 106

Figure 47 - Top. Boxplot showing dive depths of TtTag041 by SCC Phase and time of day. Bottom. Barplot showing dive rates of TtTag041 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 100.5 dB..... 109

Figure 48 - Map showing crawl model trackline of TtTag041 during the 2022 August SCC. This individual was not exposed to Navy activities..... 111

## LIST OF TABLES

Table 1 - Summary of tag programming regimes by species, year, and tag type. When there are multiple programming regimes for a single tag type and species within the same year, each regime is given a separate line. Gm = Pilot whale (*Globicephala macrorhynchus*), Pe = Melon-headed whale (*Peponocephala electra*), Pc = False killer whale (*Pseudorca crassidens*), Tt = Common bottlenose dolphin (*Tursiops truncatus*), Sb = Rough-toothed dolphin (*Steno bredanensis*). ..... **Error! Bookmark not defined.**

Table 2 - Coverage requirements for each phase by time of day and species for inclusion in statistical comparisons between phases. Gm = Pilot whale (*Globicephala macrorhynchus*), Pe = Melon-headed whale (*Peponocephala electra*), Pc = False killer whale (*Pseudorca crassidens*), Tt = Common bottlenose dolphin (*Tursiops truncatus*), Sb = Rough-toothed dolphin (*Steno bredanensis*). ..... **Error! Bookmark not defined.**

Table 3 - Tag Deployment data for 2021 and 2022 satellite tags. An “NA” in the maximum received level column indicates that there was no overlap between the specified tag and MFAS blocks. Max RL = Maximum estimated received level..... **Error! Bookmark not defined.**

Table 4 - SCC Phase times and passive acoustic monitoring data durations (times given in HST) ..... **Error! Bookmark not defined.**

Table 5 - Percentage of dive/surfacing data by phase for short-finned pilot whales. The percentage of behavioral coverage is defined as the proportion of the duration of behavioral data relative to the duration of the tag within each phase..... **Error! Bookmark not defined.**

Table 6 - A comparison of dawn diving parameters from short-finned pilot whales exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

Table 7 - A comparison of nighttime diving parameters from short-finned pilot whales exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

Table 8 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag232. .... **Error! Bookmark not defined.**

Table 9 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag233.  
..... **Error! Bookmark not defined.**

Table 10 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag234.  
..... **Error! Bookmark not defined.**

Table 11 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag235.  
..... **Error! Bookmark not defined.**

Table 12 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag234.  
..... **Error! Bookmark not defined.**

Table 13 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag237.  
..... **Error! Bookmark not defined.**

Table 14 - Percentage of dive/surfacing data by phase for false killer whale Pctag074. The percentage of behavioral coverage is defined as the proportion of the duration of behavioral data relative to the duration of the tag within each phase..... **Error! Bookmark not defined.**

Table 15 - Estimated received levels, cumulative sound exposure levels, and ship CPA for Pctag074**Error! Bookmark not defined.**

Table 16 - Percentage of dive/surfacing data by phase for melon-headed whales. The percentage of behavioral coverage is defined as the proportion of the duration of behavioral data relative to the duration of the tag within each phase..... **Error! Bookmark not defined.**

Table 17 - A comparison of nighttime diving parameters from melon-headed whales exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

Table 18 - Estimated received levels, cumulative sound exposure levels, and ship CPA for Petag029**Error! Bookmark not defined.**

Table 19 - Estimated received levels, cumulative sound exposure levels, and ship CPA for Petag031**Error! Bookmark not defined.**

Table 20 - Estimated received levels, cumulative sound exposure levels, and ship CPA for Petag032**Error! Bookmark not defined.**

Table 21 - Estimated received levels, cumulative sound exposure levels, and ship CPA for Petag033**Error! Bookmark not defined.**

Table 22 - Estimated received levels, cumulative sound exposure levels, and ship CPA for PeTag034 **Error! Bookmark not defined.**

Table 23 - Percentage of dive/surfacing data by phase for rough-toothed dolphins. The percentage of behavioral coverage is defined as the proportion of the duration of behavioral data relative to the duration of the tag within each phase..... **Error! Bookmark not defined.**

Table 24 - A comparison of daytime diving parameters from rough-toothed dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

Table 25 - A comparison of nighttime diving parameters from rough-toothed dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

Table 26 - Estimated received levels, cumulative sound exposure levels, and ship CPA for SbTag023 **Error! Bookmark not defined.**

Table 27 - Percentage of dive/surfacing data by phase for common bottlenose dolphins. The percentage of behavioral coverage is defined as the proportion of the duration of behavioral data relative to the duration of the tag within each phase..... **Error! Bookmark not defined.**

Table 28 - A comparison of daytime diving parameters from common bottlenose dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken. ... **Error! Bookmark not defined.**

Table 29 - A comparison of nighttime diving parameters from common bottlenose dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken. ... **Error! Bookmark not defined.**

Table 30 - Estimated received levels, cumulative sound exposure levels, and ship CPA for TtTag039 **Error! Bookmark not defined.**

Table 31 - Estimated received levels, cumulative sound exposure levels, and ship CPA for TtTag040 **Error! Bookmark not defined.**

Table 32 - A comparison of dusk diving parameters from short-finned pilot whales exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

Table 33 - A comparison of dawn diving parameters for false killer whale PcTag074 exposed to MFAS for phases that meet the required coverage cutoff. Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

Table 34 - A comparison of daytime diving parameters for false killer whale PcTag074 exposed to MFAS for phases that meet the required coverage cutoff. Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

Table 35 - A comparison of dusk diving parameters for false killer whale PcTag074 exposed to MFAS for phases that meet the required coverage cutoff. .... **Error! Bookmark not defined.**

Table 36 - A comparison of nighttime diving parameters for false killer whale PcTag074 exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

Table 37 - A comparison of dawn diving parameters from rough-toothed dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

Table 38 - A comparison of dusk diving parameters from rough-toothed dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where

significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

Table 39 - A comparison of dawn diving parameters from common bottlenose dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken. ... **Error! Bookmark not defined.**

Table 40 - A comparison of dusk diving parameters from common bottlenose dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken..... **Error! Bookmark not defined.**

## INTRODUCTION

Satellite tagging of cetaceans provides valuable information on habitat use, movement patterns, and dive behavior over potentially long periods and large areas, and with repeated efforts on focal species, population-level information can be developed. For example, Cascadia Research Collective (CRC) has conducted 20 years of vessel-based surveys in the Hawaiian Islands, using both photo-identification capture-recapture and satellite tagging methods (e.g., Baird et al. 2009; 2012; 2013). Through these studies, CRC has identified resident and offshore populations of multiple species of odontocetes (e.g., Baird et al. 2008a, 2008b, 2022), helped establish biologically important areas for many of these species (Baird et al. 2015a; Kratofil et al. 2023), and assessed behavioral responses to anthropogenic impacts including fisheries interactions, tourism, and Navy sonar (Baird et al. 2009, 2015b, 2020; Thorne et al. 2012; Van Cise et al. 2021). Due to the presence of a Navy testing and training range in the Hawaiian Islands, the Pacific Missile Range Facility (PMRF), not only can behavioral responses to Navy testing and training activity be assessed, but received levels of mid frequency active sonar (MFAS) can be estimated for the tagged animals in three-dimensional space (Baird et al. 2014, 2017; Henderson et al. 2021; Martin and Manzano-Roth 2012). Similar long-term studies of cetacean populations have been conducted for other species in other areas where Navy ranges are present, such as fin (*Balaenoptera physalus*) and Cuvier's beaked whales (*Ziphius cavirostris*) in southern California (Falcone et al. 2009, 2017, 2022; Schorr et al. 2017), and Blainville's beaked whales (*Mesoplodon densirostris*) in the Bahamas (Claridge 2013; Claridge and Dunn 2013; Joyce et al. 2020). These long-term, multi-modality studies record exposures and responses to real-world training scenarios and can be used to assess effects beyond a single exposure level for an individual animal, including an estimation of the effects of repeated exposures and the impacts of long-term exposures at the population level.

Continuing this work, CRC returned to the PMRF range in August 2021 and August 2022 to tag odontocetes prior to and during a two-week biannual training event, the Submarine Command Course (SCC). CRC was supported in this work by personnel from Naval Underwater Warfare Center, Newport (NUWC) and Naval Information Warfare Center, Pacific (NIWC), who directed CRC to locations of acoustic detections on the range of odontocete species of interest. Potential focal species included short-finned pilot whales (*Globicephala macrorhynchus*), false killer whales (*Pseudorca crassidens*), pygmy killer whales (*Feresa attenuata*), melon-headed whales (*Peponocephala electra*), common bottlenose dolphins (*Tursiops truncatus*), rough-toothed dolphins (*Steno bredanensis*), and when possible, Blainville's beaked whales and killer whales (*Orcinus orca*). The goals of this work were to build on the existing species' databases of home range and habitat use, and to estimate received levels of MFAS and assess any potential behavioral responses to the MFAS. This report focuses on movement patterns, estimated received levels of mid-frequency active sonar, and behavioral responses of delphinid odontocetes tagged in 2021 and 2022. Similar information on beaked whales tagged during the same field efforts are reported separately (Henderson et al. 2024).



## METHODS

### Field Operations

Field operations were undertaken between Kaua'i and Ni'ihau in August 2021, and August 2022, with tagging conducted from a 7.3-m rigid-hulled inflatable boat (RHIB). To maximize the likelihood of data being obtained for periods before, during, and after biannual SCCs, survey efforts occurred immediately prior to each SCC. Odontocetes were located both on and off the Navy's hydrophone range at PMRF. For encounters on the range, acoustic detections from the array were used to direct the RHIB to the general area where odontocetes were acoustically detected. During encounters, information was recorded on group size and start and end time and locations (see Baird et al. 2013). To identify individuals and to determine age (based on the degree of scarring and relative size) and sex (based on species-specific characteristics like sexual dimorphism, when possible), photographs of all individuals within the groups were taken. For all species except for melon-headed whales, photos were compared to long-term photo-identification catalogs (Baird et al. 2008, 2009; Mahaffy et al. 2015, 2023) to assess sighting history and the potential for repeat tagging of individuals.

Tags used included both location-only (SPOT6) and depth-transmitting SPLASH-10 or SPLASH10-F tags (Wildlife Computers, Redmond, WA) in the Low Impact Minimally Percutaneous Electronic Transmitter configuration (Andrews et al., 2008). SPLASH10-F tags also transmitted Fastloc®-GPS locations. Tags were deployed with a Dan-Inject pneumatic projector and were attached with two gas-sterilized surgical grade titanium darts (see Schorr et al. 2009; Baird et al. 2011). For smaller species (i.e., bottlenose dolphins, rough-toothed dolphins, melon-headed whales) 4.4-cm darts with one row of backward facing petals were used, while for larger species (i.e., short-finned pilot whales, false killer whales) 6.7-cm darts with two rows of backward facing petals were used.




Figure 1 – The Barking Sands Underwater Range Expansion (BSURE) is outlined in purple, the Barking Sands Tactical Underwater Range (BARSTUR) is outlined in yellow, and the Shallow Water Training Range (SWTR) is outlined in green. Ship activity primarily occurs on BSURE, but animals are considered to be on the range if they overlap with any portion of the range (Taken from Henderson et al. 2021).

## Tag Programming

SPOT6 and SPLASH10 tags were Argos location-only tags, while SPLASH10-F tags also transmitted Fastloc®-GPS locations. Tags were programmed to transmit during between 11 and 16 hours per day (depending on tag type and species, see Table 1), with transmissions prioritized during hours of the day with the greatest density of satellite overpasses, based on Argos pass predictions obtained through CLS. For SPLASH tags, transmitted dive data consisted of behavior logs recording the start and end time of dive and surface periods, and some tags were also programmed to transmit time series data every sixth day (see Table 1). All SPLASH tags were programmed to record dives greater than or equal to 50 meters and lasting longer than 30 seconds to reduce gaps in the behavior logs (Quick et al., 2019). Surface periods were considered any time when the animal did not dive below 50 m. For each dive greater than the 50-m dive depth threshold, the maximum dive depth and duration were recorded, with the start and end of the dives determined [by the wet/dry sensor; by when the animal dove below 3 m]. SPLASH10-F tags were programmed to obtain up to two Fastloc-GPS locations per hour, and 48 locations per day, with Fastloc-GPS locations set as high priority (three out of every four transmissions) and behavior logs and time series set as low priority (one out of every four transmissions). To maximize the likelihood of obtaining behavioral data before, during, and after the SCC, tags were recorded to collect behavior logs and Fastloc®-GPS locations starting approximately three days prior to the start of the SCC and ending approximately three days after the end of the SCC, and transmit behavior and Fastloc®-GPS data with a 6-day buffer. There were two shore-based Argos receivers (Wildlife Computers MOTEs) on Ni‘ihau and Kaua‘i that were also used to increase data throughput when tagged animals were within range of the receivers (Jeanniard-du-Dot et al., 2017), and an Argos goniometer was used during all small-boat operations to potentially obtain additional behavior data, times when animals were known to be at the surface, and Fastloc®-GPS locations from SPLASH10 and SPLASH10-F tags.

*Table 1 - Summary of tag programming regimes by species, year, and tag type. When there are multiple programming regimes for a single tag type and species within the same year, each regime is given a separate line. Gm = Pilot whale (*Globicephala macrorhynchus*), Pe = Melon-headed whale (*Peponocephala electra*), Pc = False killer whale (*Pseudorca crassidens*), Tt = Common bottlenose dolphin (*Tursiops truncatus*), Sb = Rough-toothed dolphin (*Steno bredanensis*).*

Species	Year	Tag type	Data	# Hours transmitting per day
Gm	2021	SPLASH10-F	Argos, Fastloc, Behavior	16
Pe	2021	SPLASH10-F	Argos, Fastloc, Behavior	16
Pc	2021	SPLASH10-F	Argos, Fastloc, Behavior	16
Sb	2021	SPLASH10	Argos, Behavior, Time Series (1 min 15 sec)	11
Sb	2021	SPOT6	Argos	15
Tt	2021	SPLASH10-F	Argos, Fastloc, Behavior	16
Gm	2022	SPLASH10-F	Argos, Fastloc, Behavior, Time Series (5 min)	14
Pe	2022	SPLASH10-F	Argos, Fastloc, Behavior, Time Series (5 min)	14
Pe	2022	SPLASH10-F	Argos, Fastloc, Behavior	16
Sb	2022	SPLASH10	Argos, Behavior, Time Series (5 min)	14
Tt	2022	SPLASH10	Argos, Behavior, Time Series (5 min)	14

## Data Processing

Argos and GPS location data were processed following methods detailed in Kratofil et al. (2023). To summarize, the Douglas-Argos Filter (Douglas et al., 2012) accessed through Movebank (Kranstauber et al., 2011) was used to filter Argos data to remove erroneous locations. Residual error values ( $< 35$ ; Dujon et al., 2014) and time errors ( $< 10$  seconds) were used to filter Fastloc-GPS locations; these were additionally processed through a general speed filter via Movebank (Kranstauber et al., 2011). Resultant Argos and GPS locations were combined, and then fitted to a continuous time correlated random walk (CTCRW) model via the *crawl* package in R (Johnson et al., 2008; Johnson & London, 2018). Models were fit to each individual trajectory; subsequent models were then used to predict locations at 5-minute intervals used to estimate received levels following Henderson et al. (2021). The CTCRW model directly incorporates known measurement error in Argos (error ellipses) and GPS (user-defined, see Henderson et al. 2021) locations (Johnson et al., 2008; McClintock et al., 2014) while allowing for location prediction at user-defined intervals based on the modeled movement process. Predicted locations at the user-defined 5-minute intervals also include estimated standard errors (meters) in both x and y directions (easting and northing), which were used to account for positional uncertainty in received-level analyses. Finally, any locations on land were re-routed around land plus a 50-meter buffer using the *pathroutr* package (London, 2020).

To verify that the tags operated as intended and experienced no malfunctioning that could invalidate dive data, the dive behavior data were examined prior to analysis as described in Henderson et al. (2021). To ascertain whether there might have been a pressure transducer failure, we assessed Depth and ZeroDepthOffset values in tag status files for indication of drift. Those exceeding  $\pm 10$  meters and  $\pm 9$  meters, respectively, were flagged as potential pressure transducer failures. We also assessed data to identify and flag extreme ascent or descent rates (greater than 2-3 meters per second) in recorded dives that could indicate tag malfunctioning by dividing twice the dive depth by the dive duration.

## Dive Analysis

The SPLASH10 and SPLASH10-F tag data included a dive behavior log, which reported start and end times of dives and surface periods, as well as maximum dive depths. Using these data, coupled with the smoothed crawl tracks in 5-min intervals, full dive cycles were modeled in a custom Matlab program. For pilot whales, estimated ascent and descent rates were derived from pre-existing TDR tag data (see Henderson et al. 2021 for more details). For the other odontocete species, minimum and maximum bottom times were estimated based on Wildlife Computer's definitions of U, V, and Square-shaped dives. Then, ascent and descent times were estimated based on the remaining time in the dive divided by two, and ascent and descent rates were determined by using the mean maximum depth provided divided by the estimated ascent and descent times. These values (minimum and maximum bottom time, ascent and descent times, and ascent and descent rates) were estimated for all dives. The dive behavior logs, MOTE transmission logs, and goniometer files were also examined for completion in the record of surface and dive periods. If data were missing, these periods were noted as well, while times when tags were detected by the MOTE, Argos, or goniometer, even if no locations were available, were utilized as known surface times.

Next, dive durations were interpolated by 30 points per dive, leading to a timestamp approximately every 30 seconds for a deep dive and approximately every 10 seconds for a shallow ( $< 300$  m) dive. Dive depths were modeled using the estimated ascent and descent rates at each interpolated timestamp, and these were combined with the surface crawl track for a full record for each whale. At each

timestamp (either at 5-min intervals when at the surface or at the finer interpolated intervals of the dives), it was also noted whether that timestamp occurred during a dive, was at the surface but interpolated, or was at the surface with either an Argos or GPS update or MOTE uplink. Finally, it was noted whether that timestamp occurred during a period of missing behavioral log data. In that case, it was assumed the animal was at the surface if the interpolated timestamp occurred within a minute of an Argos or GPS location.

### Acoustic Data, MFAS Detection, and Localization

Acoustic data were recorded for 63 of the PMRF bottom-mounted range hydrophones in August 2021 and August 2022. Of the 63 hydrophones recorded, 36 had the broadband frequency response required for detecting MFAS transmissions. Thirty-one of these hydrophones had a frequency response from 50 Hz to 48 kHz and were at depths of 2400 to 4800 m, and the remaining 5 hydrophones had a frequency response from 100 Hz to 48 kHz and were at depths of 650 to 1750 m (Martin et al., 2015). The remaining 27 hydrophones had a frequency response from 8-48 kHz and therefore were sufficient to detect most delphinid vocalizations. A custom computer-based recorder collected data at a 96 kHz sample rate with 16-bit samples.

Detection of MFAS transmissions occurred in the mid-frequency band of 1-10 kHz. Raw timeseries data were processed with a 16,384-point fast Fourier transform (FFT) and 93.75% overlap. Both long-term and short-term running average spectra of the FFTs were computed in the two detection frequency bands. These were used to determine when the signal-to-noise ratio (SNR) exceeded a user defined threshold of 15 dB. To determine if an MFAS transmission was the first arrival (e.g., not multi-path), the maximum signal level had to be greater than one-half and one-third of the detected signal frequency to exclude the second and third harmonics as detections. If the number of user-defined consecutive detections was met (75 for the lower frequency band and 25 for the higher frequency band) or the SNR threshold was no longer met, a detection queue ended, and a 5 second blanking time was initiated before resuming detection to avoid detecting reverberations.

Model-based localizations of MFAS transmissions were performed using the suite of C++ algorithms originally developed for whale calls, described in detail by Martin et al. (2015, 2023). Briefly, the onset times of automatic detections across multiple hydrophones were used as the measured time of arrival (TOA), and measured TOAs were subtracted from each other to calculate the time difference of arrivals (TDOAs). A weighted least square error (LSE) between measured and modeled TDOAs was minimized by calculating modeled TDOAs from theoretical source locations determined by an iterative spatial grid search process. The LSE was weighted by the order of TOAs with more weighting for earlier arrivals and was normalized by the number of hydrophones in a localization solution. In addition, candidate detections for a localization solution were required to have a start frequency within 12 Hz of each other and a slope of 0.5 frequency bins/time bin. Localizations with a weighted LSE of 0.15 seconds between measured and modeled TDOAs were used for this analysis.

### Acoustic Propagation Modeling

The process of estimating received levels on whales from MFAS transmissions utilizes methods was described in detail in Henderson et al. (2021). To summarize, propagation modeling was done with the Peregrine parabolic equation propagation model developed by Oasis Ltd (Heaney and Campbell, 2016), based on the range dependent acoustic model (RAM) (Collins, 1993). National Oceanic and Atmospheric Administration National Geophysical Data Center U.S. Coastal Relief Model (NOAA National Geophysical

Data Center, 2011) was used to obtain bathymetric data with 3 arc-second resolution, and the 2018 World Ocean Atlas (Locarnini et al., 2018; Zweng et al., 2018) was consulted for historical sound speed profiles. Received levels were calculated based on publicly available source levels of U.S. Navy sonars (NMFS, 2007) and modeled transmission losses. Nominal source levels at 1 m distance for hull-mounted sonar (53C) is 235 dB re 1  $\mu$ Pa. The *crawl*-modeled x and y positional error from interpolated whale track locations (described in Data Processing) were used to define a 95% confidence interval error ellipse to represent uncertainty around each modeled whale location. The error ellipse was sampled with radial slices taken systematically in azimuth, and each radial slice associated with selected MFAS transmissions were a single propagation modeling run (Figure X, also see Henderson et al. 2021 for more details on this method).




Figure 2 – Example of radial slices from ship through the error ellipse of a tag location. Blue diamonds approximate location of hydrophones for MFAS detection. Lower right shows a portion of the Kaua'i coastline.

Modeling was performed across the full depth range from 0 to a maximum of 5400 m, and the distance of the longest radial slice for an MFAS transmission was used for all radials from the same MFAS transmission for ease of analysis. Peregrine output transmission loss calculations resulted in 600 depth bins with 9 m spacing, and 1000 range bins with variable spacing based on the distance of the longest radial slice. To reduce constructive and destructive interference from modeling a single frequency of an MFAS transmission and to better characterize the bandwidth of the signal, 10 log-spaced frequencies across 200 Hz of bandwidth around an MFAS transmission ( $\pm$  100 Hz) were modeled. Figure 2 illustrates a single radial slice from a ship through the maximum range for an exposure illustrating the single slice data from the ship source over 5400 m depth and 29.6 km distance color coded by the estimated receive level (RL).




Figure 3 – Example of single slice from ship MFAS source full depth maximum distance estimated receive level in units of dB re  $1\mu\text{Pa}$ . Dark color indicates the seafloor, highlighting the steep angle limitation close to source. The red insert box illustrates potential location in this slice of the animal location using the 95% CI ellipse and depth information.

To estimate the probable 3D location of the animal at the time of MFAS transmissions requires utilizing the animal's location in the depth dimension over the full water column of the 95% CI error ellipse. Animal depths were derived from the satellite tag data (See Dive Analysis section for more details on how this was modeled). For SbTag024, there was no depth data recorded with the SPOT6 tag, therefore two depth regimes were utilized to estimate received levels. These included a shallow regime to represent exposures when the animal was within the upper 30 m of the surface, and a dive depth regime that estimated depth bins between 30 and 400 m; the latter value was derived from Shaff and Baird (2021) to be the maximum depth for rough-toothed dolphin deep dives. Tag positional update times, or times when tag position updates were attempted but failed, can be used as known times the animals are at the surface (see Dive Analysis section); these provide additional surface information for all tagged delphinids for periods when dive data were missing. When the satellite tags provided depth information, the modeled animal depth data corresponding to the time an MFAS transmission was received at the animal position was utilized, along with a percent of depth to represent uncertainties in depth. The percent of depth uncertainty used in this analysis was varied with the depth regime: for shallow depths to 54 m, each 9 m depth bin from the surface to 54 m depth were utilized; for depths from 54 m to 100 m, +/- 20% of the depth were utilized; for depths from 100 m to 400 m +/- 10% were used; and for depths > 400 m +/- 5% were used. Figure 3 shows a small red box indicating the estimated animal location for the propagation modeled single slice. Multiple slices estimated animal locations contribute to the full 3D estimate for each exposure.

Figure 4 illustrates the concept of utilizing radial azimuthal slices in a 3D manner. Each radial slice contribution to the 3D volume is considered. Given the acoustic propagation model computes the full field, this allows us to utilize all available elements of the estimated RL (three dimensions plus time along the track). Figure 5 illustrates a histogram of all available (19,872 in this example) individual depth-range elements within an animal's estimated 3D location at the time of an MFAS transmission. The

distribution is not nicely Gaussian in nature; although there is an estimated median received level of 129.3 dB, the histogram demonstrates the relatively high uncertainty in the actual exposure value for this MFAS transmission. This is also evident in Figure 3 where it is evident how multipath arrivals can contribute stronger estimated received levels while much of the ensonified area has relatively lower estimated received levels. Therefore, the estimated received level needs to be defined by a single metric and accompanied by a metric for the variation of the received level. Following Henderson et al. (2021) we will provide the median RL +/- twice the standard deviation. Furthermore, while the number of MFAS transmissions in a 5 min bin is sensitive, we are providing a stoplight (red, yellow and green) colorization of the median RL in plots to indicate relative MFAS activity in the bin with green being low, yellow moderate, and red high.




Figure 4 – 3D concept utilizing multiple radials from MFAS source through an animal’s estimated location in 3D space, using the red box insert from Figure 3.





Figure 5 – Example of all estimated receive levels for a single MFAS transmission via 25 slices from the MFAS ship to one estimated possible 95% CI location in three dimensions.

There are also diagnostic metrics available to account for the loss of propagation modeled received level elements due to a variety of situations, such as some of the possible animal locations being impossible. This could be for a variety of reasons; 1) a portion of the animal's error ellipse is shadowed from the MFAS source due to bathymetry or a land mass; 2) If the estimated water depth is less than the animal's modeled foraging dive depth; or 3) if the animal's error ellipse is within a few hundred meters of the source, the steep angle limitation precludes providing estimates. A single metric, termed the Probability of Exposure, is 1.0 for exposures with all of the 3D elements being represented in the propagation model; when this metric is  $< 1.0$  it flags exposures that might not have been fully possible (e.g., a majority of the animal's error ellipse is shadowed from the MFAS source by, for example, the islands). Figure 6 illustrates an example of this for one position from GmTag235, where a portion of the error ellipse has an acoustic path to a ship (in red), while the remainder of the ellipse is shadowed by land (in blue).



*Figure 6 - Plan view of exposure geometry for one location illustrating the situation where the animal may, or may not, have been exposed to MFAS depending on where it actually was within its 95% confidence interval location uncertainty. To account for this, an estimated probability of exposure metric is included for all estimated exposures to MFAS.*

### Ship Exposures

Each surface ship hull-mounted MFAS localization was joined to ship positional data from PMRF, nominally updated every second, if data were within one second and 400 m. During a 5-minute interval, one transmission from each individual ship transmitting sonar and its azimuthal radials were selected for propagation modeling if it was closest in time (within  $\pm 2.5$  mins) and distance to a whale update. A similar approach was utilized by Henderson et al. (2021) with the exception that in that analysis a single transmission with the absolute minimum time and distance to a whale update was selected for propagation modeling. The approach taken here requires more processing time for additional radials but



provides a continuous exposure history for each ship. This method also allows for the interpolation of all pings that occurred between the propagation-modeled exposures for each ship participating in the training activity.

### Behavioral Response and Diel Analysis

Timing and general locations of tagged animals were first assessed in relation to the timing and general location of both Phase A and Phase B, to determine whether additional assessment of potential behavioral responses were warranted. If tags ceased transmitting prior to the start of Phase A, if tagged individuals were in the acoustic shadow of Kaua'i or Ni'ihau during Phase A and B, or if tagged individuals had moved >100 km from the general area of the SCC they were not considered in the behavioral analyses. For each tag with dive behavior, the coverage of dive and surfacing data during each phase was first evaluated to provide an indication on the robustness of comparisons among phases. To do this, the duration of all dive and surfacing periods within each phase were summed up, then divided by the total durations of the respective phases. Any dive or surface periods that spanned more than one phase had their durations split (e.g., such that a surfacing period beginning in Phase A and continuing into the Interphase would have its duration split between the two periods based on when the Interphase begins). Because not all tags transmitted for the full duration of each Phase, coverage relative to the duration of each tag was evaluated. To do this, the duration of all dive and surfacing periods within each phase were summed up, with those that occurred over multiple phases split as before, and divided by the total duration of tag transmission during each phase. This allows for an assessment of gaps in the behavioral data that might influence the likelihood of detecting potential behavioral responses.

To account for the impacts of diel patterns on diving behavior, the coverage of each tag based on time of day for each SCC Phase was also calculated. Dive and surface periods were each assigned a time of day based on when they started, defined as either dawn, day, dusk, or night. Dawn was defined as the period before and after sunset, with solar angles between 6° below and above the horizon. Day was defined as the period after sunrise and prior to sunset with solar angles > 6° above the horizon. Dusk was defined as the period prior to sunset with solar angles between 6° below and above the horizon. Night was defined as the period after sunset and prior to sunrise with solar angles >6° below the horizon. The durations of surface periods that spanned more than one time of day were split (e.g., such that a surface period beginning at dawn and continuing into day would have its duration split between the two times of day based on when day begins). As before, surface periods that crossed multiple SCC phases had their durations split between phases. Due to their short duration, no dives were split based on either phase or time of day. Coverage by time of day for each tag was calculated as the total duration of dive and surfacing periods within the time of day and SCC phase of interest, divided by the total duration of the time of day within that particular phase (e.g., the dawn total duration for Phase A would represent the sum of the duration of all dawns within Phase A). For the Before and After periods, the total duration of the phase was calculated as 3 days prior to the start of Phase A, and 3 days following the end of Phase B, respectively.

Metrics calculated included the dive rate (number of dives per hour), percentage of time spent at the surface, median dive depth, and median dive duration among SCC phases and times of day for each tag to assess potential responses to MFAS exposure in their diving behavior while also accounting for known

diel patterns (see Owen et al. 2019, Shaff & Baird 2021, West et al. 2018). For all metrics, only 3 days of data following the end of Phase B (i.e., After) were used where available.

Kruskal-Wallis one-way ANOVA tests were conducted to identify significant differences in dive depth and duration among phases, and by night/day period, for each tagged individual with sufficient dive/surfacing coverage (SPLASH10 tags only), and post-hoc Dunn’s tests with a Benjamini-Hochberg correction were conducted to identify phases where pairwise significant differences were detected (e.g., statistical difference between phase A and B; significance level for both tests = 0.05). These statistical procedures were not applied to summaries on dive rates (dives per hour) nor percentage of time at surface due to the nature of how these values were calculated (i.e., only single values for each SCC phase). Sufficient coverage for each phase was defined based on species and time of day to account for species-level variation in diving behavior, with those species that have the longest dive durations requiring a higher level of coverage during the dawn and dusk periods for inclusion in statistical tests (Table 2). Additionally, due to their relatively short durations, the dawn and dusk times of day had higher coverage requirements to ensure that data from these times of day were providing an accurate representation of diving behavior.

*Table 2 - Coverage requirements for each phase by time of day and species for inclusion in statistical comparisons between phases. Gm = Pilot whale (Globicephala macrorhynchus), Pe = Melon-headed whale (Peponocephala electra), Pc = False killer whale (Pseudorca crassidens), Tt = Common bottlenose dolphin (Tursiops truncatus), Sb = Rough-toothed dolphin (Steno bredanensis).*

Species	Dawn % coverage required for inclusion in statistical testing	Day % coverage required for inclusion in statistical testing	Dusk % coverage required for inclusion in statistical testing	Night % coverage required for inclusion in statistical testing	Notes
Gm	80	50	80	50	Due to their longer dive durations, the dawn and dusk coverage requirements for this species are set at 80%.
Pe	NA	NA	NA	50	As this species dives almost exclusively at night (West et al. 2018), only night is statistically tested for this species.
Pc, Tt, Sb	70	50	70	50	

## RESULTS

### Tagging and Photo-Identification

*A summary of individuals that were tagged, including their tag duration, overlap with SCC phases, and maximum estimated RL, are given in*

Table 3. The false killer whale tagged in 2021 (PcTag074) was identified as HIPc364, a member of Cluster 3 (cf. Mahaffy et al. 2023) of the endangered main Hawaiian Islands population. This individual had previously been documented off Hawai'i Island (2009, 2014) and O'ahu (2017, 2021), but had not been previously documented off Kaua'i. All six of the tagged short-finned pilot whales from 2021 and 2022 had been previously documented. The three individuals tagged in 2021 were from two different social clusters (see Mahaffy et al. 2015). GmTag232 and GmTag233 (HIGm0242 and HIGm1166, respectively) are members of Cluster W11, known to be part of the western community (cf. Baird 2016; Van Cise et al. 2017). These individuals have been previously documented off Kaua'i in 2005 (HIGm0242) and 2008 (both individuals), and off O'ahu in 2009 (HIGm1166). GmTag234 was identified as HIGm0205 and a member of Cluster W8, previously documented off O'ahu in 2003 and 2009 and off Kaua'i in 2012. All three individuals tagged in 2022 (GmTag235, GmTag236, and GmTag237, individuals HIGm2222, HIGm2213, and HIGm0768, respectively) are members of Cluster H14. This group has not been linked by association with any other group and thus its community membership is unknown. Sighting history of the tagged individuals (HIGm0768 off Hawai'i Island in 2006 and 2013, HIGm2213 off Hawai'i Island in 2013 and O'ahu in 2016, HIGm2222 off Hawai'i Island in 2013 and 2014) suggest they are not part of the resident communities and are likely an offshore group. None of these individuals had been previously tagged.

Two of the three tagged bottlenose dolphins (TtTag040 and TtTag041) had previously been documented off Kaua'i, while the other (TtTag039) had not been previously identified. While TtTag039 had not been previously documented, it was linked by association with the resident, island-associated population through other individuals in the encounter. TtTag040 was identified as HITt0740 in the photo-identification catalog, and had been photographed on six different occasions in three different years (first in 2011). TtTag041 was identified as HITt1474 in the photo-identification catalog, and had been photographed once previously (in 2018). Both HITt0740 and HITt1474 are known to be part of the island-associated resident community (Baird et al. 2009). Only one of the three tagged rough-toothed dolphins had been previously documented. SbTag024 was identified as HISb1474 and been documented on six different occasions in four different years (first in 2008), and is known to be part of the island-associated resident population (Baird et al. 2008). The other tagged rough-toothed dolphin from 2021 (SbTag023, HISb2743 in the photo-identification catalog) was in the same encounter as HISb1474, and thus is linked by association with the resident, island-associated population. The individual tagged in 2022 (SbTag025, HISb2744 in the catalog) had not been encountered previously, but at least one other distinctive individual in the encounter was known to be part of the resident, island-associated community, and thus this individual was considered a member by association. None of the individuals of either species had been previously tagged. Although Cascadia has a melon-headed whale photo-identification catalog (Aschettino et al. 2012), the catalog has not been substantially updated since 2011. Given the size of the catalog (~2000 individuals) and the rate of mark change, it is unlikely that matches to tagged individuals would be found without also updating the catalog with photos available in the intervening years. Finally, three Blainville's beaked whales were also tagged, two in 2021 and one in 2022, but their results are discussed in a separate paper (Henderson et al. 2024) and will not be mentioned further in this report.

Table 3 - Tag Deployment data for 2021 and 2022 satellite tags. An “NA” in the maximum received level column indicates that there was no overlap between the specified tag and MFAS blocks. Max RL = Maximum estimated received level.

Tag ID	Tag type	Deployment Date/time (GMT)	End Date/time (GMT)	Duration (days)	Overlap with SCC Phases	Max RL (dB re 1 µPa)
GmTag232	SPLASH10-F	2021-08-08 21:32	2021-08-22 21:11	14.0	A, B	80.4
GmTag233	SPLASH10-F	2021-08-10 18:36	2021-08-26 21:30	16.1	A, B	81.2
GmTag234	SPLASH10-F	2021-08-10 19:01	2021-09-03 20:23	24.1	A, B	72.6
PcTag074	SPLASH10-F	2021-08-08 16:35	2021-08-20 17:42	12.0	A, B	90.5
PeTag029	SPLASH10-F	2021-08-01 17:56	2021-08-14 08:11	12.6	A	147.7
PeTag030*	SPLASH10-F	2021-08-09 18:59	2021-08-09 18:59	0	NA	NA
PeTag031	SPLASH10-F	2021-08-11 19:21	2021-09-01 09:04	20.6	A, B	136.6
PeTag032	SPLASH10-F	2021-08-13 19:24	2021-08-30 05:31	16.4	B	135.5
SbTag023	SPLASH10	2021-08-09 20:23	2021-08-22 20:12	13.0	A, B	141.9
SbTag024	SPOT6	2021-08-09 20:54	2021-08-14 09:19	4.5	A	NA
TtTag039	SPLASH10-F	2021-08-12 21:00	2021-08-25 16:10	12.8	A, B	100.0
TtTag040	SPLASH10-F	2021-08-14 19:44	2021-08-30 16:30	15.9	B	100.5
SbTag025	SPLASH10	2022-08-24 20:17	2022-09-01 08:30	7.5	B/A Mixed	NA
GmTag235	SPLASH10-F	2022-08-17 17:43	2022-09-03 21:09	17.1	A/B Mixed, B/A Mixed	129.2
GmTag236	SPLASH10-F	2022-08-17 20:00	2022-08-21 21:16	4.1	A/B Mixed	128.8
GmTag237	SPLASH10-F	2022-08-17 20:27	2022-10-02 08:42	45.5	A/B Mixed, B/A Mixed	128.8
PeTag033	SPLASH10-F	2022-08-18 18:20	2022-08-29 21:24	11.1	A/B Mixed, B/A Mixed	70.1
PeTag034	SPLASH10-F	2022-08-18 19:00	2022-08-22 21:57	4.1	A/B Mixed	72.6
PeTag035	SPLASH10-F	2022-08-24 19:24	2022-09-12 03:05	18.3	B/A Mixed	NA
PeTag036	SPLASH10-F	2022-08-24 19:49	2022-09-04 05:03	10.4	B/A Mixed	NA
TtTag041	SPLASH10	2022-08-21 20:36	2022-09-07 19:49	17.0	B/A Mixed	NA

\*Tag failed upon deployment

*In 2021 there was a two-day Unit Level Test (ULT) prior to the SCC, and then the SCC was conducted in a typical fashion, with Phase A (no surface ship or MFAS activity) occurring for three days, then a three-day break, then Phase B (which includes surface ships and MFAS) occurring for another four days. A three-day before and a three-day after period are also included in*

*Table 4 for reference. In 2022 there was a one-day Submarine Exercise (SUBEX) prior to the SCC. In this case, the SCC was conducted slightly differently than is typical, with components of Phase A and B occurring in both weeks. The first four-day portion is referred to in this report as A/B mixed, indicating it takes place during the typical Phase A period, while the second period only occurred for one day and is referred to in this report as B/A mixed, indicating it was conducted during the typical Phase B period. There was also the usual 5-day break in between the two training phases, and a 3-day before and 3-day after period included in*

Table 4 as well.

Table 4 - SCC Phase times and passive acoustic monitoring data durations (times given in HST)

Phase	Start date	End date	Duration (hrs)
Pre-ULT	7/31/2021 11:00	8/3/2021 10:57	72
ULT	8/3/2021 11:27	8/3/2021 16:42	5.3
Post-ULT/Pre-SCC	8/3/2021 16:45	8/11/2021 17:59	193.2
Phase A	8/11/2021 18:00	8/13/2021 22:20	40.3
Interphase	8/13/2021 10:21	8/17/2021 4:59	90.6
Phase B	8/17/2021 5:00	8/19/2021 14:30	57.5
Post-SCC	8/19/2021 14:31	8/22/2021 14:30	72
Pre-SUBEX	8/13/2022 2:30	8/16/2022 2:29	72
SUBEX	8/16/2022 6:01	8/16/2022 16:00	10.0
SCC A/B (mixed)	8/17/2022 6:30	8/20/2022 23:31	89.0
Interphase	8/20/2022 23:32	8/24/2022 10:29	73.3
SCC B/A (mixed)	8/24/2022 10:30	8/24/2022 15:53	5.4
Post-SCC	8/24/2022 15:54	8/27/2022 15:54	72

### Behavioral Response, Diel Analysis, and Received Level Estimation

The following sections document data available for behavior analyses, as well as movements and estimated RLs for short-finned pilot whales, false killer whales, melon-headed whales, rough-toothed dolphins, and bottlenose dolphins across two separate SCC training events, following the format from Henderson et al. (2021).

Of the 21 animals tagged at PMRF in 2021 and 2022, 16 animals remained in the area during the phases of the SCCs during which MFAS is used (Phase B in 2021, Phases A/B and B/A in 2022). Only the hull-mounted MFAS was modeled for received level estimation in this report, however in 2021 there was also helicopter-dipping and active sonobuoy MFAS present, and in 2022 there was helicopter-dipping MFAS present. The source levels of the latter two sources are much lower than that of hull-mounted MFAS and they are typically deployed in the same area of the range as the hull-mounted MFAS activity and so are not expected to be large contributors to the MFAS soundscape experienced by these whales. For all of these animals, the median estimated RL per 5 min bin (associated with a *crawl*-smoothed track location) are reported along with the  $\pm 2$  standard deviation (SD) values for those bins. As a reminder, wide  $\pm 2$  SD on the RL figures indicate a larger error ellipse on the *crawl*-smoothed track location (e.g., on the 4<sup>th</sup> to last ping on Figure 7). Bouts or blocks of MFAS are considered to be periods of MFAS separated by at least 30 min, and there are multiple blocks per Phase (Phase B or B/A). Širović et al. (2013) determined that ambient noise levels are about 60 dB re 1  $\mu$ Pa at 1 kHz; noise levels would be even lower in the mid-frequency band of 1-10 kHz. Using this value as the noise floor, and following what has been done previously at PMRF (Baird et al. 2019) and the recommendation of Schick et al.

(2019), any estimated received levels below 60 dB re 1  $\mu$ Pa will be included in figures but will otherwise be reported as “NA”.

Summary tables and figures are only included for individuals with exposures and statistically significant differences among phases; the remaining tables and figures are included in the Appendix. Sections are organized by species, and within species in the following order: summary tables, followed by narratives for each individual or group including estimated RLs, maps for those specific animals, dive behavior narratives for each individual or group, and applicable dive behavior figures.

#### Short-finned pilot whales

Six short-finned pilot whales were tagged, three in 2021 in one group, and three in 2022 in another group. Summary statistics are presented below in Table 5, Table 6, and Table 7 for all animals.

*Table 5 - Percentage of dive/surfacing data by phase for short-finned pilot whales. The percentage of behavioral coverage is defined as the proportion of the duration of behavioral data relative to the duration of the tag within each phase.*

Individual	Percentage of dive/surfacing data				
	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After
<b>GmTag232</b>					
Duration overall (days)	3.3	1.7	3.8	2.4	2.9
Days surfacing/dive data	3.3	1.7	3.8	2.3	1.7
Percentage behavioral coverage	100.0	100.0	99.2	95.0	58.4
<b>GmTag233</b>					
Duration overall (days)	1.4	1.7	3.8	2.4	6.9
Days surfacing/dive data	1.4	1.4	3.1	2.3	3.2
Percentage behavioral coverage	100.0	80.4	81.5	95.8	46.1
<b>GmTag234</b>					
Duration overall (days)	1.4	1.7	3.8	2.4	14.8
Days surfacing/dive data	1.2	1.5	3.5	2.4	3.2
Percentage behavioral coverage	89.8	88.7	93.7	99.6	21.4
<b>GmTag235</b>					
Duration overall (days)	NA	3.7	3.5	0.2	9.8
Days surfacing/dive data	NA	3.6	3.3	0.2	4.8
Percentage behavioral coverage	NA	97.5	95.1	100.0	49.4
<b>GmTag236</b>					
Duration overall (days)	NA	3.6	0.5	0.0	0.0
Days surfacing/dive data	NA	2.8	0.0	0.0	0.0
Percentage behavioral coverage	NA	78.4	8.2	0.0	0.0
<b>GmTag237</b>					
Duration overall (days)	NA	3.5	3.5	0.2	38.3
Days surfacing/dive data	NA	3.2	3.4	0.2	5.0
Percentage behavioral coverage	NA	91.2	98.0	100.0	13.1



Table 6 - A comparison of dawn diving parameters from short-finned pilot whales exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

Dive parameter per individual	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Dawn dive rate (dives/hour)</i>							
GmTag232	1.99	1.12	2.15	2.31	NA	-	
GmTag233	NA	NA	0.56	2.33	1.90	-	
GmTag234	NA	2.25	2.32	2.14	1.50	-	
GmTag235	NA	NA	2.60	NA	1.27	-	
GmTag237	NA	NA	2.70	NA	1.71	-	
<i>% time in surface periods at dawn</i>							
GmTag232	39.67	72.19	37.17	43.64	NA	-	
GmTag233	NA	NA	93.07	52.78	54.62	-	
GmTag234	NA	38.33	43.35	41.24	53.42	-	
GmTag235	NA	NA	33.90	NA	61.22	-	
GmTag237	NA	NA	23.23	NA	45.47	-	
<i>Median dive depth dawn (m)</i>							
GmTag232	751.50	663.50	655.50	623.50	NA	0.6202	NA
GmTag233	NA	NA	99.50	647.50	623.50	0.1576	NA
GmTag234	NA	903.50	719.50	655.50	599.50	0.1942	NA
GmTag235	NA	NA	679.50	NA	735.50	0.1474	NA
GmTag237	NA	NA	719.50	NA	687.50	0.8638	NA
<i>Median dive duration dawn (min)</i>							
GmTag232	18.97	14.95	16.90	14.60	NA	0.1195	NA
<b>GmTag233</b>	<b>NA</b>	<b>NA</b>	<b>7.47</b>	<b>13.10</b>	<b>14.27</b>	<b>0.0147</b>	<b>B-After</b>
GmTag234	NA	16.70	17.28	16.20	18.48	0.1596	NA
<b>GmTag235</b>	<b>NA</b>	<b>NA</b>	<b>14.95</b>	<b>NA</b>	<b>18.43</b>	<b>0.0278</b>	<b>Inter-After</b>
<b>GmTag237</b>	<b>NA</b>	<b>NA</b>	<b>17.63</b>	<b>NA</b>	<b>19.03</b>	<b>0.0202</b>	<b>Inter-After</b>



Table 7 - A comparison of nighttime diving parameters from short-finned pilot whales exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

Dive parameter per individual	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Night dive rate (dives/hour)</i>							
GmTag232	2.97	2.72	2.71	2.51	2.42	-	
GmTag233	NA	4.67	3.80	3.70	2.78	-	
GmTag234	NA	2.75	2.91	3.11	2.59	-	
GmTag235	NA	3.63	3.47	NA	3.02	-	
GmTag236	NA	3.53	NA	NA	NA	-	
GmTag237	NA	3.56	3.25	NA	3.20	-	
<i>% time in surface periods at night</i>							
GmTag232	34.82	37.18	29.86	37.48	38.31	-	
GmTag233	NA	29.11	33.85	33.35	39.83	-	
GmTag234	NA	32.35	33.73	36.71	36.25	-	
GmTag235	NA	26.55	28.98	NA	31.84	-	
GmTag236	NA	25.57	NA	NA	NA	-	
GmTag237	NA	23.56	30.96	NA	27.27	-	
<i>Median dive depth night (m)</i>							
<b>GmTag232</b>	<b>403.50</b>	<b>504.50</b>	<b>527.50</b>	<b>591.50</b>	<b>543.50</b>	<b>NA</b>	<b>Inter- Before; B-Before; After- Before</b>
<b>GmTag233</b>	<b>NA</b>	<b>227.50</b>	<b>233.50</b>	<b>255.50</b>	<b>559.50</b>	<b>&lt;0.0001</b>	<b>Inter- After; A- After; B- After</b>
<b>GmTag234</b>	<b>NA</b>	<b>575.50</b>	<b>543.50</b>	<b>511.50</b>	<b>591.50</b>	<b>0.0040</b>	<b>Inter-A; A-B; Inter- After; B- After</b>
GmTag235	NA	323.50	387.50	NA	391.50	0.4695	NA
GmTag236	NA	319.50	NA	NA	NA	NA	NA

Dive parameter per individual	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After	Kruskal- Wallis Test p- value*	Post-hoc Dunn's test significant pairs
GmTag237	NA	271.50	295.50	NA	335.50	0.8859	NA
<i>Median dive duration night (min)</i>							
<b>GmTag232</b>	<b>14.12</b>	<b>13.07</b>	<b>15.13</b>	<b>15.17</b>	<b>15.67</b>	<b>0.0188</b>	<b>No significant adjusted p-values</b>
<b>GmTag233</b>	<b>NA</b>	<b>8.93</b>	<b>10.32</b>	<b>10.97</b>	<b>13.53</b>	<b>&lt;0.0001</b>	<b>Inter-A; A-B; Inter-After; A-After; B-After</b>
<b>GmTag234</b>	<b>NA</b>	<b>14.67</b>	<b>13.97</b>	<b>12.42</b>	<b>14.87</b>	<b>0.0002</b>	<b>Inter-B; A-B; Inter-After; B-After</b>
<b>GmTag235</b>	<b>NA</b>	<b>12.08</b>	<b>12.50</b>	<b>NA</b>	<b>13.93</b>	<b>0.0003</b>	<b>Inter-After; A-After</b>
GmTag236	NA	12.82	NA	NA	NA	NA	NA
GmTag237	NA	12.60	12.57	NA	14.20	0.0641	NA

*GmTag232, GmTag233, GmTag234*

These three individuals were tagged within the same group in 2021, and generally remained associated over the overlapping period of tag attachment. Information was available on movement patterns for Before (1.4–3.3 days), Phase A (1.7 days), the interphase period (3.8 days), Phase B (2.4 days), and After (2.9–14.8 days; Table 5). All three of these individuals remained off the range for the entire duration of the SCC, though they remained in close proximity to Kaua'i.

After tag deployment south of the range, all three individuals moved along the southern coast of Kaua'i and remained in this general area throughout Phase A. After Phase A, they remained in the area for the few days of the interphase period, and then moved along the eastern coast of Kaua'i before returning to the southern coast at the start of Phase B. These individuals remained in this same area during Phase B, and were largely out of range of MFAS exposure throughout Phase B, except for a handful of exposures ranging between 47.0 and 81.2 dB re 1  $\mu$ Pa. After Phase B, these individuals remained in the same general area for a short period of time, and then moved back up along the east coast of Kaua'i. Because these individuals remained associated during their shared deployment period and had highly similar exposure levels, only one individual was mapped here (Figure 10).

GmTag232 was exposed to three different bouts of hull-mounted MFAS, although as described they were off the range and behind the island of Kaua’i for most of the SCC. They did have a few exposure bins with a probability of 1, but others were as low as 0.2 (e.g., only a portion of the error ellipse for that track location could have been exposed). In addition, distances were all longer than 72 km and all median RL values were less than 82 dB re 1  $\mu$ Pa (Table 8, Figure 7). In fact, 9 of the 14 track locations that had some MFAS exposure had RLs below the ambient noise floor of 60 dB re 1  $\mu$ Pa.

Table 8 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag232.

Minimum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	NA
Maximum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	80.4 (67.5, 93.4)
CPA of ship (km)	72.2
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	85.2




Figure 7 - Median RLs for GmTag232 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.

GmTag233 had similarly low estimated RLs and long distances for their two exposure periods (

Table 9, Figure 8). Six of the 11 locations with exposures had RLs below ambient noise levels, and 8 of the 11 locations had less than a probability of 1 of exposure.

Table 9 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag233.

Minimum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	NA
Maximum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	80.4 (69.6, 92.9)
CPA of ship (km)	72.9
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	87.2




Figure 8 - Median RLs for GmTag233 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.

GmTag234 had similar low RLs and reduced probabilities of exposure (Table 10, Figure 9). Again, of the 11 track locations with exposures, eight of those had a less than 1 probability of actually getting exposed, and seven were at or below the 60 dB re 1  $\mu$ Pa ambient noise floor.

Table 10 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag234.

Minimum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	NA
Maximum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	72.6 (59.6, 85.7)
CPA of ship (km)	72.1
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	80.2




Figure 9 - Median RLs for GmTag234 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.

## GmTag233 - August 2021




Figure 10 - Movements of GmTag233 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level, and time is given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygon represents the area of ship activity during each of the three MFAS bouts that GmTag233 was exposed to and the corresponding diamond point represents the mean ship location during the bouts. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary.

### Dive Behavior

GmTag232, GmTag233, and GmTag234 all transmitted dive behavior data for each phase (Table 6 and Table 7; and Figure 11, Figure 12, and Figure 13). However, when broken down by time of day and phase, not every time of day and phase met the required coverage (relative to the phase duration) for inclusion in the analysis.

Dawn dive rates for the three tags ranged from 0.56 dives/hr (GmTag233, Interphase) to 2.33 dives/hr (GmTag233, Phase B), with varied trends following exposure phases. Compared to Before, GmTag232 had a decreased dawn dive rate during Phase A, but subsequently increased its dive rate across the next two phases. The dawn dive rate for GmTag233 also increased during Phase B. In contrast, dawn dive rates for GmTag234 remained relatively consistent between Phase A and the Interphase, and began to decrease during Phase B. Following Phase B dawn dive rates decreased for both GmTag233 and

GmTag234. The percentage of surface time almost doubled during Phase A compared to Before (39.67% to 72.19%) for GmTag232, and decreased again during the interphase and Phase B. For GmTag233, the percentage of surface time was highest during the interphase period, after which it decreased to more consistent levels during Phase B and After. In contrast, GmTag234 maintained relatively consistent surface time percentages across phases, ranging from a minimum of 38.33% (Phase A) to a maximum of 53.42% (After). Variation in dawn dive depths between phases were not statistically significant for any tag, but there were significantly longer dives during After when compared to Phase B for GmTag233.

Day dive rates for the three tags were generally lower than dawn dive rates, and ranged from 0.35 dives/hr (GmTag233, After) to 1.89 dives/hr (GmTag234, Phase A), with varied trends following exposure phases. Compared to Before, GmTag232 had a sharply decreased day dive rate during Phase A, while GmTag233 remained relatively consistent between the two. Following Phase A, day dive rates remained fairly consistent for all tagged individuals, but decreased following Phase B for GmTag233 and GmTag234. The percentage of surface time during the day remained fairly consistent across phases for all three tags, though it did increase moderately following Phase B for GmTag233 and GmTag234. Day dive depths had statistically significant variation between phases for all tags, though only GmTag233 and GmTag234 had statistically significant pairwise post-hoc comparisons between phases. For GmTag233, there were significantly shallower dives during After when compared to Before, while for GmTag234 there were significantly deeper dives during After compared to Phase A and the interphase, and significantly deeper dives during Phase B compared to the interphase. Day dive durations also had statistically significant variation between phases for GmTag232 and GmTag234, with GmTag232 having significantly shorter dives during Before compared to Phase A and interphase, while GmTag234 had significantly shorter dives during the interphase compared to Phase B and After.

Dusk dive rates were generally higher than dawn or daytime dive rates, and ranged from 2.29 dives/hr (GmTag234, interphase) to 3.26 dives/hr (GmTag233, After), with varied trends following exposure phases. Compared to Before, GmTag232 had a decreased dusk dive rate during Phase A, and the dive rate for this tag continued to decrease during the interphase. During Phase B, dusk dive rates increased for GmTag232 and GmTag234, but fell for GmTag233. Following Phase B, GmTag233 had an increase in dusk dive rate, while GmTag234 had only a very slight decrease in dive rate. The percentage of surface time for all three tags remained comparatively low during the dusk hours, and was generally consistent across most phases, with the exception of a moderate increase in surface time for GmTag233 during Phase B (43.46%) when compared to the interphase (31.72%) and After (25%). Variation in dusk dive depths and durations between phases was not statistically significant for any tag.

Night dive rates were generally the highest of any time of day, and ranged from 2.42 dives/hr (GmTag232, After) to 4.67 dives/hr (GmTag233, Phase A), with varied trends following exposure phases. Compared to Before, GmTag232 had only a slight decrease in night dive, and the dive rate for this individual did not vary substantially from Phase A to the interphase. Following Phase A, GmTag233 had a decrease in dive rate that continued throughout the remaining duration of the SCC, while GmTag234 had a slight increase in dive rate that continued into Phase B. Following Phase B, all individuals had decreases in night dive rates, though there was considerable variation in the extent of the decrease. The percentage of surface time at night for all three tags remained largely consistent across phases, but there was significant variation in night dive depths. Night dive depths were significantly deeper during the interphase, Phase B, and After when compared to Before for GmTag232. Dive depths were also significantly deeper during After when compared to Phase A, the interphase, and Phase B for

GmTag233, and when compared to the interphase and Phase B for GmTag234. In contrast, night dive depths were significantly shallower during Phase B when compared to Phase A and the interphase for GmTag234. Night dive durations also varied significantly between phases for all tags, though only GmTag233 and GmTag234 had statistically significant post-hoc pairwise comparisons between phases. For GmTag233, Phase A had significantly shorter dive durations compared to all following phases, and After had significantly longer dive durations compared to all previous phases. For GmTag234, dive durations were significantly shorter during Phase B when compared to previous phases, and significantly longer during After when compared to the interphase and Phase B.






Figure 11 - Top. Boxplot showing dive depths of GmTag232 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag232 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 80.4 dB re 1  $\mu$ Pa.




Figure 12 - Top. Boxplot showing dive depths of GmTag233 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag233 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 81.2 dB re 1  $\mu$ Pa.




Figure 13 - Top. Boxplot showing dive depths of GmTag234 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag234 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 72.6 dB re 1  $\mu$ Pa.

*GmTag235, GmTag236, GmTag237*

These three individuals were tagged within the same group in 2022, and generally remained associated over the overlapping period of tag attachment, with received levels during exposures varying only slightly between individuals. Information was available on movement patterns for GmTag235 and GmTag237 for the A/B mixed phase (3.5 and 3.7 days, respectively), the interphase period (3.5 days), the

B/A mixed phase (0.2 days), and the after phase (9.8 and 38.3 days, respectively; Table 5). For GmTag236, movement data was available only for the A/B mixed phase (3.6 days) and the interphase period (0.5 days). All three tagged individuals were initially present on the range during the A/B mixed phase then began to move southeast, with GmTag236 ceasing transmission halfway between Kaua'i and O'ahu, and GmTag235 and GmTag237 continuing to transmit until they reached Maui Nui.

While all three tags were deployed on the range, prior to the start of exposures during the A/B mixed phase the three individuals had begun moving towards the southern boundary of the range. Following the initial exposure, the animals continued to move southward, and had left the range by the time the third exposure began, though received levels generally remained above 120 dB re 1  $\mu$ Pa during all three of these exposures. During the initial exposure, the three tags had median received levels over 100.0 dB re 1  $\mu$ Pa over a period of more than two hours, with the highest median received level reaching 125.0 dB re 1  $\mu$ Pa (for GmTag235). During the second exposure, the three tags had median received levels ranging from 115.6 to 127.6 dB re 1  $\mu$ Pa over a period of a half hour, and during the third (and longest) exposure, the three tags had median received levels ranging from 88.4 to 129.2 dB re 1  $\mu$ Pa over a period of four hours. By the time of the fourth exposure, they had moved far away enough from the range that the highest median received levels had dropped substantially, and were below the ambient noise floor. Because all three individuals remained closely associated during the periods of interest and had highly similar exposure levels, only one individual was mapped here (Figure 17).

GmTag235 was exposed to six bouts of MFAS, with the first five being on or close to the range and the last once they had moved out of the area. Of the 47 track locations with exposures, the first 40 occurred in the initial 5 bouts of MFAS and had probabilities of exposure of 1, while the remaining 7 locations occurred in the last bout of MFAS and had much lower probabilities of exposures and very low received levels, well below the ambient noise floor (Table 11, Figure 14). The estimated received levels from the first 5 bouts of MFAS were at low to moderate levels (95 to 129 dB re 1  $\mu$ Pa), but some exposure locations also had a higher level of pings per 5 min bin.

*Table 11 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag235.*

Minimum median RL value ( $\pm$ 2 SD) dB re 1 $\mu$ Pa	NA
Maximum median RL value ( $\pm$ 2 SD) dB re 1 $\mu$ Pa	129.2 (124.3, 134.1)
CPA of ship (km)	48.2
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	135.4




Figure 14 - Median RLs for GmTag235 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.

GmTag236 similarly started on the range and then moved southwest around Kaua’i and out of the area. Once again this animal was exposed to six bouts of MFAS, with the first five occurring while the animal was still on or near the range and the last bout occurring after the animal was well out of the area (Table 12, Figure 15). Therefore, the probabilities of exposure were all 1 for the first 5 bouts and very low for the last bout, with some 5 min bins having a high number of pings per bin, and again the last bout had RLs well under the ambient noise floor. RLs in the first five MFAS bouts ranged from 88 to 129 dB re 1  $\mu$ Pa.

Table 12 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag234.

Minimum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	NA
Maximum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	128.8 (124.5, 133.0)
CPA of ship (km)	48.9
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	135.0




Figure 15 - Median RLs for GmTag236 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.

GmTag237 also experienced six bouts of MFAS, with the last occurring after they had left the area and so had low probabilities of exposures and low RLs. Their median RLs while on the range were comparable to the other members of their group, ranging from 89 to 129 dB re 1  $\mu$ Pa (Table 13, Figure 16).

Table 13 - Estimated received levels, cumulative sound exposure levels, and ship CPA for GmTag237.

Minimum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	NA
Maximum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	128.8 (124.5, 133.0)
CPA of ship (km)	47.7
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	134.0




Figure 16 - Median RLs for GmTag237 in stoplight colors (green for few pings during 5 min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.




Figure 17 - Top: Movements of GmTag237 during the August 2022 SCC event, including extensive movements away from the range after exposure phases. Bottom: zoomed-in panel of the track to highlight movements that occurred before, during, and after the exposure, prior to the animal's extensive movements away from the area. Note that for this particular event, ship-based MFAS activity occurred during both A and B phases. The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level, and times given in GMT. Additionally, the RL circles are colored by "intensity" which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the MFAS bouts that GmTag237 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary.



## Dive Behavior

GmTag235 and GmTag237 both transmitted dive behavior for each phase, except for the before phase. GmTag236 only transmitted dive behavior for the A/B mixed phase, and is given no further consideration (Figure 18, Figure 19). However, when broken down by time of day and phase, not every time of day and phase met the required coverage (relative to the phase duration) for inclusion in the analysis.

Dawn dive metrics could only be calculated for the interphase and B/A mixed phase for GmTag235 and GmTag237 due to limited coverage during other phases. Though these animals had already departed the area surrounding the range by the time the interphase period began, we include discussion of these metrics, as it is possible that there may have been ongoing effects to their behavior at this point caused by exposure during the A/B mixed phase. Dawn dive rates fell sharply between the interphase and after phase for both tags, while the percentage of surface time concurrently rose. Dawn dive depths between did not vary significantly between interphase and after phase for either tag, but dive durations were significantly longer during the after phase for both tags.

Day dive rates for the three tags varied substantially between phases, and ranged from 0.66 dives/hr (GmTag237, after phase) to 3.12 dives/hr (GmTag237, A/B mixed phase). Day dive rates were highest during the A/B mixed phase for both tag GmTag235 and GmTag237, and generally decreased continuously in subsequent phases. Conversely, the percentage of surface time generally rose following the A/B mixed phase for both tags, although it did decrease slightly during the B/A mixed phase for GmTag235. Day dive depths and durations had statistically significant variation between phases for GmTag235 and GmTag237. For GmTag235, dives were significantly shallower during the A/B mixed phase and interphase compared to the after phase, and for GmTag237 dives were significantly shallower during the A/B mixed phase compared to the B/A mixed phase and after phase, as well as shallower during the interphase compared to subsequent phases. In regard to dive durations, dives were significantly shorter during the A/B mixed phase and interphase compared to both the B/A mixed phase and after phase for GmTag235 and GmTag237.

Dusk dive rates had minimal variation between phases, and ranged from 2.27 (GmTag237, after phase) to 2.82 (GmTag235, A/B mixed phase). Similarly, the percentage of surface time had minimal variation for GmTag237, although GmTag235 had a slight decrease in the percentage of surface time between the A/B mixed phase and the after phase. There was no statistically significant variation in dusk dive depths or durations between phases.

Night dive rates had moderate variation between phases and were higher than any other time of day. Night dive rates decreased in all phases following the A/B mixed phase for both GmTag235 and GmTag237. The percentage of surface time at night remained mostly consistent between phases for GmTag235, but rose slightly between the A/B mixed phase and interphase for GmTag237. There was no statistically significant variation in night dive depths between phases for either tag, but dives were significantly longer during the after phase compared to the A/B mixed phase and interphase for GmTag235.




Figure 18 - Top. Boxplot showing dive depths of GmTag235 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag235 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 129.2 dB re 1  $\mu$ Pa.




Figure 19 - Top. Boxplot showing dive depths of GmTag237 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag237 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 128.8 dB re 1  $\mu$ Pa.

## False killer whales

Only one false killer whale, from the main Hawaiian Islands insular population, was tagged off Kaua’i in 2021; summary statistics for the deployment are given in Table 14.

*Table 14 - Percentage of dive/surfacing data by phase for false killer whale PcTag074. The percentage of behavioral coverage is defined as the proportion of the duration of behavioral data relative to the duration of the tag within each phase.*

Individual	Percentage of dive/surfacing data				
	Before	Phase A	Interphase	Phase B	After
Duration overall (days)	3.5	1.7	3.8	2.4	0.7
Days surfacing/dive data	3.0	1.7	3.4	0.7	0.0
Percentage behavioral coverage	87.1	100.0	89.4	27.5	0.0

### *PcTag074*

Information was available about PcTag074’s movements in 2021 for all of Before (3.5 days), Phase A (1.7 days), the interphase period (3.8 days), Phase B (2.4 days), and the After period (0.7 days; Table 14). This individual ranged widely around both Kaua’i and Ni’ihau and O’ahu, but was only in proximity to the range during Phase B. As this animal was closest to O’ahu during Phase A, Phase B is the only period in which this animal would likely have been exposed to Navy activity.

After tagging (Before), this individual moved along the southern coast of Kaua’i and east to O’ahu, spending its time in waters around O’ahu for the entirety of Phase A and the beginning of the interphase (**Error! Reference source not found.**). Approximately halfway through the interphase, PcTag074 moved offshore and northwest towards Ni’ihau, and followed the northern coastline of Ni’ihau for the start of Phase B. Mirroring its movements post-deployment, PcTag074 gradually moved southeast along the southern edge of the range and continued to follow the southern coastline of Kaua’i during Phase B. While PcTag074 was at the southernmost part of the range, it was exposed to RLs ranging between 77.6 and 82.6 dB re 1  $\mu$ Pa. This individual continued its movements back towards O’ahu where exposures occurred during a second MFAS bout (44.6 - 90.5 dB re 1  $\mu$ Pa). The tag stopped transmitting shortly after Phase B, on the northeast side of O’ahu.

PcTag074 was exposed to five bouts of MFAS over three days. The first bout had low probabilities of exposures (0.2 – 0.65), but the rest of the 5 min bins had probabilities of 1 with the exception of one location in the 3<sup>rd</sup> bout with a low probability of 0.09. Their RLs were very low, with 10 of 23 5-min bins having estimated median levels below the ambient noise floor, and the remainder of the 5-min bins having median RLs equal to or less than 91 dB re 1  $\mu$ Pa (Table 15, Figure 20). While the first bout of MFAS occurred while the animal was closer to the range, their position was largely blocked by the island and by the bathymetry of the area. In contrast, while they were much further away for the brief second bout, the estimated RLs were higher due to a direct path in that case.

*Table 15 - Estimated received levels, cumulative sound exposure levels, and ship CPA for PcTag074*

Minimum median RL value ( $\pm$ 2 SD) dB re 1 $\mu$ Pa	NA
Maximum median RL value ( $\pm$ 2 SD) dB re 1 $\mu$ Pa	90.5 (71.7, 109.3)
CPA of ship (km)	29.0
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	98.3




Figure 20 - Median RLs for PcTag074 in stoplight colors (green for few pings during 5-min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.

### PcTag074 - August 2021




Figure 21 - Movements of PcTag074 during and after the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the three MFAS bouts that PcTag074 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary.

Dive behavior data were transmitted by the tag during all phases except for the After period. However, when broken down by time of day and phase, only the data from Before, Phase A, and the Interphase periods met the coverage requirements for inclusion in the analysis. Given that this animal was not exposed to Navy activities during any of these periods, this tag is given no further consideration for dive behavior analyses.

## Melon-headed whales

Seven melon-headed whales had tag data during the SCCs (**Error! Reference source not found.**), three in 2021 and four in 2022. However, there was only enough dive data to analyze the night-time periods for three of the animals. (Table 17).

*Table 16 - Percentage of dive/surfacing data by phase for melon-headed whales. The percentage of behavioral coverage is defined as the proportion of the duration of behavioral data relative to the duration of the tag within each phase.*

Individual	Percentage of dive/surfacing data				
	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After
<b>PeTag029</b>					
Duration overall (days)	10.4	1.7	0.5	0.0	0.0
Days surfacing/dive data	1.2	0.1	0.0	0.0	0.0
Percentage behavioral coverage	11.1	4.8	0.0	0.0	0.0
<b>PeTag031</b>					
Duration overall (days)	0.4	1.7	3.8	2.4	12.4
Days surfacing/dive data	0.0	1.6	3.7	2.4	3.2
Percentage behavioral coverage	0.0	95.8	98.1	100.0	25.7
<b>PeTag032</b>					
Duration overall (days)	NA	NA	3.8	2.4	10.2
Days surfacing/dive data	NA	NA	3.1	0.2	0.7
Percentage behavioral coverage	NA	NA	82.8	6.3	6.9
<b>PeTag033</b>					
Duration overall (days)	NA	2.6	3.5	0.2	4.8
Days surfacing/dive data	NA	1.3	0.3	0.0	0.8
Percentage behavioral coverage	NA	51.0	9.5	0.0	16.2
<b>PeTag034</b>					
Duration overall (days)	NA	2.6	1.5	0.0	0.0
Days surfacing/dive data	NA	0.1	0.0	0.0	0.0
Percentage behavioral coverage	NA	3.1	0.0	0.0	0.0
<b>PeTag035</b>					
Duration overall (days)	NA	NA	0.1	0.2	18.1
Days surfacing/dive data	NA	NA	0.0	0.0	4.2
Percentage behavioral coverage	NA	NA	0.0	0.0	23.3
<b>PeTag036</b>					
Duration overall (days)	NA	NA	NA	0.2	10.1
Days surfacing/dive data	NA	NA	NA	0.0	0.3
Percentage behavioral coverage	NA	NA	NA	0.0	2.6

Table 17 - A comparison of nighttime diving parameters from melon-headed whales exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

Dive parameter per individual	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Night dive rate (dives/hour)</i>							
PeTag031	NA	5.80	5.18	4.46	4.57	-	
PeTag032	NA	NA	5.49	NA	NA	-	
PeTag035	NA	NA	NA	NA	5.62	-	
<i>% time in surface periods at night</i>							
PeTag031	NA	29.79	33.27	45.94	39.11	-	
PeTag032	NA	NA	34.32	NA	NA	-	
PeTag035	NA	NA	NA	NA	35.70	-	
<i>Median dive depth night (m)</i>							
<b>PeTag031</b>	<b>NA</b>	<b>199.50</b>	<b>207.50</b>	<b>287.50</b>	<b>327.50</b>	<b>&lt;0.0001</b>	<b>Inter-B; A-B; Inter-After; A-After; B-After</b>
PeTag032	NA	NA	247.50	NA	NA	NA	NA
PeTag035	NA	NA	NA	NA	195.50	NA	NA
<i>Median dive duration night (min)</i>							
<b>PeTag031</b>	<b>NA</b>	<b>7.27</b>	<b>8.00</b>	<b>7.40</b>	<b>8.07</b>	<b>&lt;0.0001</b>	<b>Inter-A; Inter-B; A-After; B-After</b>
PeTag032	NA	NA	7.27	NA	NA	NA	NA
PeTag033	NA	NA	NA	NA	7.20	NA	NA



PeTag029

Information was available on movement patterns for PeTag029 in 2021 for Before (10.4 days), Phase A (1.7 days), and the interphase period (0.5 days; **Error! Reference source not found.**). During the Before period, PeTag029 ranged widely, entering and exiting the range multiple times, and was exposed to navy activities during a ULT period. While PeTag029 had been moving southeast prior to exposure during the ULT, following exposure it quickly changed direction to move northeast, then changed direction soon again and began moving northwest until it exited the range. During the ULT, PeTag029 was exposed to median received levels exceeding 135 dB re 1  $\mu$ Pa, and reaching 147.7 dB re 1  $\mu$ Pa for over two hours (Figure 23).

PeTag029 was exposed to a single long block of MFAS during the ULT prior to the SCC, but their tag stopped transmitting prior to the onset of Phase B of the SCC. They were on the range for the first few days of their tag deployment, but were off the range to the east during the 2-hour period of exposure; about 5 hours after the end of the ULT, the whale crossed the range and then began traveling west then south away from the range. All of their track locations during the exposure had probabilities of exposure of 1, and since they were just east of the training activity, their median RLs were similar throughout the exposure (137 – 148 dB re 1  $\mu$ Pa; Table 18, Figure 22), most of their 5-min bins had high numbers of pings, and distances to ships were < 20 km. This animal had the highest median and +2 SD RL values of all the whales tagged in 2021 due to its proximity to the activity. They also had the highest maximum cSEL at 158.1 dB re 1  $\mu$ Pa<sup>2</sup>.

Table 18 - Estimated received levels, cumulative sound exposure levels, and ship CPA for PeTag029

Minimum median RL value ( $\pm$ 2 SD) dB re 1 $\mu$ Pa	137.2 (124.9, 149.6)
Maximum median RL value ( $\pm$ 2 SD) dB re 1 $\mu$ Pa	147.7 (139.1, 156.2)
CPA of ship (km)	24.8
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	158.1





Figure 22 - Median RLs for PeTag029 in stoplight colors (green for few pings during 5-min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.



**PeTag029 - August 2021, MFAS exposures during ULT**




Figure 23 - Top: Movements of PeTag029 prior to the August 2021 SCC event; this tag stopped transmitting before Phase B of the SCC event, but was exposed to MFAS during a ULT period that occurred prior to Phase A of the SCC (see text for description of phases). Bottom: Movements of PeTag029 during the MFAS exposures that occurred during the ULT event. The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level. Additionally, the RL circles are colored by "intensity" which is characterized by the frequency of MFAS exposures that occurred during that given 5-

*minute exposure bin. The gray shaded rectangular polygon represents the area of ship activity during the ULT and the corresponding diamond point represents the mean ship location during the ULT. The dashed black line represents the PMRF boundary.*

Dive behavior data were transmitted by PeTag029 for Before and Phase A, but when broken down by time of day and phase, the data coverage for both phases failed to meet the required coverage for inclusion in the dive behavior analysis.

*PeTag031, PeTag032*

PeTag031 and PeTag032 were tagged in the same group in 2021, and generally remained associated over the overlapping period of tag attachment. Information was available on movement patterns for PeTag031 for Before (0.4 days), and for both tags for Phase A (1.7 and 0.04 days), the interphase period (3.8 days), Phase B (2.4 days), and the After period (12.4 and 10.2 days; **Error! Reference source not found.**). Both individuals moved on and off the range multiple times over the course of the SCC, but consistently remained in close proximity to the range.

PeTag031 moved towards the east of the range after tag deployment (Figure 26). At the start of Phase A, this individual moved southwest through the southern portion of the range and in the canyon area between Kaua’i and Ni’ihau. During the interphase period (between Phase A and B), PeTag031 spent much of its time in this area between the two islands; PeTag032 was deployed in this area at this time (Figure 27). Both tagged individuals continued using this area, with some movements farther north of the channel including use of the southwest portion of the range. Both whales were in the area just north of Ni’ihau prior to the start of Phase B. Both individuals were exposed to their highest MFAS levels (maximum median RL of 136.6 dB re 1  $\mu$ Pa for PeTag031, 135.5 dB re 1  $\mu$ Pa for PeTag032) for approximately one hour at the start of Phase B when they entered the southwestern edge of the range. As they continued moving northwest into the range, they were exposed to another bout of MFAS (approximately one hour long, similar levels as first bout). They both gradually moved northeast and exited the eastern portion of the range where they were exposed to one last bout of MFAS that lasted just over six hours. Received levels were slightly lower than the first two bouts (between 105 and 128 dB re 1  $\mu$ Pa), and generally followed a decreasing trend over time. After this bout, they both moved west back inside the range and continued moving northwest outside of the range for the remainder of Phase B (no exposures during this period). In the three days following Phase B, both individuals moved north of the range, entering and exiting the northern portion of the range twice. At the end of the three days post-SCC, both individuals were just outside of the northwestern area of the range (Figure 26, Figure 27).

PeTag031 was on or relatively near the range for the full duration of their tag attachment period. They were exposed to six bouts of MFAS, and all 5-min bins had a probability of exposure of 1. Median RLs ranged from 108 to 137 dB re 1  $\mu$ Pa (Table 19, Figure 24). They did move further away from the area of training activity between the 2nd and 3<sup>rd</sup> bouts, as median RLs were about 10-20 dB lower than they were for the 1<sup>st</sup> bout.

*Table 19 - Estimated received levels, cumulative sound exposure levels, and ship CPA for PeTag031*

Minimum median RL value ( $\pm$ 2 SD) dB re 1 $\mu$ Pa	108.4 (83.3, 133.5)
Maximum median RL value ( $\pm$ 2 SD) dB re 1 $\mu$ Pa	136.6 (120.9, 152.2)

CPA of ship (km)	0.8
Maximum overall cSEL dB re 1 $\mu\text{Pa}^2$	142.0




Figure 24 - Median RLs for Pe31 in stoplight colors (green for few pings during 5-min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.

PeTag032 had similar exposure patterns and estimated RLs, with all exposures having a probability of 1 and median RLs ranging between 108 and 135 dB re 1  $\mu\text{Pa}$  (Table 20, Figure 25). Again, their first two bouts of MFAS had higher median RLs values than the last three bouts, indicating the animals likely moved away from the main area of activity.

Table 20 - Estimated received levels, cumulative sound exposure levels, and ship CPA for PeTag032

Minimum median RL value ( $\pm 2$ SD) dB re 1 $\mu\text{Pa}$	107.9 (102.9, 113.0)
Maximum median RL value ( $\pm 2$ SD) dB re 1 $\mu\text{Pa}$	135.5 (122.3, 148.7)
CPA of ship (km)	5.3
Maximum overall cSEL dB re 1 $\mu\text{Pa}^2$	141.8




Figure 25 - Median RLs for PeTag032 in stoplight colors (green for few pings during 5-min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.

### PeTag031 - August 2021




Figure 26 - Movements of PeTag031 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the three MFAS bouts that PeTag031 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary.

## PeTag032 - August 2021




Figure 27 - Movements of PeTag032 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the three MFAS bouts that PeTag032 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary.

### Dive behavior

Dive behavior data were transmitted by PeTag031 for Phase A, and by both PeTag031 and PeTag032 for the interphase, Phase B, and After. However, only the data from PeTag031 for all phases, and PeTag032 for the interphase period met the coverage requirements for inclusion in the analysis (Figure 28). Given that only PeTag031 offered multiple phases to compare, PeTag032 was given no further consideration in the dive analysis. As melon-headed whales dive almost exclusively at night, the dive behavior analysis is also restricted to this time of day.

Night dive rates for PeTag031 were highest during Phase A, and decreased during the interphase and Phase B, then rose slightly during After. Conversely, the percentage of surface time at night was lowest during Phase A, and increased during the interphase and Phase B, then dropped slightly during After. Night dives increased in depth following Phase A, and were significantly deeper during Phase B and After compared to all previous phases. Night dive durations however, had a more mixed trend. Night dives were shorted during Phase A, slightly longer during the interphase, slightly shorter during Phase B, and longest during After. Durations were significantly longer during the interphase and After compared to Phase A and Phase B.




Figure 28 - Top. Boxplot showing dive depths of PeTag031 by SCC Phase and time of day. Bottom. Barplot showing dive rates of PeTag031 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 136.6 dB.



*PeTag033, PeTag034*


PeTag033 and PeTag034 were both tagged in the same group in 2022 and remained in close proximity for at least the initial part of their deployments. Information was available on movements of both tags for the A/B mixed phase (2.6 days), the interphase period (3.5 days for PeTag033, and 1.5 days for PeTag034), and for PeTag033 for the B/A mixed phase (0.2 days) and after phase (4.8 days; **Error! Reference source not found.**).

Both PeTag033 and PeTag034 moved south and to offshore waters shortly after tag deployment (Figure 31, Figure 32). As they continued south, they were exposed during several MFAS bouts albeit received levels were very low given the large distance from the range (all under 70 dB re 1 μPa). PeTag034’s deployment ended shortly after Phase A in offshore waters south of Ni’ihau. PeTag033 turned north after the exposures and moved northeast through waters between Kaua’i and Ni’ihau for the interphase period. For the second mixed phase A/B period, this individual started heading east towards O’ahu (no exposures) and continued this movement through the end of the study period (3-days post SCC). Insufficient data was available to include either tag in the dive behavior analysis.

Because PeTag033 immediately left the range area immediately after being tagged, all estimated RLs were at or below 70 dB re 1 μPa, and most were below the ambient noise floor (Table 21, Figure 29).

*Table 21 - Estimated received levels, cumulative sound exposure levels, and ship CPA for PeTag033*

Minimum median RL value (± 2 SD) dB re 1 μPa	NA
Maximum median RL value (± 2 SD) dB re 1 μPa	70.1 (63.5, 76.7)
CPA of ship (km)	124.2
Maximum overall cSEL dB re 1 μPa <sup>2</sup>	79.6



*Figure 29 - Median RLs for PeTag033 in stoplight colors (green for few pings during 5-min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the ± 2 SD values.*

PeTag034’s estimated RLs were also quite low, with a max of 73 dB re 1  $\mu$ Pa and most below the ambient noise floor (Table 22, Figure 30). They also had several 5-min bins with probabilities less than 1 of exposure.

Table 22 - Estimated received levels, cumulative sound exposure levels, and ship CPA for PeTag034

Minimum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	NA
Maximum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	72.6 (61.3,84.0)
CPA of ship (km)	164.3
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	77.4




Figure 30 - Median RLs for PeTag034 in stoplight colors (green for few pings during 5-min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.

### PeTag033 - August 2022




Figure 31 - Movements of PeTag033 during the August 2022 SCC event, including extensive movements away from the range after exposure phases. Note that for this particular event, ship-based MFAS activity occurred during both A and B phases. The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the MFAS bouts that PeTag033 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary.

## PeTag034 - August 2022




Figure 32 - Movements of PeTag034 during the August 2022 SCC event, including extensive movements away from the range after exposure phases. Note that for this particular event, ship-based MFAS activity occurred during both A and B phases. The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by "intensity" which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the MFAS bouts that PeTag034 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary.

### Rough-toothed dolphins

While three rough-toothed dolphins were tagged (Table 23), only one animal had any exposures to MFAS. SbTag025 did have some overlap with Phase B/A mixed, but did not received any MFAS exposures. Diel dive data were assessed for two of the animals (Table 24 and Table 25).

*Table 23 - Percentage of dive/surfacing data by phase for rough-toothed dolphins. The percentage of behavioral coverage is defined as the proportion of the duration of behavioral data relative to the duration of the tag within each phase.*

Individual	Percentage of dive/surfacing data				
	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After
<b>SbTag023</b>					
Duration overall (days)	2.3	1.7	3.8	2.4	2.8
Days surfacing/dive data	1.9	1.7	3.8	2.4	0.7
Percentage behavioral coverage	83.2	100.0	100.0	100.0	24.5
<b>SbTag024</b>					
Duration overall (days)	2.3	1.7	0.5	0.0	0.0
Days surfacing/dive data	NA	NA	NA	NA	NA
Percentage behavioral coverage	NA	NA	NA	NA	NA
<b>SbTag025</b>					
Duration overall (days)	NA	NA	NA	0.2	7.3
Days surfacing/dive data	NA	NA	NA	0.2	5.8
Percentage behavioral coverage	NA	NA	NA	100.0	80.2

Table 24 - A comparison of daytime diving parameters from rough-toothed dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

Dive parameter per individual	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Day dive rate (dives/hour)</i>							
SbTag023	0.00	0.24	0.18	0.32	NA	-	
SbTag025	NA	NA	NA	NA	0.11	-	
<i>% time in surface periods at day</i>							
SbTag023	100.00	99.06	98.81	98.85	NA	-	
SbTag025	NA	NA	NA	NA	99.67	-	
<i>Median dive depth day (m)</i>							
SbTag023	NA	76.50	159.50	82.50	NA	0.1363	NA
SbTag025	NA	NA	NA	NA	76.50	NA	NA
<i>Median dive duration day (min)</i>							
<b>SbTag023</b>	<b>NA</b>	<b>2.18</b>	<b>3.95</b>	<b>2.20</b>	<b>NA</b>	<b>0.0019</b>	<b>Inter-A; Inter-B</b>
SbTag025	NA	NA	NA	NA	1.95	NA	NA

Table 25 - A comparison of nighttime diving parameters from rough-toothed dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

Dive parameter per individual	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Night dive rate (dives/hour)</i>							
SbTag023	4.16	3.22	2.20	3.30	NA	-	
SbTag025	NA	NA	NA	NA	2.19	-	
<i>% time in surface periods at night</i>							
SbTag023	73.72	80.79	87.76	81.63	NA	-	
SbTag025	NA	NA	NA	NA	91.71	-	
<i>Median dive depth night (m)</i>							
<b>SbTag023</b>	<b>81.50</b>	<b>73.50</b>	<b>91.50</b>	<b>92.50</b>	<b>NA</b>	<b>0.0005</b>	<b>Inter-A; A-B; Inter-Before</b>
SbTag025	NA	NA	NA	NA	71.50	NA	NA
<i>Median dive duration night (min)</i>							
<b>SbTag023</b>	<b>3.83</b>	<b>3.65</b>	<b>3.40</b>	<b>3.32</b>	<b>NA</b>	<b>0.0154</b>	<b>Inter-Before; B-Before</b>
SbTag025	NA	NA	NA	NA	2.23	NA	NA

### *SbTag023*

Information was available about SbTag023's movements in 2021 for Before (2.3 days), Phase A (1.7 days), the interphase period (3.8 days), Phase B (2.4 days), and After (2.8 days; Table 24). During this time, SbTag023 remained on or in close proximity to the range throughout the course of the SCC.

SbTag023 was tagged prior to the start of Phase A; this individual largely remained on the southern end of the range prior to and during Phase A (Figure 34). At the end of Phase A, SbTag023 had exited the southernmost end of the range and remained in this area (slope between Ni'ihau and Kaua'i) for most of

the interphase period, before moving north back into the range and then just outside the southwestern portion of the range prior to the start of Phase B. SbTag023 re-entered the range after the start of Phase B, and moved northeast along the southeastern area of the range where it was exposed to its highest received levels during two MFAS bouts (maximums 141.3 and 141.9 dB re 1  $\mu$ Pa for each, approximately one hour break in between). This individual continued moving northeast and was on the edge and just outside of the southeastern portion of the range during the last MFAS bout that it was exposed to (levels between 105 - 130 dB re 1  $\mu$ Pa). After this bout, SbTag023 re-entered the range and moved southwest across the range through the end of Phase B. During the three days post-SCC, SbTag023 largely remained on the southwestern edge of the range.

Because SbTag023 remained on or close to the range for their entire tag attachment period, all of their 5-min bins had probabilities of 1 for exposure, and their estimated median RLs ranged from 107 – 142 dB re 1  $\mu$ Pa (Table 26, Figure 33). They were exposed to six bouts of MFAS, and the first two bouts had higher median RLs than the last four bouts by 10 – 20 dB. This animal had the highest estimated median and + 2 SD RL values for the animals tagged in 2022.

Table 26 - Estimated received levels, cumulative sound exposure levels, and ship CPA for SbTag023

Minimum median RL value ( $\pm$ 2 SD) dB re 1 $\mu$ Pa	106.8 (94.7, 118.9)
Maximum median RL value ( $\pm$ 2 SD) dB re 1 $\mu$ Pa	141.9 (121.2, 156.6)
CPA of ship (km)	13.9
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	150.5




Figure 33 - Median RLs for SbTag023 in stoplight colors (green for few pings during 5-min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm$  2 SD values.



### SbTag023 - August 2021




Figure 34 - Movements of SbTag023 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times are given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the three MFAS bouts that SbTag023 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary.

#### Dive Behavior

Dive behavior data were available for SbTag023 for all phases (Table 24, Table 25, Figure 35). However, when broken down by time of day and phase, not every time of day and phase met the required coverage (relative to the phase duration) for inclusion in the analysis.

Behavior data for the dawn hours were available for Phase A, the interphase period, and Phase B, but the tagged individual remained exclusively at the surface during Phase A and the Interphase. Day dive metrics were available for all phases except After. No day dives were recorded during Before, and day dive rates for the remaining phases had only slight variation. The day dive rate fell slightly between Phase A and the interphase, and rose to its highest during Phase B. The percentage of surface time during day hours remained high across all phases, never dipping below 98%. There was no statistically

significant variation in day dive depths, but dives were significantly longer during the interphase compared to Phase A and Phase B.

Dusk dive metrics were available for Phase A, the interphase period, and Phase B, but no dusk dives were recorded during Phase B. The dusk dive rate was highest during Phase A and dropped sharply during the interphase. The percentage of surface time during dusk hours was lowest during Phase A and rose during the interphase. Dusk dive depths and durations did not vary significantly between phases.

Night dive metrics were available for all phases except After. Night dive rates were highest during Before, dropped during Phase A, continued to drop during the interphase, and rose during Phase B. The percentage of surface time during night hours was lowest during Before, rose during Phase A and the interphase, and dropped again during Phase B. Night dives were significantly deeper during the interphase compared to Before, and significantly deeper during the interphase and Phase B compared to Phase A. Night dives were also significantly longer during Before compared to the interphase or Phase B.





Figure 35 - Top. Boxplot showing dive depths of *SbTag023* by SCC Phase and time of day. Bottom. Barplot showing dive rates of *SbTag023* by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 141.9 dB.

### Common bottlenose dolphins

Three bottlenose dolphins had tag data during the SCCs, two in 2021 and one in 2022 (Table 27). Diel dive analyses could be conducted for one animal from each year (**Error! Reference source not found.** and Table 29).

*Table 27 - Percentage of dive/surfacing data by phase for common bottlenose dolphins. The percentage of behavioral coverage is defined as the proportion of the duration of behavioral data relative to the duration of the tag within each phase.*

Individual	Percentage of dive/surfacing data				
	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After
<b>TtTag039</b>					
Duration overall (days)	NA	1.0	3.8	2.4	5.7
Days surfacing/dive data	NA	0.7	3.0	2.0	1.4
Percentage behavioral coverage	NA	75.3	80.4	83.8	25.5
<b>TtTag040</b>					
Duration overall (days)	NA	NA	2.8	2.4	10.7
Days surfacing/dive data	NA	NA	NA	NA	NA
Percentage behavioral coverage	NA	NA	NA	NA	NA
<b>TtTag041</b>					
Duration overall (days)	NA	NA	3.0	0.2	13.8
Days surfacing/dive data	NA	NA	3.0	0.2	5.0
Percentage behavioral coverage	NA	NA	100.0	100.0	36.1

Table 28 - A comparison of daytime diving parameters from common bottlenose dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

Dive parameter per individual	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Day dive rate (dives/hour)</i>							
TtTag039	NA	NA	0.85	0.69	NA	-	
TtTag041	NA	NA	0.82	1.11	0.75	-	
<i>% time in surface periods at day</i>							
TtTag039	NA	NA	90.06	94.15	NA	-	
TtTag041	NA	NA	87.59	89.87	88.32	-	
<i>Median dive depth day (m)</i>							
<b>TtTag039</b>	<b>NA</b>	<b>NA</b>	<b>123.50</b>	<b>71.50</b>	<b>NA</b>	<b>0.0218</b>	<b>Inter-B</b>
<b>TtTag041</b>	<b>NA</b>	<b>NA</b>	<b>607.50</b>	<b>116.50</b>	<b>591.50</b>	<b>0.0102</b>	<b>Inter-B/A; B/A-After</b>
<i>Median dive duration day (min)</i>							
<b>TtTag039</b>	<b>NA</b>	<b>NA</b>	<b>7.50</b>	<b>5.17</b>	<b>NA</b>	<b>0.0068</b>	<b>Inter-B</b>
<b>TtTag041</b>	<b>NA</b>	<b>NA</b>	<b>9.67</b>	<b>5.57</b>	<b>10.20</b>	<b>0.0183</b>	<b>Inter-B/A; B/A-After</b>

Table 29 - A comparison of nighttime diving parameters from common bottlenose dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

Dive parameter per individual	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Night dive rate (dives/hour)</i>							
TtTag039	NA	NA	4.01	5.08	3.83	-	
TtTag041	NA	NA	2.95	NA	2.65	-	
<i>% time in surface periods at night</i>							
TtTag039	NA	NA	65.40	54.64	70.07	-	
TtTag041	NA	NA	67.84	NA	75.15	-	
<i>Median dive depth night (m)</i>							
TtTag039	NA	NA	115.50	97.50	109.50	0.0884	NA
TtTag041	NA	NA	199.50	NA	137.50	0.1106	NA
<i>Median dive duration night (min)</i>							
<b>TtTag039</b>	<b>NA</b>	<b>NA</b>	<b>4.67</b>	<b>5.23</b>	<b>4.78</b>	<b>0.0472</b>	<b>B-After</b>
<b>TtTag041</b>	<b>NA</b>	<b>NA</b>	<b>6.60</b>	<b>NA</b>	<b>4.70</b>	<b>0.0142</b>	<b>Inter-After</b>

### TtTag039

Information was available about TtTag039's movements in 2021 for Phase A (1.0 days), the interphase period (3.8 days), Phase B (2.4 days), and After (5.7 days; Table 27). TtTag039 spent time during each phase on the range, but also moved extensively around the island of Kaua'i.

Movements of TtTag039 relative to SCC phases are shown in Figure 37. After tag deployment, TtTag039 spent its time during Phase A in the nearshore waters along the northern Kaua'i coast. During the interphase, TtTag039 began moving southwest along the easternmost edge of the south end of the range, continued following the coast (counter-clockwise) and remained in the northeastern coast of Kaua'i through the end of the interphase. At the start of Phase B, this individual began moving south along the east coast of the island where it was exposed to one bout of MFAS for approximately one hour

(received levels remained under 100 dB re 1  $\mu$ Pa). TtTag039 continued south and followed the coastline back around to the range and near its deployment location through the end of Phase B. In the three days after Phase B, this individual moved along the north-northwest coastline of Kaua'i.

TtTag039 only had one short bout of exposure, with only nine 5-min bins with estimated RLs equal to or less than 100 dB re 1  $\mu$ Pa and all had probabilities of exposures less than 0.29 (Table 30, Figure 36).

Table 30 - Estimated received levels, cumulative sound exposure levels, and ship CPA for TtTag039

Minimum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	90.0 (73.0, 107.0)
Maximum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	100.0 (83.5, 116.4)
CPA of ship (km)	34.8
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	106.4




Figure 36 - Median RLs for TtTag039 in stoplight colors (green for few pings during 5-min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.

### TtTag039 - August 2021




Figure 37 - Movements of TtTag039 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the three MFAS bouts that TtTag039 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary.

#### Dive Behavior

Dive behavior data were available for TtTag039 for all phases except Before (**Error! Reference source not found.**, Table 29, Figure 38). However, when broken down by time of day and phase, not every time of day and phase met the required coverage (relative to the phase duration) for inclusion in the analysis. Coverage was insufficient during dawn to calculate any metrics, and only metrics for the interphase period during dusk hours could be calculated. As a result, analysis was restricted to the day and night hours.

Day dive metrics were available for the interphase and Phase B. The day dive rate was highest during the interphase, and dropped slightly during Phase B, while the percentage of surface time during day hours



was lowest during the interphase, and rose slightly during Phase B. Day dives were significantly deeper and longer during the interphase than Phase B.

Night dive metrics were available for the interphase, Phase B, and After. The night dive rate was highest during Phase B, and dropped sharply during After, while the percentage of surface time was lowest during Phase B, and rose sharply during After. Night dive depths did not have statistically significant variation between phases, but night dives were significantly shorter during After compared to Phase B.




Figure 38 - Top. Boxplot showing dive depths of TtTag039 by SCC Phase and time of day. Bottom. Barplot showing dive rates of TtTag039 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 100.0 dB.

*TtTag040*


Information was available about TtTag040’s movements in 2021 for the interphase period (3 days), Phase B (0.2 days), and After (13.8 days; Table 27). TtTag040 was not closely associated with TtTag039 for the entirety of its deployment period, although exhibited similar movement patterns and generally remained along the nearshore coasts of Kaua’i.

TtTag040 was tagged during the interphase period; during this time, this individual moved along the western coast of Kaua’i (southeastern edge of the range), and then eventually along the northern coast (Figure 40). At the start of Phase B, this individual was on the eastern side of Kaua’i and was exposed to two bouts of MFAS, with exposure periods (when received levels were computed) lasting about an hour each; received levels were under 100 dB re 1 μPa during both bouts. TtTag040 continued to use the nearshore waters on the east coast of Kaua’i throughout the rest of Phase B. In the three days after Phase B, this individual moved west along the southern coast of the island, then back along the east coast and eventually the north coast of Kaua’i.

TtTag040 was exposed to three separate bouts of MFAS over two different days, they also had less than 0.77 probability of exposure for any of their 5-min bins, and had estimated median RLs of 100 dB re 1 μPa or less (Table 31, Figure 39).

*Table 31 - Estimated received levels, cumulative sound exposure levels, and ship CPA for TtTag040*

Minimum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	85.4 (78.2, 92.7)
Maximum median RL value ( $\pm 2$ SD) dB re 1 $\mu$ Pa	100.5 (82.3, 118.7)
CPA of ship (km)	54.5
Maximum overall cSEL dB re 1 $\mu$ Pa <sup>2</sup>	108.2



*Figure 39 - Median RLs for Tt040 in stoplight colors (green for few pings during 5-min bin, yellow for moderate number of pings, red for a high number of pings) with error bars giving the  $\pm 2$  SD values.*

### TtTag040 - August 2021




Figure 40 - Movements of TtTag040 during the August 2021 SCC event (see text for description of phases). The maximum, median estimated received levels (RLs) that occurred during each 5-minute exposure bin are plotted as open circles, with the size of the circle scaled to RL level and times given in GMT. Additionally, the RL circles are colored by “intensity” which is characterized by the frequency of MFAS exposures that occurred during that given 5-minute exposure bin. The shaded rectangular polygons represent the area of ship activity during each of the three MFAS bouts that TtTag040 was exposed during and the corresponding diamond points represent the mean ship location during each bout. Note that After is restricted to three days after the end of the SCC. The dashed black line represents the PMRF boundary.

## DISCUSSION

This report describes the movement patterns, diel dive behavior, and estimated RLs for 15 of 21 satellite tagged odontocetes on and near PMRF during the two-week long SCC training events in August 2021 and 2022. These data add to the growing catalog of odontocetes that have been tagged concurrently with these training events since 2011. While received levels for these tagged animals have been reported since 2014, in 2021 a major re-analysis was conducted of 9 of 12 short-finned pilot whales, 4 of 7 rough-toothed dolphins, and 2 bottlenose dolphins using updated methods to estimate received levels more accurately (Henderson et al. 2021). These methods, including estimating RLs for the full error ellipse of each *crawl*-interpolated track position every 5-min and utilizing modeled dive depth information, allow for more rigorous estimations of RL in three dimensions (four, including time along the track). These methods were replicated in this report and support better assessments of behavioral response. New statistical methods of behavioral response analyses for both horizontal movement and dive behavior are being developed for satellite tags (e.g., Hewitt et al. 2022); by building up an extensive catalog of individuals and species, these aggregated methods may be applicable in the future for the species at PMRF as well.

The 15 tagged animals with MFAS exposures in 2021 and 2022 had relatively low RLs. PeTag029 had the highest levels, with a maximum median SPL of 148 dB re 1  $\mu$ Pa (up to 156 dB re 1  $\mu$ Pa with 2 SD), and a maximum cSEL of 158 dB re 1  $\mu$ Pa<sup>2</sup>. Two other melon-headed whales received maximum median RL values of 136-137 dB re 1  $\mu$ Pa, two short-finned pilot whales had RLs of 129 dB re 1  $\mu$ Pa, and one rough-toothed dolphin had maximum median RLs up to 142 dB re 1  $\mu$ Pa. The remainder of the animals' RLs were less than 100 dB re 1  $\mu$ Pa, and many of the median 5-min bin values were below the ambient noise floor of 60 dB re 1  $\mu$ Pa (at and above 1 kHz, Širović et al. 2013). These levels were lower than those for most of the animals re-analyzed in 2021, where median RLs ranged from 101 dB to as high as 177 dB re 1  $\mu$ Pa (Henderson et al. 2021), with the + 2 SD extending as high as 195 dB re 1  $\mu$ Pa, although these high values were at the tail end of the RL histograms and were very unlikely. In fact, in 2022, only three of the short-finned pilot whales were on the range during the SCC, and only during the A/B mixed phase, after which they left the range, and they only received exposures once off the range. Similarly, the false killer whale tagged in 2021 was also off the range during their periods of exposure, as were the two bottlenose dolphins tagged in 2021 and the two melon-headed whales tagged in 2022. The rough-toothed dolphin tagged in 2021 remained on the range for almost the whole SCC, and two of the melon-headed whales from 2021 moved on and off the range throughout the SCC, while the animal with the highest exposure (PeTag029) was on and off the range during the ULT prior to the SCC but then moved far out of the area before the SCC began. While the latter four tags could be candidates for a statistical analysis of their movement behavior, the rest of the animals had low enough received levels and were far enough away from the range that behavioral responses, if they occurred, were likely to be too subtle to be detected by coarse analyses of satellite tag data.

However, dive behavior were analyzed with a statistical approach for the animals who had enough data in each temporal period (dawn, day, dusk, night) and SCC period (Before, Phase A or A/B, Interphase, Phase B or B/A, and After). Differences in dive rates, time in surface periods, dive depth, and dive duration were compared across phases and diel periods, and dives were also identified as occurring on or off the range. However, the resulting changes in dive behavior across years, species, and individuals,

even among those in the same groups, were variable and contrasting, making it difficult to identify consistent patterns of response to Navy training activity or MFAS.

Of the pilot whales, only GmTag233, GmTag235, and GmTag237 had significant differences in their median dive durations in the dawn diel period; for GmTag233 significant differences occurred between Phase B and the After period, while for GmTag235 and GmTag237 differences occurred between the Interphase and the After period. During the daytime period, several of the short-finned pilot whales had significant differences in both their median dive depth and dive duration across SCC phases. While there were individual differences, the two most common phases with differences were the Interphase to After period, and the Phase A to After periods. However, the differences in dive behavior also varied between individuals, even within the same phases. For example, GmTag233 had deeper daytime dives (median depths > 500 m) in all phases except After, when their median dive depths were around 250 m. In contrast, GmTag234, GmTag235, and GmTag237 had very shallow median dive depths during Phase A and the Interphase, but then deep median depths during Phase B and After. Similarly, several of the short-finned pilot whales had significant differences in their median dive depths and durations for their nighttime dives across SCC phases. The 2021 pilot whales had variability in their dive depths across almost all phases, but again patterns diverged among individuals, with GmTag232 having all deep nighttime dives that got progressively deeper across phases, while GmTag234 had moderately deep (> 500 m) median depths that got progressively shallower during the SCC then deeper again After, and GmTag233 had shallow nighttime median dive depths (< 250 m) during the SCC but median dive depths jumped to > 500 m in the After period. The short-finned pilot whales tagged in 2021 were off the range during the SCC and many of their exposures were partially blocked, and so changes in dive behavior were more likely due to their location around the island and relative prey patterns in those areas rather than training activity. However, the short-finned pilot whales tagged in 2022 were on the range at the start of the A/B mixed phase; their exposures occurred at the southern end of the range when the animals were already moving south and away from the area. Exposure levels were highest during the first period of MFAS, up to a median RL of 125 dB re 1  $\mu$ Pa, and remained at similar levels while the animals were in the same general area during the second MFAS period. However, by the third MFAS period the animals had begun moving further south and so exposure levels were reduced. While dive rates were not significantly different across SCC phases, it may be worth noting that dive rates were highest during this A/B mixed Phase, when the animals were on the range near training activity and MFAS, and then rates went down across subsequent phases. In addition, both GmTag235 and GmTag237 had significantly shallower dives during the A/B mixed Phase than the B/A mixed phase (and After phase for GmTag237) when the animals had moved well away from the area on the far side of Kaua'i.

Similar changes and variability in dive behavior were found for short-finned pilot whales tagged prior to 2021 (Henderson et al. 2021). There were no significant differences in dive rates or time at the surface for any of the individuals, but a few animals had significant differences in daytime and nighttime median dive depths and durations. The patterns again varied among individuals, with some animals having variability in both daytime and nighttime dive depths across SCC phases (GmTag081, GmTag214), and other showing distinct changes across phases. For example, GmTag153 had very shallow median daytime and shallower (around 250 m) nighttime dive depths, with deeper dives in both diel periods in the After phase. In contrast, GmTag152's median dive depths were already deep (> 250 m at night and > 500 m during the day) and got deeper across SCC phases during both the daytime and nighttime periods, while GmTag214's median dive depths were deeper than 250 m both day and night during the SCC and

then became shallower than 250 m After. Therefore, across a decade of effort and with 15 individuals tagged, no clear and consistent patterns have emerged in changes to short-finned pilot whale dive behavior during exposure to Navy training activity and MFAS. In fact, in Henderson et al. (2021) a few of the short-finned pilot whales had potentially high levels of MFAS exposure because they approached active sources on the range, perhaps demonstrating either habituation to MFAS by resident animals, or indicating that the area is particularly important for foraging, and thus the animals may be more willing to tolerate MFAS or other training activity. Based on 24 satellite tag deployments on individuals from the western community of island-associated short-finned pilot whales in Hawai'i between 2008 and 2021, a Biologically Important Area was designated around Kaua'i, encompassing much of the southern part of the PMRF (Kratofil et al. 2023; see page 144, Supplemental File A).

For melon-headed whales, only nighttime dives could be assessed statistically, and only PeTag031 had significant differences in their median dive depths and durations across SCC phases. This animal remained on or near the range during all Phases and so had multiple bouts of MFAS exposure, although none exceeded median RLs of 137 dB re 1  $\mu$ Pa. Their median dive depths increased significantly across all phases from around 200 m during Phase A to over 300 m during the After period. Their dive rates also decreased across phases. In the same year, PcTag029 had the highest exposure levels as they were directly east of the ULT training activity and remained there for the duration of the ULT. As dive behavior was only compared across SCC phases it is unknown if there were any short-term changes in their dive behavior during this period. They did cross the range and then travel to the west and then south once the ULT was finished, which could have been a subsequent avoidance response of the area although they did not move away during the actual period of exposure. The two melon-headed whales tagged in 2022 left the area right after tagging and so all exposures were at low RLs and were partially blocked by land.

Of the three rough-toothed dolphins tagged in 2021, only two had enough dive data to conduct statistical analyses, and only one, SbTag023, had significant differences in their daytime median dive duration and nighttime median dive depth and dive durations. Median daytime dive depths were shallow (< 100 m) during both Phase A and B and then deepened to almost 200 m during the Interphase period. The pattern was less clear at night, although dives were slightly shallower in Phase A than in any other period, and were slightly deeper in the Interphase and Phase B than during Phase A. This animal's dive rates were highest in the Before period and lowest during the Interphase. This was also the only rough-toothed dolphin with MFAS exposures and had the second-highest median RLs in this study, up to 142 dB re 1  $\mu$ Pa, although they generally remained in the southern portion of the range for the exposure periods. They were tagged on the range and then continued to move on and off the range and into the channel between Ni'ihau and Kaua'i for the duration of the SCC. Similar results were found for rough-toothed dolphins in Henderson et al. (2021); only two of those four animals had sufficient dive data for a statistical analysis, and significant differences were found for both animals for their nighttime median dive depths and durations. These animals were tagged in the Interphase in 2016. SbTag017 had shallower (< 100 m) nighttime median dive depths during the Interphase and Phase B and then deeper (> 100 m) nighttime dives in the After period. SbTag018 had slightly shallower nighttime median dive depths in the Interphase and then slightly deeper median dive depths in Phase B and After, although in all three phases median dive depths were < 100 m. These animals again occupied the southern portion of the range or the area off the range in the channel between Ni'ihau and Kaua'i, and they had maximum median RLs of 148 dB re 1  $\mu$ Pa and 157 dB re 1  $\mu$ Pa (Henderson et al. 2021). Rough-toothed

dolphins generally occupy the same habitat at the southern end of the range (see Figure 5 in Kratofil et al. 2023), and are therefore not likely to experience much higher RLs than these as most training activity at PMRF takes place in the northern portion of the range, as can be seen in Figure 34.

Three common bottlenose dolphins were tagged in 2021 and 2022, but only two had sufficient dive data for analysis. Both the median dive depths and durations were statistically different across SCC phases for TtTag039 and TtTag041 during the day, while only durations were different for both at night. Both variables differed for TtTag039 between the Interphase and Phase B during the day, with deeper median and overall depths during the Interphase than during Phase B, and shorter dives at night After than during Phase B. However, this animal and TtTag040 circled Kaua'i throughout the whole SCC, and the few exposures that occurred were partially blocked by land and had low RLs. Therefore, like the pilot whales tagged in 2021 the changes in dive behavior for these animals were more likely due to variability in prey than any response to Navy training. For TtTag041, both dive depth and duration differed between the Interphase and Phase B, and between Phase B and After during the day, and duration differed between the Interphase and After at night. They had much deeper median dive depths (> 500 m) during the Interphase and After than during Phase B, when median dive depths were < 250 m. This animal also circumnavigated Kaua'i throughout the SCC and had no MFAS exposures, therefore again differences in their dive behavior were more likely due to foraging patterns than to Navy training. The two bottlenose dolphins analyzed in Henderson et al. (2021) also had significant differences in their median dive depths and durations both day and night across SCC phases. Both animals had deeper daytime than nighttime dives throughout the SCC, although TtTag034 had slightly deeper median daytime dives during Phase B than the Interphase or After period, while TtTag035 had slightly shallower daytime median dive depths during Phase B than the Interphase or After. Both animals had median nighttime dive depths that decreased from the Interphase (> 300 m) through Phase B and further into the After period, although not significantly for TtTag034. These animals actually remained on the range-side of Kaua'i during Phase B and therefore had median RLs up to 146 dB re 1  $\mu$ Pa. Much like the rough-toothed dolphins, bottlenose dolphins generally remain in the southern portion of the range (Kratofil et al. 2023, see page 36 in Supplemental File A), and even closer to shore than rough-toothed dolphins, and so again will not likely have exposure levels much higher than those.

The false killer whale, all six short-finned pilot whales, two bottlenose dolphins, and one rough-toothed dolphin were all resights of previously identified animals, and all but the three pilot whales from 2022 are known to be part of island-associated resident populations. The resident pilot whales, bottlenose dolphins, and rough-toothed dolphin have also been observed off the island of Kaua'i, and in fact the latter two species are specifically Kaua'i-island associated populations. Similarly, of the tagged delphinids assessed in Henderson et al. (2021), all but two of the short-finned pilot whales were from the Western Hawaiian population, and all of the rough-toothed and bottlenose dolphins were from the Kaua'i/Ni'ihau population. The two short-finned pilot whales that were from the pelagic population had also been previously photo-identified. Therefore, these data can begin to shed light on behavioral responses by animals that have likely been exposed more than once to Navy training activity and to MFAS. As there were no obvious changes in movement behavior, and as none of the changes in dive behavior were consistent enough to point to being responses to MFAS, none of these animals at least appear to be sensitized to MFAS. Likewise, in Henderson et al. (2021) there were no overt behavioral responses, and some animals approached active MFAS sources. While in Henderson et al. (2021) there were instances where the direction of travel abruptly changed to move away from MFAS, the direction

of travel often seems to change abruptly, as can be seen in multiple individuals in both that study and in this one. As these data are aggregated, and as more statistical tools are developed, these nuances in behavior may be able to be teased out further. While we must consider the argument by Forney et al. (2017) that populations such as these that are relegated to finite areas may have no choice other than to co-exist with anthropogenic stressors, and therefore no obvious response does not mean there isn't a stress response or other long-term, population level effect, at this time it does not appear as though resident odontocetes are responding to Navy training activity with MFAS at PMRF. However, future studies of stress hormones in these populations during these SCC-associated tagging efforts is recommended, as are drone studies comparing body composition of Kaua'i-associated populations compared to those associated with other Hawaiian islands. These types of studies may be able to detect responses that satellite tagging studies cannot. That said, these kind of long-term tagging studies that build up databases of multiple species are also invaluable, as behavior could change over time, and new methods are continually being developed that may be more sensitive to changes in satellite tag behavior than currently can be detected.



## LITERATURE CITED

- Andrews, R.D., Pitman, R.L. and Ballance, L.T., 2008. Satellite tracking reveals distinct movement patterns for Type B and Type C killer whales in the southern Ross Sea, Antarctica. *Polar Biology*, 31:1461-1468.
- Aschettino, J.M., R.W. Baird, D.J. McSweeney, D.L. Webster, G.S. Schorr, J.L. Huggins, K.K. Martien, S.D. Mahaffy, and K.L. West. 2012. Population structure of melon-headed whales (*Peponocephala electra*) in the Hawaiian Archipelago: evidence of multiple populations based on photo-identification. *Marine Mammal Science* 28(4):666-689. <https://doi.org/10.1111/j.1748-7692.2011.00517.x>
- Baird, R.W. 2016. The lives of Hawai'i's dolphins and whales: natural history and conservation. University of Hawai'i Press, Honolulu, Hawai'i. <https://doi.org/10.1515/9780824865931>.
- Baird, R.W. and Webster, D.L., 2020. Using dolphins to catch tuna: Assessment of associations between pantropical spotted dolphins and yellowfin tuna hook and line fisheries in Hawai'i. *Fisheries Research*, 230, 105652. <https://doi.org/10.1016/j.fishres.2020.105652>.
- Baird, R.W., Mahaffy, S.D. and Lerma, J.K., 2022. Site fidelity, spatial use, and behavior of dwarf sperm whales in Hawaiian waters: using small-boat surveys, photo-identification, and unmanned aerial systems to study a difficult-to-study species. *Marine Mammal Science*, 38(1):326-348. <https://doi.org/10.1111/mms.12861>.
- Baird, R.W., A.M. Gorgone, D.J. McSweeney, D.L. Webster, D.R. Salden, M.H. Deakos, A.D. Ligon, G.S. Schorr, J. Barlow and S.D. Mahaffy. 2008a. False killer whales (*Pseudorca crassidens*) around the main Hawaiian Islands: long-term site fidelity, inter-island movements, and association patterns. *Marine Mammal Science* 24:591-612. <https://doi.org/10.1111/j.1748-7692.2008.00200.x>.
- Baird, R.W., D.L. Webster, S.D. Mahaffy, D.J. McSweeney, G.S. Schorr and A.D. Ligon. 2008b. Site fidelity and association patterns in a deep-water dolphin: rough-toothed dolphins (*Steno bredanensis*) in the Hawaiian Archipelago. *Marine Mammal Science* 24(3):535-553. <https://doi.org/10.1111/j.1748-7692.2008.00201.x>
- Baird, R.W., A.M. Gorgone, D.J. McSweeney, A.D. Ligon, M.H. Deakos, D.L. Webster, G.S. Schorr, K.K. Martien, D.R. Salden, and S.D. Mahaffy. 2009. Population structure of island-associated dolphins: evidence from photo-identification of common bottlenose dolphins (*Tursiops truncatus*) in the main Hawaiian Islands. *Marine Mammal Science* 25(2):251-274. <https://doi.org/10.1111/j.1748-7692.2008.00257.x>
- Baird, R.W., Hanson, M.B., Schorr, G.S., Webster, D.L., McSweeney, D.J., Gorgone, A.M., Mahaffy, S.D., Holzer, D.M., Oleson, E.M. and Andrews, R.D., 2012. Range and primary habitats of Hawaiian insular false killer whales: informing determination of critical habitat. *Endangered Species Research*, 18(1):47-61. <https://doi.org/10.3354/esr00435>
- Baird, R.W., Webster, D.L., Aschettino, J.M., Schorr, G.S. and McSweeney, D.J., 2013. Odontocete cetaceans around the main Hawaiian Islands: Habitat use and relative abundance from small-boat sighting surveys. *Aquatic Mammals*, 39(3):253-269. <https://doi.org/10.1578/AM.39.3.2013.253>

Baird, R.W., Martin, S.W., Webster, D.L. and Southall, B.L., 2014. Assessment of modeled received sound pressure levels and movements of satellite-tagged odontocetes exposed to mid-frequency active sonar at the Pacific Missile Range Facility: February 2011 through February 2013. Prepared for US Pacific Fleet, submitted to NAVFAC PAC by HDR Environmental, Operations and Construction, Inc. *Prepared for US Pacific Fleet*.

Baird, R.W., Cholewiak, D., Webster, D.L., Schorr, G.S., Mahaffy, S.D., Curtice, C., Harrison, J. and Van Parijs, S.M., 2015a. 5. Biologically Important Areas for Cetaceans Within US Waters-Hawai'i Region. *Aquatic Mammals*, 41(1):54-64. <https://doi.org/10.1578/AM.41.1.2015.54>

Baird, R.W., Mahaffy, S.D., Gorgone, A.M., Cullins, T., McSweeney, D.J., Oleson, E.M., Bradford, A.L., Barlow, J. and Webster, D.L., 2015b. False killer whales and fisheries interactions in Hawaiian waters: evidence for sex bias and variation among populations and social groups. *Marine Mammal Science*, 31(2): 579-590. <https://doi.org/10.1111/mms.12177>

Baird, R.W., Martin, S.W., Manzano-Roth, R., Webster, D.L. and Southall, B.L., 2017. Assessing exposure and response of three species of odontocetes to mid-frequency active sonar during Submarine Commanders Courses at the Pacific Missile Range Facility: August 2013 through February 2015. Prepared for US Pacific Fleet, submitted to NAVFAC PAC by HDR Environmental, Operations and Construction, Inc.

Baird, R.W., Henderson, E.E., Martin, S.W., Southall, B.L. and Cascadia Research Collective, 2019. Assessing odontocete exposure and response to mid-frequency active sonar during Submarine Command Courses at the Pacific Missile Range Facility: 2016-2018. Prepared for Commander, Pacific Fleet, under Contract No. N62470-15-D-8006 Task Order KB16 issued to HDR Inc., Honolulu, HI.

Claridge, D.E., 2013. Population ecology of Blainville's beaked whales (*Mesoplodon densirostris*) (Doctoral dissertation, University of St Andrews).

Claridge, D.E., Dunn, C.A. 2013. Population consequences of acoustic disturbance of Blainville's beaked whales at AUTECH. Office of Naval Research.

Collins, M. (1993). "A split-step Pade solution for the parabolic equation method," *Journal of the Acoustical Society of America* 93, 1736-1742.

Falcone, E.A., Keene, E.L., Keen, E.M., Barlow, J., Stewart, J., Cheeseman, T., Hayslip, C. and Palacios, D.M. 2022. Movements and residency of fin whales (*Balaenoptera physalus*) in the California Current System. *Mammalian Biology*, 102(4): 1445-1462.

Falcone, E.A., Schorr, G.S., Douglas, A.B., Calambokidis, J., Henderson, E., McKenna, M.F., Hildebrand, J. and Moretti, D. 2009. Sighting characteristics and photo-identification of Cuvier's beaked whales (*Ziphius cavirostris*) near San Clemente Island, California: a key area for beaked whales and the military? *Marine Biology*, 156:2631-2640.

Falcone, E.A., Schorr, G.S., Watwood, S.L., DeRuiter, S.L., Zerbini, A.N., Andrews, R.D., Morrissey, R.P. and Moretti, D.J. 2017. Diving behaviour of Cuvier's beaked whales exposed to two types of military sonar. *Royal Society Open Science*, 4(8), p.170629.

- Heaney, K.D. and Campbell, R.L., 2016. Three-dimensional parabolic equation modeling of mesoscale eddy deflection. *The Journal of the Acoustical Society of America*, 139(2), pp.918-926.
- Henderson, E.E., C.R. Martin, R.W. Baird, M.A. Kratofil, S.W. Martin, and Southall, B.L. 2021. FY20 Summary Report on the Received Level Analysis of Satellite Tagged Odontocetes. NIWC Pacific Technical Report.
- Henderson, E.E., M.A. Kratofil, R.W. Baird, C.R. Martin, A.E. Harnish, S.W. Martin, and Southall, B.L.. 2024. Assessing exposure and response of satellite-tagged Blainville's beaked whales on the Pacific Missile Range, Hawai'i. NIWC Pacific Technical Report?
- Hewitt, J., Gelfand, A.E., Quick, N.J., Cioffi, W.R., Southall, B.L., DeRuiter, S.L. and Schick, R.S., 2022. Kernel density estimation of conditional distributions to detect responses in satellite tag data. *Animal Biotelemetry*, 10(1), p.28.
- Jeanniard-du-Dot, T., Holland, K., Schorr, G.S. and Vo, D., 2017. Motes enhance data recovery from satellite-relayed biologgers and can facilitate collaborative research into marine habitat utilization. *Animal Biotelemetry*, 5, pp.1-15.
- Johnson, D.S., and J.M. London. 2018. Crawl: An R package for fitting continuous-time correlated random walk models to animal movement data. Zenodo doi:10.5281/zenodo.596464
- Johnson, D.S., J.M. London, M.-A. Lea, and J.W. Durban. 2008. Continuous-time correlated random walk model for animal telemetry data. *Ecology* 89:1208-1215 doi:10.1890/07-1032.1
- Joyce, T.W., Durban, J.W., Claridge, D.E., Dunn, C.A., Hickmott, L.S., Fearnbach, H., Dolan, K. and Moretti, D., 2020. Behavioral responses of satellite tracked Blainville's beaked whales (*Mesoplodon densirostris*) to mid-frequency active sonar. *Marine Mammal Science*, 36(1), pp.29-46.
- Kranstauber, B., A. Cameron, R. Weinzierl, T. Fountain, S. Tilak, M. Wikelski, and R. Kays. 2011. The Movebank data model for animal tracking. *Environmental Modelling & Software* 26(6):834-835 doi:10.1016/j.envsoft.2010.12.005
- Kratofil, M.A., Harnish, A.E., Mahaffy, S.D., Henderson, E.E., Bradford, A.L., Martin, S.W., Lagerquist, B.A., Palacios, D.M., Oleson, E.M. and Baird, R.W., 2023. Biologically Important Areas II for cetaceans within US and adjacent waters—Hawai'i Region. *Frontiers in Marine Science*, 10, 1053581. <https://doi.org/10.3389/fmars.2023.1053581>.
- Locarnini M, Mishonov A, Baranova O, Boyer T, Zweng M, Garcia H, Reagan Jr, Seidov D, Weathers K, Paver C, Smolyar I. 2018. World Ocean Atlas 2018, Volume 1: Temperature. NOAA Atlas NESDIS 81, 52pp.
- London J. M. 2021. Pathroutr: an r package for (re-)routing paths around barriers (version v0.2.1) (Zenodo). doi: 10.5281/zenodo.4321827
- Mahaffy, S.D., R.W. Baird, A.E. Harnish, T. Cullins, S.H. Stack, J.J. Currie, A.L. Bradford, S.R. Salden, and K.K. Martien. 2023. Identifying social clusters of endangered main Hawaiian Islands false killer whales. *Endangered Species Research* 51:249-268. <https://doi.org/10.3354/esr01258>.

Mahaffy, S.D., R.W. Baird, D.J. McSweeney, D.L. Webster and G.S. Schorr. 2015. High site fidelity, strong associations and long-term bonds: short finned pilot whales off the island of Hawai'i. *Marine Mammal Science* 31(4):1427-1451. <https://doi.org/10.1111/mms/12234>

Martin CR, Henderson EE, Martin SW, Helble TA, Alongi GC, Guazzo RA (2023) FY22 annual report on Pacific Missile range facility marine mammal monitoring. NIWC Pacific.

Martin, S.W., C.R. Martin, B. Matsuyama, and E.E. Henderson. 2015. Minke whale (*Balaenoptera acutorostrata*) responses to navy training. *Journal of the Acoustical Society of America* 137(5): 2533-2541.

McClintock, B.T., Johnson, D.S., Hooten, M.B., Ver Hoef, J.M. and Morales, J.M., 2014. When to be discrete: the importance of time formulation in understanding animal movement. *Movement ecology*, 2, pp.1-14.

McSweeney, D.J., R.W. Baird and S.D. Mahaffy. 2007. Site fidelity, associations and movements of Cuvier's (*Ziphius cavirostris*) and Blainville's (*Mesoplodon densirostris*) beaked whales off the island of Hawai'i. *Marine Mammal Science* 23(3):666-687. <https://doi.org/10.1111/j.1748-7692.2007.00135.x>

Miller P, Kvadsheim P, Lam F, Tyack PL et al. 2015. First indications that northern bottlenose whales are sensitive to behavioural disturbance from anthropogenic noise. *R Soc Open Sci* 2:140484

NMFS 2008. Taking and importing of marine mammals; U.S. Navy training in the Southern California Range Complex; proposed rule. 36 p.

NOAA National Geophysical Data Center. 2003. U.S. Coastal Relief Model Vol.7 - Central Pacific. NOAA National Centers for Environmental Information. <https://doi.org/10.7289/V50Z7152>

Owen, K., R.D. Andrews, R.W. Baird, G.S. Schorr and D.L. Webster. 2019. Lunar cycles influence the diving behavior and habitat use of short-finned pilot whales around the main Hawaiian Islands. *Marine Ecology Progress Series* 629:193-206. <https://doi.org/10.3354/meps13123>

Schick, R.S., Bowers, M., DeRuiter, S., Friedlaender, A., Joseph, J., Margolina, T., Nowacek, D.P. and Southall, B.L., 2019. Accounting for positional uncertainty when modeling received levels for tagged cetaceans exposed to sonar. *Nueva Sociedad*, (284), pp.675-690.

Schorr, G.S., Falcone, E.A., Rone, B.K. and Keene, E.L., 2019. Distribution and demographics of Cuvier's beaked whales and fin whales in the Southern California Bight. *Contract*, 66604(18-Q), p.2187.

Schorr, G.S., R.W. Baird, M.B. Hanson, D.L. Webster, D.J. McSweeney and R.D. Andrews. 2009. Movements of satellite-tagged Blainville's beaked whales off the island of Hawai'i. *Endangered Species Research* 10:203-213.

Shaff, J.F., and R.W. Baird. 2021. Diel and lunar variation in diving behavior of rough-toothed dolphins (*Steno bredanensis*) off Kaua'i, Hawai'i. *Marine Mammal Science* 37(4):1261-1276. <https://doi.org/10.1111/mms.12811>.

Širović, A., Wiggins, S.M. and Oleson, E.M., 2013. Ocean noise in the tropical and subtropical Pacific Ocean. *The Journal of the Acoustical Society of America*, 134(4), pp.2681-2689.

Thorne, L.H., Johnston, D.W., Urban, D.L., Tyne, J., Bejder, L., Baird, R.W., Yin, S., Rickards, S.H., Deakos, M.H., Mobley Jr, J.R. and Pack, A.A., 2012. Predictive modeling of spinner dolphin (*Stenella longirostris*) resting habitat in the main Hawaiian Islands. p. e43167.

Van Cise, A. M., K. K. Martien, S. D. Mahaffy, R. W. Baird, D. L. Webster, J. H. Fowler, E. M. Oleson, P. A. Morin. 2017. Familial social structure and socially driven genetic differentiation in Hawaiian short-finned pilot whales. *Molecular Ecology* 26(23):6730-6741. <https://doi.org/10.1111/mec.14397>.

Van Cise, A.M., Baird, R.W., Harnish, A.E., Currie, J.J., Stack, S.H., Cullins, T. and Gorgone, A.M., 2021. Mark-recapture estimates suggest declines in abundance of common bottlenose dolphin stocks in the main Hawaiian Islands. *Endangered Species Research*, 45, pp.37-53.

West, K.L., W.A. Walker, R.W. Baird, D.L. Webster and G.S. Schorr. 2018. Stomach contents and diel diving behavior of melon-headed whales (*Peponocephala electra*) in Hawaiian waters. *Marine Mammal Science* 34(4):1082-1096. <https://doi.org/10.1111/mms.12507>.

Zweng, M., Reagan, J., Seidov, D., Boyer, T., Locarnini, R., Garcia, H., Mishonov, A., Baranova, O., Paver, C., and Smolyar, I. 2018. *World Ocean Atlas 2018, Volume 2: Salinity*, A. Mishonov Technical Ed.; NOAA Atlas NESDIS, pp. 82, 50.

## APPENDIX

### Additional dive behavior results and movement narratives

This section contains all diving behavior tables and figures for tags/TODs where there is no statistically significant difference between phases, as well as all narratives/figures for tags where there was no exposure. Sections are broken down by species, and are organized with tables first, then dive behavior figures, then narratives, then maps.

#### Short-finned pilot whales

*Table 32 - A comparison of dusk diving parameters from short-finned pilot whales exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.*

Dive parameter per individual	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Dusk dive rate (dives/hour)</i>							
GmTag232	3.08	2.63	2.44	2.71	NA	-	
GmTag233	NA	NA	2.98	2.58	3.26	-	
GmTag234	NA	NA	2.29	2.64	2.52	-	
GmTag235	NA	2.82	NA	NA	2.67	-	
GmTag237	NA	2.68	2.65	NA	2.27	-	
<i>% time in surface periods at dusk</i>							
GmTag232	27.16	23.69	25.80	23.82	NA	-	
GmTag233	NA	NA	31.72	43.46	25.00	-	
GmTag234	NA	NA	31.27	29.26	29.55	-	
GmTag235	NA	40.47	NA	NA	26.52	-	
GmTag237	NA	39.89	44.47	NA	35.33	-	
<i>Median dive depth dusk (m)</i>							
GmTag232	535.50	607.50	591.50	575.50	NA	0.2181	NA
GmTag233	NA	NA	631.50	607.50	607.50	0.8532	NA
GmTag234	NA	NA	623.50	591.50	623.50	0.6797	NA
GmTag235	NA	497.50	NA	NA	551.50	0.1436	NA
GmTag237	NA	179.50	527.50	NA	559.50	0.1388	NA
<i>Median</i>							

Dive parameter per individual	Before	Phase A/ A/B Mixed	Interphase	Phase B/ B/A Mixed	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>dive duration dusk (min)</i>							
GmTag232	14.92	17.28	18.50	16.53	NA	0.1235	NA
GmTag233	NA	NA	13.87	13.27	14.67	0.9682	NA
GmTag234	NA	NA	17.52	15.63	16.57	0.0669	NA
GmTag235	NA	13.97	NA	NA	16.63	0.0875	NA
GmTag237	NA	14.17	13.80	NA	19.10	0.2874	NA




Figure 41 - Top. Boxplot showing dive depths of GmTag236 by SCC Phase and time of day. Bottom. Barplot showing dive rates of GmTag236 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 128.8 dB.



False killer whale

Table 33 - A comparison of dawn diving parameters for false killer whale PcTag074 exposed to MFAS for phases that meet the required coverage cutoff. Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

<b>Dive parameter per individual</b>	<b>Before</b>	<b>Phase A</b>	<b>Interphase</b>	<b>Phase B</b>	<b>After</b>	<b>Kruskal-Wallis Test p-value*</b>	<b>Post-hoc Dunn's test significant pairs</b>
<i>Dawn dive rate (dives/hour)</i>	1.49	0.56	0.00	NA	NA	-	
<i>% time in surface periods at dawn</i>	91.03	92.31	100.0	NA	NA	-	
<i>Median dive depth dawn (m)</i>	143.50	751.50	NA	NA	NA	0.1573	NA
<i>Median dive duration dawn (min)</i>	3.25	8.30	NA	NA	NA	0.1573	NA

Table 34 - A comparison of daytime diving parameters for false killer whale PcTag074 exposed to MFAS for phases that meet the required coverage cutoff. Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

<b>Dive parameter per individual</b>	<b>Before</b>	<b>Phase A</b>	<b>Interphase</b>	<b>Phase B</b>	<b>After</b>	<b>Kruskal-Wallis Test p-value*</b>	<b>Post-hoc Dunn's test significant pairs</b>
<i>Day dive rate (dives/hour)</i>	0.68	0.37	0.34	NA	NA	-	
<i>% time in surface periods at day</i>	96.04	97.26	97.43	NA	NA	-	
<i>Median dive depth day (m)</i>	151.50	137.50	159.50	NA	NA	0.5492	NA
<i>Median dive duration day (min)</i>	3.10	4.08	3.77	NA	NA	0.1740	NA

Table 35 - A comparison of dusk diving parameters for false killer whale PcTag074 exposed to MFAS for phases that meet the required coverage cutoff.

<b>Dive parameter per individual</b>	<b>Before</b>	<b>Phase A</b>	<b>Interphase</b>	<b>Phase B</b>	<b>After</b>	<b>Kruskal-Wallis Test p-value*</b>	<b>Post-hoc Dunn's test significant pairs</b>
<i>Dusk dive rate (dives/hour)</i>	1.84	0.00	2.06	NA	NA	-	
<i>% time in surface periods at dusk</i>	85.47	100.00	84.89	NA	NA	-	
<i>Median dive depth dusk (m)</i>	163.50	NA	303.50	NA	NA	0.6847	NA
<i>Median dive duration dusk (min)</i>	4.10	NA	3.90	NA	NA	0.8075	NA

Table 36 - A comparison of nighttime diving parameters for false killer whale PcTag074 exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

<b>Dive parameter per individual</b>	<b>Before</b>	<b>Phase A</b>	<b>Interphase</b>	<b>Phase B</b>	<b>After</b>	<b>Kruskal-Wallis Test p-value*</b>	<b>Post-hoc Dunn's test significant pairs</b>
<i>Night dive rate (dives/hour)</i>	0.26	0.19	0.16	NA	NA	-	
<i>% time in surface periods at night</i>	98.46	98.20	98.63	NA	NA	-	
<i>Median dive depth night (m)</i>	79.50	129.50	123.50	NA	NA	0.199	NA
<i>Median dive duration night (min)</i>	3.00	5.33	3.80	NA	NA	0.0773	NA




Figure 42 - Top. Boxplot showing dive depths of PcTag074 by SCC Phase and time of day. Bottom. Barplot showing dive rates of PcTag074 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 90.5 dB.

## Melon-headed whales

### *PeTag035, PeTag036*

These individuals were tagged in the same group in 2022. For PeTag035, information is available on movement patterns for the interphase period (0.1 days), the B/A mixed phase (0.2 days), and the after phase (18.1 days). While PeTag035 was tagged on the range, it moved off the range during Phase B, and continued to move southwest away from the range over the remainder of its deployment. Information is available on movement patterns for PeTag036 for the B/A mixed phase (0.2 days) and the after phase (10.1 days).

Dive behavior data for both tags were only transmitted during the after phase, after any potential exposure had occurred, and these tags are hence given no further consideration in the dive behavior analysis.





Figure 43 - Top. Map showing crawl model trackline of PeTag035 during the 2022 August SCC. Bottom. Map showing crawl model trackline of PeTag036 during the 2022 August SCC.

Rough-toothed dolphins

Table 37 - A comparison of dawn diving parameters from rough-toothed dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

Dive parameter per individual	Before	Phase A	Interphase	Phase B	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Dawn dive rate (dives/hour)</i>							
SbTag023	NA	0.00	0.00	0.77	NA	-	
SbTag025	NA	NA	NA	NA	0.38	-	
<i>% time in surface periods at dawn</i>							
SbTag023	NA	100.00	100.00	97.89	NA	-	
SbTag025	NA	NA	NA	NA	98.65	-	
<i>Median dive depth dawn (m)</i>							
SbTag023	NA	NA	NA	85.50	NA	NA	NA
SbTag025	NA	NA	NA	NA	97.50	NA	NA
<i>Median dive duration dawn (min)</i>							
SbTag023	NA	NA	NA	1.65	NA	NA	NA
SbTag025	NA	NA	NA	NA	2.13	NA	NA



Table 38 - A comparison of dusk diving parameters from rough-toothed dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

Dive parameter per individual	Before	Phase A	Interphase	Phase B	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Dusk dive rate (dives/hour)</i>							
SbTag023	NA	1.90	0.28	0.00	NA	-	
SbTag025	NA	NA	NA	NA	0.55	-	
<i>% time in surface periods at dusk</i>							
SbTag023	NA	85.66	98.32	100.00	NA	-	
SbTag025	NA	NA	NA	NA	95.65	-	
<i>Median dive depth dusk (m)</i>							
SbTag023	NA	131.50	147.50	NA	NA	0.6547	NA
SbTag025	NA	NA	NA	NA	139.50	NA	NA
<i>Median dive duration dusk (min)</i>							
SbTag023	NA	5.17	3.67	NA	NA	0.6547	NA
SbTag025	NA	NA	NA	NA	4.77	NA	NA





Figure 44 - Top. Boxplot showing dive depths of SbTag025 by SCC Phase and time of day. Bottom. Barplot showing dive rates of SbTag025 by SCC Phase and time of day.

*SbTag024*

Information was available about *SbTag024*'s movements in 2021 for Before (2.3 days), Phase A (1.7 days), and the interphase period (0.5 days). During this time, *SbTag024* spent most of its time on the range, though the deployment duration did not overlap with any potential MFAS exposures.




Figure 45 - Map showing crawl model trackline of *SbTag024* during the 2021 August SCC.

*SbTag025*

Information was available about SbTag025's movements in 2022 for a very short part of the interphase (0.01 days), the B/A mixed phase (0.2 days), and the after phase (7.3 days). During this time, SbTag025 remained on or near the range. This individual was not exposed to MFAS.

Dive behavior data were available for SbTag025 for only the B/A mixed phase and the after phase. However, when broken down by time of day and phase, only the after phase data met the required coverage (relative to the phase duration) for inclusion in the dive analysis, and hence this tag was given no further consideration in the dive behavior analysis.




Figure 46 - Map showing crawl model trackline of SbTag025 during the 2022 August SCC. This individual was not exposed to Navy activities.

Common bottlenose dolphins

Table 39 - A comparison of dawn diving parameters from common bottlenose dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

Dive parameter per individual	Before	Phase A	Interphase	Phase B	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Dawn dive rate (dives/hour)</i>							
TtTag041	NA	NA	2.92	NA	2.35	-	
<i>% time in surface periods at dawn</i>							
TtTag041	NA	NA	54.69	NA	55.94	-	
<i>Median dive depth dawn (m)</i>							
TtTag041	NA	NA	591.50	NA	615.50	0.4366	NA
<i>Median dive duration dawn (min)</i>							
TtTag041	NA	NA	10.30	NA	11.10	0.0933	NA

Table 40 - A comparison of dusk diving parameters from common bottlenose dolphins exposed to MFAS for phases that meet the required coverage cutoff. Kruskal-Wallis one-way ANOVA significant results (i.e., significant differences among phases were detected) are shown in bold. Pairs of phases where significant differences were detected are listed in the associated post-hoc Dunn's test column (level of significance 0.05). Values for dive rates and percentage time in surface periods represent single values for each individual for each period, thus no statistical testing was undertaken.

Dive parameter per individual	Before	Phase A	Interphase	Phase B	After	Kruskal-Wallis Test p-value*	Post-hoc Dunn's test significant pairs
<i>Dusk dive rate (dives/hour)</i>							
TtTag039	NA	NA	2.47	NA	NA	-	
TtTag041	NA	NA	2.42	NA	2.91	-	
<i>% time in surface periods at dusk</i>							
TtTag039	NA	NA	74.61	NA	NA	-	
TtTag041	NA	NA	53.35	NA	51.89	-	
<i>Median dive depth dusk (m)</i>							
TtTag039	NA	NA	327.50	NA	NA	NA	NA
TtTag041	NA	NA	719.50	NA	647.50	0.0515	NA
<i>Median dive duration dusk (min)</i>							
TtTag039	NA	NA	6.50	NA	NA	NA	NA
TtTag041	NA	NA	11.55	NA	10.53	0.0707	NA





Figure 47 - Top. Boxplot showing dive depths of TtTag041 by SCC Phase and time of day. Bottom. Barplot showing dive rates of TtTag041 by SCC Phase and time of day. Maximum estimated RL from MFAS for this individual was 100.5 dB.

### *TtTag041*

Information was available about TtTag041's movements in 2022 for the interphase period (3.0 days), the B/A mixed phase (0.2 days), and the after phase (13.8 days). TtTag041 was deployed on the range during the interphase period, then moved to the east side of Kaua'i where it spent the duration of the B/A mixed phase, then continued to circumnavigate the island by moving southwest and back onto the range.

Dive behavior data were available for TtTag041 for the interphase period, the B/A mixed phase, and the after phase. However, when broken down by time of day and phase, not every time of day and phase met the required coverage (relative to the phase duration) for inclusion in the analysis. Coverage was sufficient for the interphase and after phase for each time of day, as well as for daytime during the B/A mixed phase.

The dawn dive rate was highest during the interphase, and dropped slightly in the after phase, while the percentage of surface time during dawn hours remained fairly consistent between the two phases. There was no statistically significant variation between phases for dawn dive depths or durations.

The day dive rate was highest during the B/A mixed phase, and lowest during the after phase, while the percentage of surface time during day hours remained fairly consistent between phases. Day dives were significantly shallower and shorter during the B/A mixed phase than during either the interphase or after phase.

The dusk dive rate was highest during the after phase, and the percentage of surface time during dusk hours remained fairly consistent between the two phases. There was no statistically significant variation between phases for dusk dive depths or durations.

The night dive rate was highest during the interphase, and the percentage of surface time during night hours rose slightly during the after phase. There was no statistically significant variation in night dive depth between phases, but night dives were significantly shorter during the after phase than during the interphase.






Figure 48 - Map showing crawl model trackline of TtTag041 during the 2022 August SCC. This individual was not exposed to Navy activities.