

SPLASH: Structure of Populations, Levels of Abundance and Status of Humpback Whales in the North Pacific

Final report for Contract AB133F-03-RP-00078

Cascadia Research
Waterstreet Building
218½ W 4th Avenue
Olympia, Washington 98501
www.cascadiaresearch.org

For

U.S. Dept of Commerce
Western Administrative Center
Seattle, Washington

by

John Calambokidis, Erin A. Falcone, Terrance J. Quinn, Alexander M. Burdin, Phillip J. Clapham, John K.B. Ford, Christine M. Gabriele, Richard LeDuc, David Mattila, Lorenzo Rojas-Bracho, Janice M. Straley, Barbara L. Taylor, Jorge Urbán R., David Weller, Briana H. Witteveen, Manami Yamaguchi, Andrea Bendlin, Dominique Camacho, Kiirsten Flynn, Andrea Havron, Jessica Huggins, and Nora Maloney

Contracting Technical Representatives

Jay Barlow and Paul R. Wade

May 2008

EXECUTIVE SUMMARY

Humpback whales were hunted commercially in the North Pacific until 1966 and remain on the endangered species list at the time of this report. The degree to which they have recovered from whaling in the North Pacific is difficult to determine because of the lack of accurate abundance estimates for this wide-ranging species. New methods such as photographic identification and analysis of skin and blubber biopsy samples have provided tools to examine the status of humpback whales. These animals undergo one of the longest migrations of any mammal and, within the North Pacific, their population structure and migrations appear to be complex. To effectively study and protect a species that travels widely across international borders requires a high level of collaboration among researchers and governments.

SPLASH (Structure of Populations, Levels of Abundance and Status of Humpbacks) represents one of the largest international collaborative studies of any whale population ever conducted. It was designed to determine the abundance, trends, movements, and population structure of humpback whales throughout the North Pacific and to examine human impacts on this population. This study involved over 50 research groups and more than 400 researchers in 10 countries. It was supported by a number of agencies and organizations including the National Marine Fisheries Service, the National Marine Sanctuary Program, National Fish and Wildlife Foundation, Pacific Life Foundation, Department of Fisheries and Oceans Canada, and Commission for Environmental Cooperation with additional support from a number of other organizations and governments for effort in specific regions. Results presented here include a comprehensive analysis of individual identification photographs. Additional analysis of human impacts, ecosystem markers (e.g., stable isotopes) and the genetic structure of populations are underway or planned pending further funding.

Field efforts were conducted on all known winter breeding regions for humpback whales in the North Pacific during three seasons (2004, 2005, 2006) and all known summer feeding areas during two seasons (2004, 2005). A total of 18,469 quality fluke identification photographs were taken during over 27,000 approaches of humpback whales. After reconciling all within and cross-regional matches (from both the primary match and rechecks), a total of 7,971 unique individuals were cataloged in SPLASH. A total of 6,178 tissue samples were also collected for genetic studies of population structure, with fairly even representation of wintering and feeding areas.

Migratory movements and population structure of humpback whales in the North Pacific were found to be more complex than had been previously described; a high degree of structure, however, was also apparent. Migrations between feeding and wintering areas were documented based on 873 whales that were seen on both a wintering and feeding areas. The overall pattern showed that coastal wintering regions of the western (Asia) and eastern (mainland Mexico and Central America) North Pacific were the primary wintering areas for the lower latitude coastal feeding regions. The wintering areas off Hawaii and the Revillagigedo Archipelago were the primary wintering regions for the more central and northern latitude feeding areas.

Even though the SPLASH study collected data from all known wintering and feeding areas for humpback whales in the North Pacific, the SPLASH data did suggest the likely existence of missing wintering areas that have not been previously described. Humpback whales

that feed off the Aleutians and in the Bering Sea were not well represented on any of the sampled wintering areas and must be going to one or more unsampled winter locations. Thus, it is likely that SPLASH has revealed a new breeding ground for humpback whales. While it would be logical to assume that this region would be located in the eastern central North Pacific, the complexities of the migratory pathways revealed here indicate that this is not certain.

Individual whales showed high rates of return to specific wintering and feeding areas, suggesting strong site fidelity to both habitats. Interchange of whales between feeding areas both within and between seasons was unusual and all but a few of these were between adjacent areas. Similarly, whales tended to return to the wintering region they had inhabited previously, although the geographic scale of this varied by region. Whales showed frequent interchange among areas within the Hawaiian Islands but only rarely switched between broader regions. Some wintering areas that were sampled, especially Ogasawara and Baja Mexico, appeared to be transitory areas rather than migratory destinations. These findings are consistent with preliminary analyses of the genetic structure population showing a high degree of maternally-directed fidelity to both breeding and feeding grounds but a complex relationship between seasonal habitats.

Using several methods, the abundance of humpback whales was estimated to be just under 20,000 for the entire North Pacific, an estimate that is about double estimates made previously. The non-stratified Chapman-Petersen estimates of abundance were 18,000 to 21,000. Among geographically stratified models, the model assuming non-Markovian movements with capture probability proportional to sample size across years provided the best overall fit to the data indicated an abundance of 17,558 for wintering areas and 19,056 for the feeding areas. The average of these two estimates (18,302) represented the best estimate of overall abundance of humpback whales in the North Pacific, excluding calves. Over 50% of this population was estimated to winter in Hawaiian waters with large populations also inhabiting Mexican waters. The abundance estimates of humpback whales wintering in Asia and Central America were fairly low (1,000 or less). Among feeding areas, regional estimates differed greatly among models. Average estimates of abundance ranged from about 100-700 for Russia, 6,000-14,000 for the Bering Sea and Aleutians, 3,000-5,000 each for the Gulf of Alaska and the combined Southeast Alaska and Northern British Columbia area, 200-400 for Southern British Columbia-Northern Washington, and 1,400-1,700 for California-Oregon.

The SPLASH estimate represents a dramatic increase in abundance from other post-whaling estimates for the overall North Pacific, yet is consistent with a moderate rate of recovery for a depleted population. Comparison of the SPLASH estimate of 18,302 for all feeding and wintering areas to the estimate of 9,819 obtained for 1991-93 in a previous study suggests a 4.9% annual increase over this 13-year period. Going back to the estimate of 1,400 whales at the end of whaling for humpbacks in 1966, a 6.8% annual increase over the 39-year period would be required to reach the current SPLASH abundance. For Hawaii, three methods were used to compare estimates to determine trends since the early 1990s and yielded very similar annual rate of increase from 5.5 to 6.0%.

While the overall humpback whale abundance and trends in the North Pacific are encouraging, some areas should be of concern, especially Asia. The western-most feeding and wintering areas were distinct from the rest of the North Pacific with a very low level of interchange between Asian wintering or feeding areas and those in the central and eastern North

Pacific. Abundance estimates in this area are low (below historical levels based on the number taken in this region) and whales along the Asian coast appear to be subject to a high level of incidental mortality.

INTRODUCTION

Humpback whale (*Megaptera novaeangliae*) populations were depleted due to commercial exploitation and are listed as endangered under the Endangered Species Act today. North Pacific humpback whale populations were thought to have numbered about 15,000 prior to commercial exploitation in the twentieth century (Rice 1978), however, this estimate was based on whaling data that may have been inaccurate. Approximate numbers in the North Pacific following the cessation of commercial whaling have been estimated at 1,400 (Gambell 1976) and 1,200 (Johnson and Wolman 1984).

The most recent estimate of North Pacific humpback whale abundance was conducted using capture-recapture statistics with photo-identification data from the early 1990s (Calambokidis *et al.* 1997, 2001, In prep.). This study was a retrospective analysis using data collected between 1990 and 1993 by 16 research groups from all areas of the North Pacific where photo-identification studies had been conducted. It yielded estimates of 6,000- 10,000 whales. This estimate is now 10 years old and a number of areas where whales are now known to be present were not represented in the study. Genetic data derived from biopsy samples also were not part of that study. Data from photo-identification and genetic studies have provided some information on North Pacific stock structure, verifying a high degree of site fidelity to feeding areas and the presence of individuals from multiple feeding areas at each wintering area but few individuals that move between wintering area; however, only limited data exist on the numbers, sizes, and ranges of most feeding areas in the North Pacific (Baker *et al.* 1986, 1993, 1994, 1998; Calambokidis *et al.* 1996, 2001).

We report on the first-ever comprehensive field study of humpback whales throughout the North Pacific. Termed SPLASH (Structure of Populations, Levels of Abundance and Status of Humpbacks), this work was the result of an international collaborative research effort conducted throughout the North Pacific which involved over 50 research groups and more than 400 researchers, and which was supported by a number of agencies and organizations. Primary support for the overall project and specifically for data collection in three of the five field seasons (Winter 2004, Summer 2004, and Summer 2005) came under a contract from NOAA's National Marine Fisheries Service and the National Marine Sanctuary Program. Major support including that to help fund the sampling in Winter 2005 and Winter 2006 came from the National Fish and Wildlife Foundation, Pacific Life Foundation, Department of Fisheries and Oceans Canada, and Commission for Environmental Cooperation. Significant support also came from a number of other organizations and governments for effort in specific regions.

OVERALL OBJECTIVES

A dedicated sampling program and subsequent analyses of humpback whales at wintering and feeding areas within the North Pacific were conducted to address the following objectives:

- Collect a representative sample of photo-identification photographs of humpback whale populations throughout the North Pacific.

- Collect biopsy samples to provide a better understanding of population structure and migratory interchange using genetic markers.
- Estimate humpback whale overall abundance for the North Pacific basin using capture-recapture models.
- Estimate the abundance of humpback whales for specific wintering and feeding areas.
- Examine trends in abundance.

Additionally, data were gathered to examine human impacts on humpback whales including incidence of entanglement and ship strikes and other population parameters including reproductive rates, mortality rates, age/sex structure, pregnancy rates.

METHODS AND STUDY DESIGN

SPLASH sampling was conducted by an international collaborative group of more than 50 research groups and 400 researchers coordinated by a Steering Committee that included coordinators for each of the regions (Table 1) sampled, as well as principals in the funding, coordination, and analysis of SPLASH.

Table 1. Summary of organizations coordinating SPLASH-dedicated surveys and compiling opportunistic data contributions.

Research Group	Full Name	Survey Regions
Feeding areas		
ASLC	Alaska Sealife Center	Russia
CRC	Cascadia Research	U.S. West Coast, British Columbia, Bering Sea
DFOC	Department of Fisheries and Oceans, Canada	British Columbia
GBNP	Glacier Bay National Park	Southeast Alaska
NGOS	North Gulf Oceanic Society	Northern Gulf of Alaska, Bering Sea
NMML	U.S. National Marine Mammal Laboratory	Southeast Alaska, Northern Gulf of Alaska, Western Gulf of Alaska, Bering Sea
SWFSC	Southwest Fisheries Science Center	SPLASH cruise of BC, SEAK, Gulf of Alaska, Aleutians and Bering Sea in 2004 and US West Coast in 2005 (CSCAPE)
UAFK	University of Alaska Fairbanks	N Gulf of Alaska, W Gulf of Alaska
UASE	University of Alaska Southeast	Southeast Alaska
Wintering areas		
OMC	Ogasawara Marine Center	Asia including Philippines
HIWS	Hawaiian Islands Humpback Whale National Marine Sanctuary.	Hawaii
UABCS	Universidad Autonoma de Baja California Sur	Mexico
CRC	Cascadia Research	Central America

Sampling effort was designed to obtain a large and broadly distributed sample from each geographic region including both wintering and feeding areas and was collected in a manner to best achieve random sampling of all age and sex classes. Somewhat different sampling procedures were used on feeding versus wintering areas, particularly the type of research vessel used for sampling area. Data were collected during three seasons at wintering areas (2004, 2005, and 2006) and in two seasons (2004 and 2005) at feeding areas. In all regions, the goal was to apportion effort in a manner that was proportional to the anticipated density of animals. The study was designed to provide broad coverage of all known feeding areas (Figure 1) and wintering areas of humpback whales and within each area to sample as wide a geographic area as possible. Sampling within areas was conducted over a broad time period, especially in the wintering areas, to cover the full duration of the season (Table 2). This approach helped to avoid sampling bias regarding residency time and timing of arrival on the wintering regions.




Figure 1. Survey track lines and effort during surveys of feeding areas in 2004 and 2005.

Table 2. Summary of survey effort broken down by season and area.

Region	Start Date	End Date	Vessel Days	Research Groups	Fluke Identifications	Fluke Ident/SPLASH ID	Unique Individ.	Samples
Winter 2004								
Asia-PHI	26-Feb-04	01-May-04	33	WWFP	56	50	27	3
Asia-OK	17-Feb-04	05-Mar-04	12	OCA	85	72	43	0
Asia-OG	10-Jan-04	13-May-04	47	OMC	294	205	114	45
Hawaii	02-Dec-03	12-May-04	174	CFWS, HIWS, HMMC, HWRF, MMRC, OWSI, TDI, WHTR	1170	839	697	540
MX-REV	24-Jan-04	16-Apr-04	169	COR, HSU, UNAM	1352	1206	317	151
MX-Baja	20-Jan-04	03-Apr-04	60	UABCS, UNAM	215	183	182	176
MX-ML	30-Nov-03	08-Apr-04	226	COVISI, FIBB, UNAM	527	417	223	77
Cent Am	14-Jan-04	22-Mar-04	41	CRC	27	23	18	12
Total					3,726	2,995	1,621	1,004
Summer 2004								
Russia	18-Jul-04	19-Aug-04	16	ASLC	65	57	40	30
W Aleut.	16-Aug-04	22-Aug-04	8	SWFSC	18	15	12	9
E Aleut.	21-Jul-04	15-Sep-04	20	NMML, SWFSC	76	66	51	38
Bering	09-Jun-04	10-Sep-04	62	NGOS, NMML, SWFSC	415	302	228	106
WGOA	21-Jul-04	19-Sep-04	29	NMML, SWFSC, UAFK	400	334	223	120
NGOA	07-May-04	08-Oct-04	209	NGOS, UAFK, NMML, SWFSC	1359	1174	730	247
SEAK	01-May-04	14-Dec-04	246	CRC, GBNP, NMML, SWFSC, UASE, UAFB	1933	1566	808	347
NBC	01-Jan-04	27-Nov-04	254	DFOC, SWFSC	1050	868	421	106
NWA-SBC	18-Apr-04	23-Nov-04	46	CRC, DFOC, SWFSC	171	106	76	26
CA-OR	04-Apr-04	13-Dec-04	108	CRC	598	421	253	65
Total					6,085	4,909	2,842	1,094
Winter 2005								
Asia-PHI	17-Feb-05	02-May-05	60	WWFP	86	65	35	6
Asia-OK	18-Feb-05	25-Mar-05	20	OCA	122	114	55	0
Asia-OG	27-Dec-04	13-May-05	53	OMC	524	292	123	60
Hawaii	15-Dec-04	29-Apr-05	212	ANZO, CFWS, HAMR, HIWS, HMMC, HWRF, MMRC, OWSI, TDI, WHTR	1718	1102	846	668
MX-REV	08-Feb-05	22-Apr-05	132	COR, HSU, UNAM	781	568	193	123
MX-Baja	08-Jan-05	30-May-05	103	CRC, UABCS	382	198	157	97
MX-ML	16-Nov-04	16-Mar-05	261	COVISI, CRC, FIBB, UNAM	741	462	266	150
Cent Am	17-Jan-05	18-Mar-05	54	CRC	82	69	48	15
Total					4,436	2,870	1,723	1,119
Summer 2005								
Russia	22-Jun-05	14-Aug-05	44	ASLC	101	98	72	43
Bering	25-Jun-05	10-Sep-05	39	CRC, NGOS, NMML	755	475	301	164
WGOA	31-May-05	11-Sep-05	15	NMML, UAFK	284	205	111	81
NGOA	09-Jan-05	25-May-06	183	NGOS, NMML, UAFK	1117	755	427	292
SEAK	09-Jan-05	11-Jan-06	167	CRC, GBNP, NMML, UASE	1545	1096	482	145
NBC	24-Jan-05	26-Jan-06	236	DFOC	826	608	236	56
NWA-SBC	25-Feb-05	27-Dec-05	169	CRC, DFOC, SWFSC	424	329	152	33
CA-OR	08-Mar-05	05-Dec-05	221	CRC, SWFSC	848	680	319	74
Total					5,900	4,246	2,100	888
Winter 2006								
Asia-PHI	19-Feb-06	30-Apr-06	79	WWFP	88	47	26	0
Asia-OK	12-Jan-06	30-Mar-06	95	OCA	633	424	152	94
Asia-OG	27-Nov-05	24-May-06	63	OMC	775	285	119	259
Hawaii	14-Nov-05	30-Apr-06	321	ANZO, CFWS, HAMR, HIWS, HMMC, HWRF, MMRC, OWSI, TDI, WHTR	3590	1488	1026	1174
MX-REV	30-Jan-06	22-Apr-06	159	COR, HSU, UNAM	929	532	186	198
MX-Baja	10-Dec-05	17-May-06	52	UABCS	198	120	87	121
MX-ML	09-Dec-05	31-May-06	195	COVISI, FIBB, UABCS, UNAM	777	475	328	204
Cent Am	05-Dec-05	19-Mar-06	55	CRC	101	78	46	23
Total					7,091	3,449	1,970	2,073
All seasons total					27,238	18,469	10,256	6,178
Unique after internal matches							7,971	

Data collection methods

Photo-identification

Photographs of pigmentation patterns and scarring on the ventral surface of tailflukes, together with serration patterns along the trailing edge (Figure 1), were used to individually identify whales (*e.g.*, Katona *et al.* 1979). To obtain photographs, whales were photographed with digital SLR cameras equipped with telephoto lenses.

Biopsy sampling

Skin and attached blubber tissue samples were collected for genetic analysis using a small stainless steel biopsy dart fired from a crossbow or modified rifle or air-powered gun. Each dart was fitted with a flange or “stop” that regulated penetration of the bolt/dart and caused recoil after sampling. Flotation material secured to the shaft of the bolt/dart allowed it to float on the surface and be retrieved after sampling. Crossbows, most commonly with a draw of 68 kg (150 lbs), and veterinary rifles using either compressed air or blank charges with adjustable pressure were used for sample collection. Depending on field conditions, samples were preserved by freezing immediately after sampling or by immersion in a saturated solution of salt (many samples were also stored in ethanol if freezing was not available). At least half of each biopsy tissue sample (skin and attached blubber if blubber is obtained in the sample) was submitted to the marine mammal tissue archive at the Southwest Fisheries Science Center. An initial analysis of population structure using a representative subset of 2,000 samples is underway with funding from National Fisheries and Wildlife Foundation and the Marine Mammal Institute of Oregon State University.

Regional sampling effort

A variety of approaches were used in the SPLASH sampling summarized in Table 2 and are briefly described by region below. Regional strata were assigned at the outset, based on areas of effort and concentrations of sightings and then modified or pooled based on preliminary results as described in the text. Region names and abbreviations and how they were pooled for the mark-recapture analyses are listed in Table 3.

Asia

The winter distribution of humpback whales in the western North Pacific is centered off the Ogasawara Islands, Ryukyu (Okinawa) Islands, Taiwan, the Philippines, and the Mariana Islands. Humpback whales in this geographic region are distributed over a large area along this chain of islands. Past photographic identification has been conducted in this region began in the 1990s (Darling and Mori 1993, Yamaguchi *et al.* 2002, Acebes 2001, Acebes *et al.* 2007). SPLASH effort in this region was coordinated by the Ogasawara Marine Center (Manami Yamaguchi). Sampling was conducted in the following areas: 1) Ogasawara including Chichi-

jima, Haha-jima (50km from Chichi-jima) and Muko-jima (70km from Chichi-jima) by OMC, 2) Okinawa including Okinawa mainland and Zamami Islands (40km from Okinawa mainland) by the Okinawa Expo Aquarium, and 3) Philippines around the Babuyan Islands by World Wildlife Fund-Philippines (Acebes *et al.* 2007). Effort was conducted primarily from shore-based small boats.

Table 3. Summary of area designations and abbreviations used in SPLASH.

Area (subarea)	Abbreviation	6x6 model abbrev.
Russia (pooled regions)		
Kamchatka	Kamchatka	Russia-Kam
Commander Islands	Commander Is	Al-Ber
Gulf of Anadyr		Al-Ber
Aleutian Islands	Aleut	Al-Ber
Western Aleutians	W Aleut	Al-Ber
Eastern Aleutians	E Aleut	Al-Ber
Bering Sea	Bering	Al-Ber
Gulf of Alaska	GOA	GOA
Western Gulf of Alaska	WGOA	GOA
Northern Gulf of Alaska	NGOA	GOA
Southeast Alaska	SEAK	SEAK-NBC
Northern British Columbia	NBC	SEAK-NBC
Southern British Columbia-	SBC-NWA	SBC-NWA
Northern Washington		
California-Oregon	CA-OR	CA-OR
Asia	Asia	Asia
Philippines	Asia-PHI	Asia
Japan-Okinawa	Asia-OK	Asia
Japan-Ogasawara	Asia-OG	Asia
Hawaii	HI	HI
Kauai	Kauai	HI
Oahu	Oahu	HI
Penguin Bank	PB	HI
Molokai	Molokai	HI
Maui	Maui	HI
Big Island	Big Is	HI
Mexico	MX	
Mainland	MX-ML	Mx-Main
Baja	MX-Baja	Baja
Revillagigedo	MX-REV	Mx-Rev
Central America	Cent Am	CentAm

Hawaii

Sampling for the SPLASH project in Hawaii was coordinated and funded by NOAA's Hawaiian Islands Humpback Whale National Marine Sanctuary (Dave Mattila and Alan Ligon). The Sanctuary and its partner, the State of Hawaii, contracted local humpback whale researchers, who have been consistently active in the region, to collect geographically and temporally representative data for SPLASH. These represent eight teams: on Kauai (1), Oahu (1), Penguin Bank (1), Maui (4) and the Big Island (1).

Mexico

There are three main winter aggregations in the Mexican Pacific: the southern end of Baja California Peninsula (Baja); central portions of the Pacific coast of mainland Mexico (Mainland); and the Revillagigedo Archipelago (Revillagigedo)(Urban and Aguayo, 1987). Jorge Urbán was the regional coordinator for the field sampling in Mexico. Field sampling in Baja was primarily conducted by UABCS coordinated by Ursula Gonzalez, off mainland Mexico by UNAM and UABCS and off the Revillagigedos by Jeff Jacobsen and included effort by personnel associated with Humboldt State University, Cornell University, and UNAM. The primary platforms for dedicated effort were inflatable boats at Revillagigedo and pangas at mainland and Baja with 2-4 people, with additional effort as part of broader surveys from larger platforms. Most of the effort off mainland Mexico was conducted around Bahia de Banderas with some surveys conducted near the area around the Islas Tres Marias. An important part of the effort in this region was the contribution of two groups who compiled identification photographs obtained opportunistically during whale watching operations in and around Bahia de Banderas. These were COVISI (coordinated by Maria Eugenia Rodriguez and Eduardo Lugo of Wildlife Connections) and FIBB (coordinated by Astrid Frisch of Ecotours de Mexico).

Central America

The waters from southern Mexico south along the Central America coast are used as a wintering area for humpback whales coming almost exclusively from feeding areas off California (Steiger *et al.* 1991; Calambokidis *et al.* 2000; Rasmussen *et al.* 2001). Cascadia Research coordinated surveys in this region. To cover this broad low-density area, surveys were conducted in a number of ways: 1) dedicated surveys were made from small chartered boats, 2) a network of local collaborators were set up to obtain opportunistic identifications, and 3) identification photographs were collected during several weeks of surveys in collaboration with Oceanic Society Expeditions trips in southern Costa Rica. Dedicated small-boat surveys were conducted in Guatemala, El Salvador, Nicaragua, and Costa Rica.

US West Coast including California, Oregon, and Washington

Surveys conducted along a broad area of the U.S. west coast were coordinated by Cascadia Research and SWFSC using several platforms and with additional contributions from a number of sources included:

1. Dedicated small boat surveys using 5-6m RHIBs based from various harbors all along the coast.
2. CSCAPE cruise conducted by SWFSC in 2005 off the coast of California, Oregon, and Washington.
3. Opportunistic identifications obtained by naturalists, researchers, and boaters including the Naturalist Corps working with the Channel Islands National Marine Sanctuary.
4. SPLASH sampling was conducted in June based from a NOAA ship and deployed RHIB during surveys off Washington conducted with the Olympic Coast National Marine Sanctuary (OCNMS).

Washington and southern British Columbia were treated as a single region because the concentration of whales straddles the border and the same general areas were being sampled from effort originating on either side of the border.

British Columbia

DFO coordinated sampling effort in British Columbia which included small-boat surveys, ship surveys and opportunistic identifications. A joint DFO and Cascadia survey aboard the charter vessel *Curve of Time* with two deployed RHIBs was conducted off northern Vancouver Island and off the Queen Charlotte Islands each August. Additional identifications were obtained as a part of ship surveys conducted by DFO generally in spring and fall of each year. Small-boat effort was conducted off northern British Columbia from a variety of vessels that operated widely along the mainland coast or were based at Langara Island. SPLASH effort off southern British Columbia was conducted from a small charter boat operating along the southwest side of Vancouver Island. Surveys were conducted in West Hecate Strait in collaborations with Parks Canada.

Southeast Alaska – coastal effort

Three primary research groups collected data in SPLASH in Southeast Alaska: University of Alaska Southeast, Glacier Bay National Park and Preserve and National Marine Mammal Laboratory. UASE vessel charters worked the inside waters of Peril Straits, Lynn Canal, Chatham Strait, Stephens Passage, Frederick Sound and the outside near-shore waters from Dixon Entrance to Cross Sound. Researchers from the Glacier Bay National Park and Preserve surveyed Glacier Bay and Icy Strait. The NOAA vessel *John Cobb* surveyed the inside waters of southern Southeast Alaska.

Northern and Western GOA

Coastal effort in the Gulf of Alaska and Aleutians was conducted through a subcontract to the North Gulf Oceanic Survey (NGOS, Craig Matkin) and coordinated by Kate Wynne and Bree Witteveen (UAFK). Sampling was conducted successfully in a number of areas including 1) Prince William Sound and Kenai area, 2) Barren Islands, 3) Kodiak Is. area, 4) Shumagin Islands, and 5) eastern Aleutian Islands.

Russia

Surveys for humpback whales on feeding grounds off Russia were conducted primarily based from a larger charter vessel and smaller deployed boats under subcontract to North Pacific Wildlife Consulting LLC (Alexander Burdin). The primary survey areas included Anadyr Gulf, Bering Island, Litke Strate, Karaginsky Island, Dezhnevia Bay and the Commander Islands.

NMML ship surveys in Alaska waters

The National Marine Mammal Laboratory (NMML) conducted a range of large-scale ship surveys using the NOAA vessel *Oscar Dyson* and other platforms in the waters off Alaska especially in 2005. These covered broad areas of the Gulf of Alaska, Aleutian Islands, and Bering Sea including offshore areas missed by the coastal-based work described above.

SWFSC ship surveys - 2004

A large scale ship survey was conducted in 2004 by SWFSC that encompassed waters off British Columbia, SE Alaska, Gulf of Alaska, Aleutians, and the Bering Sea. This survey provided somewhat systematic coverage of broad portions of the feeding areas of humpback whales, especially in offshore waters.

Management of regional submissions

At the end of each field season (or incrementally throughout the field season), regional coordinators received copies of all photographs, datasheets, vessel track lines, and tissue samples collected. In some cases, the contributors entered their data into a digital format and provided this to the regional coordinator. Additionally, some contributors reconciled their photographs and provided a set of best flukes for each unique individual that they encountered that season as well as a complete archive of all photographs taken.

Regional coordinators were responsible for the compilation of regional data from a season into a unified database including effort, sighting, identification, and sample data fields requested prior to submission to Cascadia Research. Data were either entered from hard copies of the field notes provided by the contributors, or imported from digital formats. Regional coordinators then reviewed photograph archives from each sighting to identify (or verify) selections of the best photos (fluke, left and right flanks, and tailstocks) of each individual, and these filenames were imported into the database. In most cases, where software was available, a selection of sighting data was also imported into the metadata fields associated with each image file.

The best fluke photographs from each sighting within an area were compared to each other. Regional coordinators included low quality fluke photos in this preliminary match stage to identify as many within area resightings of individual whales as possible. At the completion of regional reconciliation, a copy of the best fluke photograph of each whale for that season was

copied into a separate folder, and an area-season working ID number was assigned to each. This number was assigned to all identifications of the whale within the area, and was also updated in database records where the identity of the biopsy sampled whale was known within a sighting. A digital copy of this preliminary catalog was provided to the central matching office, along with the combined data for all contributors from the area. Tissue samples were sent separately to the lab at SWFSC for archiving, extraction, and/or distribution.

The seasonal collection from each area, including archives of all photographs collected, all vessel track lines, the combined regional database, preliminary catalog, and hard copies of the original field datasheets if available, were sent to Cascadia for reference and archiving.

Data compilation and catalog development

Each regional database was imported into the combined seasonal SPLASH sighting database. SPLASH data were compiled using a Microsoft Access relational database which was designed specifically for the SPLASH project and which was comprised of six related tables for field data. The top tier table contained a record for each vessel survey day that included regional information, contributing organization, personnel, effort type (SPLASH dedicated or opportunistic) and total survey hours. Related to this table by research group (contributing organization), date, and vessel were tables with time and position data for effort-related events, such as changes in environmental conditions and whale sightings. The remaining two tables, which contained records for each individual identified and samples collected during a sighting, were linked to the sightings table by research group, date, vessel, and a unique sighting number for that vessel-day.

Upon receipt, the preliminary catalog was reviewed for metadata completeness and accuracy, and fields were added or reformatted as necessary for each best fluke image to include linked information for research group, photographer, sighting information, and the working ID number of the pictured whale. The preliminary catalog was converted to a digital print layout using the *Fotoslate* plug-in of the *ACDSee* photo management software package. The layout applied custom settings to each image that cropped and adjusted the exposure of each fluke and converted each image to grayscale prior to printing, without actually altering the original image file, thus avoiding any loss in quality that may have resulted from sequentially resaving JPG files.

Layouts were printed with associated metadata on a label. The layouts were cut into individual fluke photos, which were then sorted into the numeric color categories (1-5, from lightest to darkest) that were used to organize all SPLASH collections at the central matching level. Some color categories were further separated into subcategories by proportion of pigmentation and, in the case of all black flukes, by a hierarchical classification of mark types. During this step, incidental matches within a collection were sometimes encountered. If a significant number of internal matches were found, a complete re-reconciliation of the preliminary catalog was undertaken. As part of the initial color-sorting procedure, very poor quality flukes were rejected from the collection.

After sorting and reconciliation, remaining best fluke photos from each region were coded on a scale of 1-5 for five quality features (proportion visible, vertical angle, lateral angle, focus/sharpness, and exposure) and three characteristic features (distinctiveness of the trailing edge, degree of scarring, and presence of killer whale rake marks) using the same process developed previously (Calambokidis *et al.* 1997, 2000). As part of the coding process, each printed fluke photo was verified in the database, and other resightings of the pictured whale were reviewed for accuracy and to verify that the fluke photograph in the preliminary catalog was the best of season. After an entire collection was coded, flukes that received a score of 4 or higher in any quality category or scored 3 in more than three quality categories were rejected (however there were some exceptions to this in the case of small collections, where marginal flukes were included in the match to augment sample size). In most cases, if a rejected fluke was at or near the level of acceptability, all images of the whale from its initial sighting were reviewed to ensure that the best fluke of the sighting had accurately been selected per SPLASH criteria, and in some cases the original best fluke was replaced. All best flukes and rejected flukes were quality coded in the database; known resighting flukes were not coded.

After coding, the best fluke photos were placed in archival-quality clear plastic sleeves within their assigned color category. Any flukes that were known to be from calves (and thus subject to significant change in appearance over time) or which had ambiguous coloration (and were thus difficult to confidently assign to a color category) were flagged in the database and a copy of the photo was placed in a supplemental category at the end of each catalog for special attention during matching. Once assembled, the final catalog was reviewed for consistency in color-coding and quality screening, and each individual was assigned a six-digit SPLASH ID number, which reflected the season and region in which it was assigned to the whale. SPLASH IDs were then updated into all identifications of the whale in the identifications table, and each fluke identification record was classified as either the catalog best for the pictured whale, a regional resighting for that season, or a reject in the coding table.

Photographic matching

All SPLASH comparisons were conducted manually by a team of six experienced humpback whale fluke matchers. Because it was not feasible to manually compare every whale in every catalog against every other whale as the SPLASH collection became progressively larger, a series of protocols were developed to expedite the matching process. These protocols increased the likelihood of finding matches quickly, removed whales from the match process once found in an earlier catalog, and systematically limited the number of flukes against which each whale was ultimately compared; subsequently they introduced a certain potential to miss matches by exclusion. Extensive collateral data were collected with each comparison to record which photos were actually compared, as well as to later assess any biases in match rate associated with factors such as matcher, fluke color, or collection.

A series of regional catalogs were organized for each season, although the Summer 2004 SWFSC cruise and NMML cruises were combined in a single catalog. Regional catalogs from a

current season were sequentially compared against all earlier catalogs¹, prioritizing comparison against the regions of highest known match rates (either from other studies or previous seasonal comparisons). In this way, each regional catalog was always compared first to the same region in a previous year (*e.g.*, Hawaii 2005 against Hawaii 2004), and then to the feeding or wintering area catalogs known to have the highest interchange rates (*e.g.*, Hawaii 2005 against SEAK 2004). Once a whale was found in an earlier catalog, the previously assigned SPLASH ID was recorded in the match log, along with the relative quality of the new photo to the previous photograph, and any changes in the fluke itself. If the newer photograph of the whale was of comparable or lesser quality to the older photo and there were no significant changes in the fluke, the newer was covered in its catalog and excluded from further comparison against other collections. If the new fluke photograph was substantially better quality and/or different than the earlier photo, it remained uncovered and continued through the comparison to all other earlier collections.

The fluke photograph being matched was compared to all flukes in the corresponding color category of the earlier catalog, as well as all flukes in the color categories preceding and following, and all flukes in the calf/ambiguous section. Because it was sometimes more difficult to accurately assign color to the darkest flukes, all whales in the 4C section were compared against the 4B section and all the 5 sub-categories, and all whales in the 5 sub-categories were also compared against the 4C section. At matcher discretion, a given fluke was compared to additional color categories beyond those dictated by protocol if either its coloration or quality warranted a broader search.

When comparisons to all previous season catalogs were completed for a current season, the remaining uncovered photos in each regional catalog were compared against each other to identify any same-season movements among whales new to the SPLASH collection. All match logs for the season were compiled in preparation for collapsing catalogs and updating the SPLASH IDs assigned that season to the lower number for all whales that had been identified previously. As a final step to completing a seasonal match, any better/changed photos of whales seen previously were moved into the earliest catalog in which the match was found, photographs of whales seen previously that were not better were moved out of the active matching section of the newer catalog, and whales new to SPLASH were condensed in their current catalog in preparation for the next seasonal comparison. Consequently, there was only ever one active photo of each whale, its best to date, which was being compared against in subsequent seasons.

Upon completion of the final season of matching (Winter 2006), catalogs were not collapsed as in previous seasons. Instead, SPLASH IDs were updated on the photos and in the database, and original catalogs were reconstructed so that selective matches between collections could be conducted using the same photos initially compared. In this way, a whale seen in multiple seasons and regions would be represented by its best of collection photo in every catalog it had been in, but with its unified, lowest SPLASH ID number. Additionally a single

¹ The one exception to this pattern was Winter 2004, which was compared against Summer 2004, rather than vice versa, to facilitate simultaneous comparisons by multiple matchers working within two collections that were still relatively small.

digital catalog was compiled containing the single best photograph of each individual identified in SPLASH.

Evaluation of rates of missed matches and double checks

A complete double match of the entire SPLASH collection was not possible due to the number of photographs, so several experiments were conducted to quantify the error rates associated with the initial match. Nine comparisons among collections were completely rechecked: five rechecks were conducted throughout the matching process to verify and refine matching protocols, and four were conducted after the final match was completed.

A more systematic assessment of error rate was included in the final Winter 2006 comparison by “seeding” the Winter 2006 catalogs with 266 known matches to earlier catalogs. The Winter 2006 catalog included roughly 10% “seeds” (relative to its initial size), which were evenly distributed across color categories, quality scores, matching regions, and seasons. The seeded matches were assigned false SPLASH IDs and printed with mock labels so that the matchers were blind to which photographs were seeds. In this way, a rate at which known matches were missed was calculated for each matcher and factors which contributed to a higher than usual miss rate could be identified as Winter 2006 was compared to the 21 previous SPLASH catalogs.

In total 246 of 266 seeded matches were found (92%), consistent with expectations and similar to the match success rate found in a previous study using similar quality scoring criteria as employed here (Calambokidis *et al.* 1997). The average score of the five quality criteria used to rate photographs was only slightly higher (poorer quality) for those that were missed ($n = 20$, mean = 2.16, SD = 0.28) than those that were found ($n = 246$, mean = 2.03, SD = 0.33). Although this difference was not significant, the results of specific quality characteristics proved relevant. We looked at the success rate in finding those seeded matches where the photograph was at our cut-off for acceptance; match success rate ranged from high of 96% for photographs with no quality scores at three (the poorest acceptable rating) to a low of 82% when three of the criteria were rated as a three (no more than three scores of three were allowed or the photograph to be included in the SPLASH comparison). The clarity of the photograph (*i.e.*, quality score for focus) appeared to be the most important factor in whether a match was missed, with the largest difference in this category between the photographs where matches were found and those that were missed (mean score of 2.1 vs. 2.6, a highly significant difference, ANOVA, $p = 0.005$). Similarly, when the quality scores of both the seeded match and the whale it matched to were pooled, both the quality score for exposure and focus were significantly better for whales where the match was found versus those that were missed (ANOVA, $p = 0.04$ and $p = 0.001$ for exposure and focus, respectively).

Selective rechecking also indicated an overall match success rate of close to 90% for both initial and secondary comparisons. A total of 148 out of 165 matches were found (90%) in either an initial matching or in a recheck. Two matches were missed due to protocol (color category not checked) and discovered incidentally. The influence of photograph quality was assessed for the remaining 31 missed matches. Photo quality did not appear to be a factor in 10 of 31 missed matches (39%), where both the photo being matched and the catalog photo had better than

average quality scores (<1.9) and did not score a 3 in any category. For the remaining 21 cases, 11 misses were attributed to low quality in both photos, 5 involved a low quality photo in the catalog, and 5 were attributed to a low quality photo in hand. No specific quality fault appeared unusually high among missed match photos.

Based on the relatively high success rate in finding matches (at least 90%) no additional exclusions from the analysis were made based on quality. Additional matches that were found in the rechecks were included in the SPLASH sample because a correction factor for missed matches was not being applied.

ANALYTICAL METHODS

We examined abundances of humpback whales using several capture-recapture methods including simple Chapman/Petersen models and a more complex multi-strata model to estimate the abundance of humpback whales in feeding and wintering areas of the North Pacific in conjunction with calculated migration rates among these areas. We used a geographically stratified mark-recapture model (Hilborn 1990) similar to that used in the past analyses of the 1990-93 humpback photo-identification data (Calambokidis *et al.* 1997). In this approach, parameters for migration between feeding and wintering areas, survival, and capture probability were estimated in a likelihood setting. Abundance was estimated by dividing the number of sampled animals in an area by the estimated capture probability. Details of the methodology as applied to feeding areas in Southeast Alaska are given in Straley *et al.* (In press).

For mark-recapture abundance estimates, known calves were excluded from their first winter and feeding season because these animals can be harder to identify and accurately match. For the Hilborn models we also restricted the number of feeding and wintering areas to six each (with five seasons) as summarized in Table 3. Because the model did not allow for an animal to be seen in multiple areas in the same period, in the event an animal was seen in more than one area it was assigned to the area where it was seen closest to the middle of the season (1 March for wintering areas and 1 August for feeding areas). In the few cases where whales were identified on feeding areas in winter months, they were assigned to the previous or following feeding season using 1 March as a cut-off (even though they may have been submitted and compared as if they were part of a different season).

The Hilborn method estimates the size of a geographically stratified population that moves between areas and sampled during different time periods (Hilborn 1990; Quinn and Deriso 1999, Chapter 10; Calambokidis *et al.* 1997). For a given release group in a given year stratified by area, a model was constructed to predict the matrix of recaptures by release area and recovery area for each time period. In this application, two sets of areas were defined: six wintering areas and six feeding areas, and given time-periods denoted [Winter 2004 (W04), Summer 2004 (S04), Winter 2005 (W05), Summer 2005 (S05), and Winter 2006 (W06)]. It is assumed that survival is equal to 1, because of the short duration of the study (2 years) and thus estimates of abundance would not be appreciably biased. The key information derived from mark recaptures were estimates of capture probabilities and probabilities of movement between different areas. We denoted capture probability as p_{tj} for time period t and capture area j .

Two different assumptions about movement were evaluated. The first assumption was that movement followed a Markov process, in which the probability that a mark was in a given area depended on where the mark had been the previous season. This required having two movement matrices: Θ_{WS} for winter-to-summer movement and Θ_{SW} for summer-to-winter movement, each of size 6x6. The second assumption was that movement followed a non-Markov process, in which the probability that a mark was in a given area was not dependent on season but rather depended on where the mark had been the previous year. In other words, winter recaptures depended on where the whales had been the previous winter, and summer recaptures depended on where the whales had been the previous summer. This required having two additional movement matrices: Θ_{WW} for winter-to-winter movement and Θ_{SS} for summer-to-summer movement.

For this application, there were release groups in each of the first four time periods. The first release group in W04 had four sets of mark-recapture matrices in subsequent time periods, and each subsequent release group had one less mark-recapture matrix. A schematic showing which movement matrices were used in each period is shown in Table 4.

Table 4. Depiction of which movement matrices were applied to recapture periods for each release group, (a) Markovian movement, (b) Non-Markovian movement.

(b)	Recapture period				(a)	Recapture period			
Release group	S04	W05	S05	W06	Release group	S04	W05	S05	W06
W04	Θ_{WS}	Θ_{WW}	Θ_{SS}	Θ_{WW}	W04	Θ_{WS}	Θ_{SW}	Θ_{WS}	Θ_{SW}
S04		Θ_{SW}	Θ_{SS}	Θ_{WW}	S04		Θ_{SW}	Θ_{WS}	Θ_{SW}
W05			Θ_{WS}	Θ_{WW}	W05			Θ_{WS}	Θ_{SW}
S05				Θ_{SW}	S05				Θ_{SW}

A recursion process was used to follow the movement of the number of marked whales (release group) from a given time period. The release group k consisted of newly identified whales not previously seen. The recursion started with a diagonal matrix of releases in each area. Movement was assumed constant over time for parsimony.

For the Markov process, the formula for the predicted number of marked whales in area j that came from area i at the next season was found from the number of marked whales $M_{m \rightarrow i, t}$ in area i that came from all areas m in year t , given by:

$$M_{i \rightarrow j, t} = \sum_m M_{m \rightarrow i, t} \theta_{WS, i \rightarrow j} \text{ for winter-to-summer movement,}$$

$$\text{and } M_{i \rightarrow j, t+1} = \sum_m M_{m \rightarrow i, t} \theta_{SW, i \rightarrow j} \text{ for summer-to-winter movement.}$$

(The time subscript denotes year, so that summer follows winter in the same year t , and winter the next year ($t+1$) follows summer.) These equations can be compactly written in matrix notation as $\mathbf{M}_{S,t} = \mathbf{M}_{W,t} \Theta_{WS}$ and $\mathbf{M}_{W,t+1} = \mathbf{M}_{S,t} \Theta_{SW}$, respectively, in which the subscripts S and W denote summer and winter.

For the non-Markov process, the predicted number of marked whales in the season following release used these same equations. Thereafter, the predicted number of marked whales in winter depended on the previous winter, such that $\mathbf{M}_{W,t+1} = \mathbf{M}_{W,t} \Theta_{WW}$. Similarly, the predicted number of marked whales in summer was determined from $\mathbf{M}_{S,t+1} = \mathbf{M}_{S,t} \Theta_{SS}$.

The predicted number of recaptures was then found by multiplying the predicted number of whales by the capture probability in area j , or $m_{i \rightarrow j,t} = M_{i \rightarrow j,t} p_j$. This prediction was then compared to the observed number of marked recaptures from the release group in a likelihood setting to estimate parameters. Estimates of movement and capture probability parameters were obtained numerically by maximizing the likelihood, here assumed to be a product of Poisson distributions. All calculations were done in Excel using its Solver optimizer.

Once capture probabilities were estimated (\hat{p}_{ij}) for each area j , the general law of estimating abundance (Seber, 1982) was used to produce the estimates of abundance as $\hat{N}_{ij} = n_{ij} / \hat{p}_{ij}$, where n_{ij} is the number of animals examined for marks at time period t in area j .

Four different modeling scenarios made different assumptions about how probability of capture varied over time. The first scenario assumed that capture probabilities were constant over time [p(.)]. The second scenario assumed that capture probabilities varied over time [p(t)]. The third scenario assumed that capture probability was proportional to sample size each year [p(n)]. Effectively, model p(n) resulted in a single estimate of abundance for each area, since $N = n/p$. In the Markov case, the fourth scenario was a modification of p(n), in which the capture probability for SBC/NWA was constant over time, in an attempt to constrain it to a realistic value. In the non-Markov case, a fourth scenario was examined for analytical purposes, in which winter capture probabilities were proportional to sample size and summer capture probabilities were constant [p(S., Wn)]

Model selection followed the procedures outlined in Burnham & Anderson (2002), including the use of AICc (Akaike Information Criterion, corrected, p. 51) as a model comparison statistic. Central to AICc is the calculation of the difference Δ between a given model and the model with the lowest AICc value. Models with $\Delta < 4$ were considered to be very similar, while models with $\Delta > 10$ were considered unlikely to be correct.

RESULTS AND DISCUSSION

A total of 18,469 fluke identification photographs of acceptable quality were taken during over 27,000 humpback whale approaches. After reconciling all within and cross-regional matches (from both the primary match and rechecks), a total of 7,971 unique individuals were cataloged in SPLASH and 6,178 tissue samples collected (see Table 2). A total of 4,516 individuals were identified at wintering regions in at least one of the three seasons and 4,328 individuals were seen at least once at feeding areas in one of the two years.

Interchange within wintering areas

Interchange within some wintering regions, like Hawaii, was extensive (Table 5). Although most of the Hawaii sample came from the Maui sub-area, identifications from Big Island and Kauai at the eastern and western end of the region showed a high rate of interchange with Maui. Although there were some indications of subtle differences in migratory destinations among some sub-areas discussed later, for most of the analyses reported here, the Hawaii sub-areas are treated as a single unit.

Table 5. Rates of interchange among the Hawaii subareas for 2004-2006. Numbers along the diagonal show the total number of unique identifications within that subarea and numbers along upper right portion of the matrix show number of individuals seen in multiple sub-areas.

Area	Kauai	Oahu	PB	Molokai	Maui	Big Island
Kauai	203	1	0	4	29	2
Oahu		89	0	5	20	9
PB			34	3	4	3
Molokai				201	61	12
Maui					1526	99
Big Island						507

Interchange of whales among the three principal wintering regions sampled (Asia, Hawaii, and Mexico) was relatively low although there were a few animals seen in different major regions in different years (Table 6). Two whales were seen in both Asia (one each in Ogasawara and Philippines) and Hawaii. Similarly, 17 whales were seen in both Hawaii and one of the Mexican wintering sub-areas in different years (14 to the Revillagigedos, 2 to Baja, and 1 to mainland).

Interchange among the three sub-areas of Mexico and Central America was more complex. Of 562 whales from the Revillagigedos included in the SPLASH study, 112 (20%) were sighted at the Revillagigedos in more than one winter season and 48 (8.5%) were resighted at other wintering areas (Table 6). The highest number of wintering area matches was to Baja (22 whales), followed by equal numbers of matches to mainland Mexico and Hawaii (14 each; Table 6). Two whales were sighted at the Revillagigedos and both of the other Mexican wintering sub-areas during the study; no whales were observed at the Revillagigedos and Central America or Asia. The comparable match rates to mainland Mexico and Hawaii is interesting, given that the distance between the Revillagigedos and mainland Mexico (approximately 580km) is only

slightly greater than the distance between Isla Socorro and Isla Clarion, the two most separated islands in the Revillagigedos chain (approximately 400km), and the distance between the Revillagigedos and the easternmost of the Hawaiian Islands is 4,600km. No same-season transits were observed between the Revillagigedos and Hawaii, but 3 same-season transits were observed between the Revillagigedos and mainland Mexico. In comparison, there were 22 same-season transits between islands at the Revillagigedos, despite that the majority of the Revillagigedos effort was limited to Isla Socorro only.

A closer connection was found between the whales that winter off Central America and those that winter off Mexico. Central America is the southernmost humpback whale wintering area in the North Pacific, and SPLASH represents the first time a relatively large number of whales from Central America have been compared widely across the ocean basin and especially to a concurrent sample from Mexico. Nine whales from Central America were sighted in mainland Mexico and two whales from Central America were sighted in Baja during the course of the study (Table 6). Three same-season movements between Central America and mainland Mexico were documented. In all cases the whales were sighted first at mainland Mexico and then later in Central America (28, 37, 52 days transit). The average sighting dates of whales seen in both areas (in the same or different seasons) was significantly earlier off mainland Mexico (28 December) than off Central America (5 February) ($t = 3.97$, $P = 0.002$), however this may partly be the result of the timing of effort, which was limited in Central America. The two whales that were photographed in both Baja and Central America were sighted later in the season off mainland Mexico (on 26 and 30 March), suggesting that southbound whales en route to Central America may pass too early in the season to be captured in Baja, or do not spend a significant amount of time in the area. It is also possible that Central America whales vary their migratory route north and southbound.

Table 6. Interchange among wintering areas 2004-2006. Numbers along the diagonal show the total number of unique identifications within that area and numbers along upper right portion of the matrix show number of individuals seen in multiple areas.

Area	Asia-							
	PHI	Asia-OK	Asia-OG	HI	MX-REV	MX-Baja	MX-ML	Cent Am
Philippines	77	5	5	1	0	0	0	0
Okinawa		215	10	0	0	0	0	0
Ogasawara			294	1	0	0	0	0
Hawaii				2317	14	2	1	0
MX-REV					562	22	14	0
MX-Baja						406	66	2
MX-ML							690	9
Cent Am								105

Interchange and movements among and within feeding areas

Even though multiple sightings of the same individuals were made on the feeding areas, most of these showed only infrequent exchanges among areas, which supported an overall conclusion of a high degree of site fidelity to feeding areas. There were cases of this interchange both within and between 2004 and 2005 seasons (Figure 2).




Figure 2. Summary of resightings of humpback whales on feeding areas both within and between the 2004 and 2005 seasons.

Same-season movement among the feeding areas was best evaluated in 2004 when survey efforts were broadest due to the four-month SPLASH cruise conducted by SWFSC supplementing more localized effort in British Columbia, SE Alaska, Gulf of Alaska, Aleutians, and Bering Sea (Table 7). Whales that were seen on multiple days were generally resighted in the same area and in only 12 cases were resighting locations more than 400 nmi apart. Overall, most resightings of whales in the feeding areas were within the same area and in only 42 cases were whales seen in different regions. Out of 789 whales seen on more than one day, 42 (5%) were seen in more than one defined area. Even among these 42, all but three were resightings made in adjacent areas. Adjacent areas with relatively high rates of within-season interchange included SE Alaska and N British Columbia (19), SE Alaska and N Gulf of Alaska (10), North and West Gulf of Alaska (4), and Bering Sea and E Aleutians (5).

Interchange between feeding areas was relatively uncommon; most individuals that were sighted in Summer 2004 and 2005 were sighted in the same area in both years (Table 8). Most cross-regional resightings of whales again occurred between adjacent areas. The highest rates of adjacent-area resightings occurred between SE Alaska and N British Columbia, however even here, cross-regional resightings were much less frequent than resightings within each area. Other adjacent areas with intermediate rates of interchange included the N and W Gulf of Alaska and the N British Columbia and S British Columbia/Washington. There were only three examples of shifts in location between years that were farther than the nearest adjacent area: 1) one whale seen in N British Columbia in 2004 and resighted in N Gulf of Alaska in 2005, 2) one whale from SE Alaska in 2004 resighted in W Gulf of Alaska in 2005 and 3) one whale seen of W Gulf

of Alaska in 2004 and resighted off California on 18 November 2005, possibly on its southern migration.

Table 7. Summary of interchange of humpback whales across feeding areas during SPLASH effort in the 2004 feeding season. Numbers along the diagonal indicate photo-IDs in the area.

	W		E		SBC-					
	Russia	Aleut	Bering	Aleut	WGOA	NGOA	SEAK	NBC	NWA	CA-OR
Russia	40									
W Aleut	0	12								
Bering	0	0	228							
E Aleut	0	0	5	51						
WGOA	0	0	0	0	223					
NGOA	0	0	0	1	4	730				
SEAK	0	0	1	0	0	10	808			
NBC	0	0	0	0	0	1	19	421		
SBC-										
NWA	0	0	0	0	0	0	0	1	76	
CA-OR	0	0	0	0	0	0	0	0	0	253

Table 8. Interchange among feeding areas based on sightings in 2004 (rows) and 2005 (column headings).

		2005							
		Russia	Bering	WGOA	NGOA	SEAK	NBC	SBC-NWA	CA-OR
2004	IDs	72	301	111	427	482	236	152	319
Russia	40	10	0	0	0	0	0	0	0
Aleut-Bering	291	0	41	0	0	0	0	0	0
WGOA	223	0	0	33	1	0	0	0	1
NGOA	730	0	0	6	119	1	0	0	0
SEAK	808	0	0	1	4	175	16	0	0
NBC	421	0	0	0	1	13	74	4	0
SBC-NWA	76	0	0	0	0	0	1	21	1
CA-OR	253	0	0	0	0	0	0	1	47

Identifications from Russian waters came from three distinct areas, the Commander Islands at the western end of the Aleutians, an area off the east side of Kamchatka, and in the Gulf Anadyr at the north end of the Bering Sea. These areas were sampled in both 2004 and 2005 with the exception of the Gulf Anadyr that was sampled only in 2005. All 10 of the resightings of humpback whales in Russian waters between 2004 and 2005 were made within these sub-areas. Nine whales were resighted off Kamchatka between years, and the other was resighted at the Commander Islands (Table 9). While the small sample size would limit the expected number of resightings, the finding that none of the 10 resightings involved interchange among these areas indicated site fidelity within these areas and some degree of separation between them.

Table 9. Summary of matches made between the three subareas in Russian waters in 2004 and 2005 showing all resightings between years were within the same subarea

Area 2005	IDs	2004	
		Commanders	Kamchatka
Commander Is	7	1	0
Gulf of Anadyr	27	0	0
Kamchatka	38	0	9

Migrations between wintering and feeding areas

Migrations between feeding and wintering areas were documented based on 873 whales that were seen at both wintering and feeding regions (Table 10, Figure 3). Movement patterns were complex but indicated a high degree of population structure. The overall pattern showed that wintering areas on both sides of the Pacific (Asia in the west and mainland Mexico and Central America in the east) are the primary wintering areas for the lower latitude coastal feeding areas on the same side of the Pacific. The regions off Hawaii and the Revillagigedo Archipelago were the primary wintering areas for the more central- and northern-latitude feeding areas. Within this broad overall pattern, however, were more complex and sometimes surprising movements and structure that are discussed in more detail below.




Figure 3. Locations of SPLASH identifications. Lines connect sequential resightings of the same individual.

Table 10. Identifications and connections made between wintering areas (columns) and feeding areas (rows). Seasons are pooled for both wintering and feeding areas. Sum reflects the total of the whales matching between areas and Overall reflects the number of individuals matching to any area (in a few cases the same whale matched to multiple areas).

Region	Daily IDs		Asia-PHI	Asia-OK	Asia-OG	Hawaii	MX-REV	MX-Baja	MX-ML	Cent Am	Sum	Overall
	Unique		151	448	602	3205	2009	465	1222	140		
Russia	128	102	6	14	5	4	1	0	0	0	30	29
Aleutians	64	63	0	1	0	4	0	2	0	0	7	7
Bering	728	491	0	1	5	44	11	8	11	0	80	77
WGOA	516	301	0	0	2	26	13	7	4	0	52	51
NGOA	1792	1038	0	0	1	124	44	20	21	0	210	200
SEAK	2382	1115	0	0	0	215	9	3	8	0	235	235
NBC	1183	583	0	0	0	99	8	5	4	0	116	114
NWA-SBC	380	207	0	0	0	20	2	8	22	3	55	53
CA-OR	881	525	0	0	0	0	0	20	97	26	143	133
Sum			6	16	13	536	88	73	167	29	928	
Overall matches			6	16	13	516	87	70	164	29		873

Some feeding areas in the central North Pacific had very low match rates to any wintering ground (Figure 4). The lowest proportion of animals going to any sampled wintering area occurred in the Aleutian region where only about 11% of the whales identified matched to any wintering area. Humpback whales identified in neighboring areas including the Commanders, Bering Sea, and Western Gulf of Alaska had just slightly higher proportions with 15-17% matching to wintering areas. This is in contrast with areas like SE Alaska where almost all animals go to Hawaii and over 21% match a wintering area, California-Oregon where most animals migrate from mainland Mexico and Central America and over 25% match a wintering area, and Russia where close to 40% match Asian wintering areas. One likely explanation for this pattern is the existence of an unknown wintering ground not sampled in SPLASH that serves as a destination for these animals. We also cannot rule out that this pattern could also be created by a combination of small sample sizes for the Aleutians and Russia feeding areas, under-sampled wintering areas or use of a known wintering ground outside the sampled period, although this seems unlikely given the sampling strategy and the high proportion (>20%) of animals from most wintering areas present on some feeding areas (Russia for the Asia wintering areas, SE Alaska for Hawaii, and California-Oregon for mainland Mexico and Central America).

With the exception of Asian wintering areas, over 15% of animals at wintering areas were identified on feeding areas (Figure 5). At all three Asian wintering areas less than 10% of the identified whales had also been seen on a feeding ground and this was less than 5% for Ogasawara. This supports the idea that the SPLASH sample under-represents the feeding areas that are the main destinations for animals wintering in Asia. This is not too surprising given the limited effort in the difficult to study areas in the western Aleutians, Bering Sea, and waters off Russia.




Figure 4. The proportion of whales on feeding areas that were seen on different wintering areas.




Figure 5. The proportion of whales on wintering areas that were seen on different feeding areas.

Humpback whales that were identified on feeding areas in Russian waters migrated primarily to Asian wintering areas, but strong differences in winter destinations existed among whales at the three Russian sub-areas (Table 11). Humpback whales identified along the east side of Kamchatka had high match rates to the three Asian wintering areas (especially Philippines and Okinawa) with 38% of these identified whales having been seen in at least one of these three wintering areas and no matches to Hawaii (Table 11). In sharp contrast, humpback whales identified off the Commander Islands and farther north in the Bering Sea in the Gulf Anadyr, had very low overall incidence of matches to any wintering area, and those that were found included more matches to Hawaii and Mexico than to Asian wintering areas. The match rates and destinations for the animals from the Commanders and northern Russia were much more similar to those for the Aleutians and Bering Sea (US side) than they were for the area off Kamchatka (Table 11). For this reason we used an alternate regional assignment for some of the mark-recapture estimates that reclassified identifications from the Commander Is and the Gulf of Anadyr to the Aleutians and Bering Sea (where they would reasonably belong though across an international border) and categorized the Kamchatka identifications as a separate region.

Table 11. Summary of migratory destinations of humpback whales from Russian waters showing differences by sub-area and similarities to the Aleutian and Bering areas in US waters.

	IDs	Asia- PHI 77	Asia- OK 215	Asia- OG 294	Hawaii 2317	MX 1558	Any area	%
Gulf of Anadyr	27	0	1	0	3	0	4	15%
Kamchatka	58	6	13	4	0	0	22	38%
Commanders	17	0	0	1	1	1	3	18%
All Russia	102	6	14	5	4	1	29	28%
Aleutians	63	0	1	0	4	0	7	11%
Bering	491	0	1	5	44	27	77	16%

This study also provided the first clear insight into the migratory destinations of whales that winter in the Revillagigedos. Eighty-seven of 562 whales from the Revillagigedos (15.4%) were sighted at a feeding area, a substantial increase in the feeding area matching rates reported from all previous studies of 3% or less. The whales from the Revillagigedos were seen in all sampled feeding areas except California-Oregon and the south side of the Aleutians. The majority of movements observed were to N Gulf of Alaska (n=44) and W Gulf of Alaska (n=13), with match rates declining to the west and east to a low of one match to Russia and two matches to Washington-S British Columbia. The movement observed between the Revillagigedos in March 2005 and the Commander Islands in the Russian Bering Sea in August that same year is the first documented, and represents a minimum migratory distance of 7,925 km.

Current Abundance

Chapman/Petersen mark-recapture estimates using all pooled regions for each of the five SPLASH seasons yielded consistent abundances increasing from 18,347 to 21,452 (Table 12). While one reason for these increased estimates could be a reflection of an increasing population, the rate of increase is higher than would be expected (see later sections) and such fluctuations over a relatively short time scale may be the result of other sampling factors in conjunction with an increasing population. Because calves (where identified) were excluded in their initial year, these estimates are of the non-calf portion of the population. While these are closed population models and there is some violation to closure from natality and mortality, the time intervals are less than a year so these violations should result in only a small upward bias in the estimates. This would also be offset by the small downward bias introduced by excluding calves.

Table 12. Chapman/Petersen mark-recapture estimates using pooled regions for each of the five SPLASH seasons. Estimates are based on four successive paired periods alternating between winter and feeding seasons.

Sample periods	Unique Ids			Estimate	CV	CV-Jk*
	n1	n2	Recapt.			
Winter 2004 - Summer 2004	1,588	2,724	235	18,347	0.06	0.13
Summer 2004 - Winter 2005	2,724	1,685	247	18,525	0.06	0.53
Winter 2005 - Summer 2005	1,685	2,021	169	20,052	0.07	0.20
Summer 2005 - Winter 2006	2,021	1,930	181	21,452	0.07	0.17

*Jackknife estimate of variance treating entire regions as samples, sometimes high due to the large influence of eliminating some larger samples like Hawaii

As expected, mark-recapture estimates based on paired years exclusively from two of the three winter seasons or exclusively from the two summer feeding seasons yielded unrealistically low estimates of abundance (Table 13). Estimates from only winter areas tend to be biased because males are more likely to return and are identified in greater numbers than females (Brown *et al.* 1995, Craig and Herman 1997, 2000, Craig *et al.* 2003) and juveniles are probably underrepresented (Robbins 2007). These biases have been shown to result in dramatic underestimates of abundance (Calambokidis *et al.* 1997, Smith *et al.* 1999). While the sex ratio on the feeding grounds are more representative of the population overall (Clapham *et al.* 1995), the estimate based only feeding areas samples would be biased downward by heterogeneity in capture probability created by the tendency to capture the same animals in the same areas in similar effort in the two years. The knowledge of these biases was the reason the focus for accurate abundance was on estimates using the combination of wintering and feeding areas.

Mark-recapture estimates of abundance and capture probabilities based on the Hilborn model with Markovian movement are compared in Table 14 and key parameters summarized in Tables 15 and 16. These indicated the p(t) model, in which capture probabilities varied over time, had the lowest AIC and AICc scores. However, all four models had AICc scores within 7 of the p(t) model, indicating that none of these models could be ruled out.

Table 13. Petersen mark-recapture estimates using pooled regions for each of the five SPLASH seasons comparing only Winter to Winter and feeding to feeding for pairs of years.

Sample periods	Unique Ids			Estimate	CV	CV-Jk*
	n1	n2	Recapt.			
Winter 2004 - Winter 2005	1,588	1,685	298	8,959	0.05	0.28
Winter 2005 - Winter 2006	1,685	1,930	278	11,668	0.05	0.31
Summer 2004 - Summer 2005	2724	2021	544	10,109	0.03	0.40

*Jackknife estimate of variance treating entire regions as samples

Table 14. Comparison of key parameters for four Hilborn models with Markovian movement. Values in bold reflect lowest model score.

Parameter	p(.)	p(t)	p(n)	p(n)(2)
log likelihood	-974	-953	-976	-976
Observations	360	360	360	360
Parameters	84	96	84	83
Constraints	12	12	12	12
Unconstrained parameters	72	84	72	71
AIC	2,092	2,074	2,097	2,094
BIC	2,372	2,400	2,377	2,370
AICc	2,129	2,126	2,133	2,130

All of the Hilborn models gave fairly similar estimates of abundance and movement rates for most areas, although there were difficulties with some of the smaller areas, especially the Washington/S British Columbia feeding area. Overall, average estimates for either all feeding or all wintering areas in the four models ranged from about 16,000 to 17,000 (Table 15), with the p(t) model giving estimates of 16,184 for the combined wintering areas and 16,744 for the combined feeding areas. Among wintering areas, Hawaii was estimated at close to 8,000 (approximately half the population), the three Mexican areas totaled about 6,600 (with Revillagigedos and Baja the largest at about 2,600 and Mainland Mexico at about 1,400), Asia was estimated at about 1,000 and Central America at about 500. Among feeding areas, the combined SE Alaska and N British Columbia area had the highest abundance at about 6,000 followed by Gulf of Alaska (about 4,000), Aleutians-Bering Sea (about 3,000), California-Oregon (about 2,000), Russia (encompassing the other feeding areas for the Asia wintering area but not the Aleutians) (~1,200) and Washington/S British Columbia (~200).

A key element of the Hilborn model is the movement rates from wintering to feeding areas and return (Table 16). While many of these movement rates appear quite reasonable and provide the best integrated measure of whale migration between these areas, a few were surprising. In particular, all the models estimated high movement rates between Baja and the Aleutians/Bering Sea areas despite a relatively low number of matches between these areas. This may be a result of the low proportion of animals in the Aleutians/Bering Sea for which a wintering destination was found, forcing the model to adjust for this. One other surprising high movement rate was for Washington/S BC to Central America, which may be an artifact of the

Table 15. Summary of results of capture probabilities and abundance estimates for four models: 1) $p(\cdot)$ where capture probabilities are constant over time for each region, 2) $p(t)$ where capture probabilities vary over time, 3) $p(n)$ which assumes that capture probability is proportional to sample size each year referenced to Summer 2004 and Winter 2005 (effectively results in a single estimate of abundance), and 4) $p(n)(2)$ same as $p(n)$ except capture probability for SBC/NWA kept constant for Summer 2004 and 2005.

Capture probabilities and abundance																
Model $p(\cdot)$																
Capture probs p									Capture probs p							
Winter	Asia	HI	Mx-Rev	Baja	MX-Main	CentAm	Sum	Summer	Russia-Kam	Al-Ber	GOA	SEAK-NBC	SBC-NWA	CA-OR	Sum	
All	0.241	0.114	0.073	0.044	0.202	0.088		All	0.034	0.106	0.158	0.145	0.497	0.132		
Estimated abundance									Estimated abundance							
2004	759	6103	4211	4016	1038	205	16331	2004	738	2715	5825	8061	145	1879	19363	
2005	866	7370	2625	3419	1246	512	16037	<u>2005</u>	<u>1122</u>	<u>3063</u>	<u>3332</u>	<u>4766</u>	<u>274</u>	<u>2295</u>	<u>14852</u>	
<u>2006</u>	<u>1190</u>	<u>8935</u>	<u>2502</u>	<u>1882</u>	<u>1567</u>	<u>512</u>	<u>16587</u>	Average	930	2889	4578	6414	209	2087	17108	
Average	938	7469	3112	3105	1284	409	16319	Average	930	2889	4578	6414	209	2087	17108	
Model $p(t)$																
Capture probs p									Capture probs p							
Winter	Asia	HI	Mx-Rev	Baja	MX-Main	CentAm	Sum	Summer	Russia-Kam	Al-Ber	GOA	SEAK-NBC	SBC-NWA	CA-OR	Sum	
2004	Capture prob's in W04 cannot be estimated							2004	0.017	0.091	0.216	0.153	0.434	0.147		
2005	0.217	0.122	0.080	0.079	0.221	0.128	2005	0.042	0.109	0.139	0.145	0.497	0.128			
2006	0.256	0.110	0.065	0.025	0.191	0.071										
Estimated abundance									Estimated abundance							
2004	Abundance in W04 cannot be estimated							2004	1464	3180	4280	7622	166	1690	18402	
2005	965	6872	2408	1876	1140	351	13612	<u>2005</u>	<u>895</u>	<u>2978</u>	<u>3795</u>	<u>4772</u>	<u>274</u>	<u>2372</u>	<u>15087</u>	
<u>2006</u>	<u>1119</u>	<u>9196</u>	<u>2832</u>	<u>3316</u>	<u>1658</u>	<u>636</u>	<u>18757</u>	Average	1179	3079	4038	6197	220	2031	16744	
Average	1042	8034	2620	2596	1399	493	16184	Average	1179	3079	4038	6197	220	2031	16744	
Model $p(n)$																
Capture probs p									Capture probs p							
Winter	Asia	HI	Mx-Rev	Baja	MX-Main	CentAm	Sum	Summer	Russia-Kam	Al-Ber	GOA	SEAK-NBC	SBC-NWA	CA-OR	Sum	
2004	0.171	0.084	0.117	0.072	0.149	0.037		2004	0.025	0.094	0.233	0.215	0.497	0.112		
2005	0.195	0.101	0.073	0.061	0.179	0.092		2005	0.038	0.106	0.133	0.127	0.938	0.137		
2006	0.268	0.123	0.069	0.034	0.226	0.092										
Estimated abundance									Estimated abundance							
2004								2004	989	3074	3962	5427	145	2206	15803	
2005	1071	8261	2637	2447	1406	489	16312	<u>2005</u>	<u>989</u>	<u>3074</u>	<u>3962</u>	<u>5427</u>	<u>145</u>	<u>2206</u>	<u>15803</u>	
<u>2006</u>	<u>1071</u>	<u>8261</u>	<u>2637</u>	<u>2447</u>	<u>1406</u>	<u>489</u>	<u>16312</u>	Average	989	3074	3962	5427	145	2206	15803	
Average	1071	8261	2637	2447	1406	489	16312	Average	989	3074	3962	5427	145	2206	15803	
Model $p(n)(2)$																
Capture probs p									Capture probs p							
Winter	Asia	HI	Mx-Rev	Baja	MX-Main	CentAm	Sum	Summer	Russia-Kam	Al-Ber	GOA	SEAK-NBC	SBC-NWA	CA-OR	Sum	
2004	0.171	0.084	0.117	0.071	0.155	0.033		2004	0.025	0.094	0.233	0.215	0.905	0.112		
2005	0.195	0.101	0.073	0.061	0.186	0.083		2005	0.038	0.106	0.133	0.127	0.905	0.137		
2006	0.268	0.123	0.069	0.033	0.234	0.083										
Estimated abundance									Estimated abundance							
2004								2004	989	3080	3963	5427	80	2208	15747	
2005	1071	8258	2638	2452	1356	542	16318	<u>2005</u>	<u>989</u>	<u>3080</u>	<u>3963</u>	<u>5427</u>	<u>150</u>	<u>2208</u>	<u>15817</u>	
<u>2006</u>	<u>1071</u>	<u>8258</u>	<u>2638</u>	<u>2452</u>	<u>1356</u>	<u>542</u>	<u>16318</u>	Average	989	3080	3963	5427	115	2208	15782	
Average	1071	8258	2638	2452	1356	542	16318	Average	989	3080	3963	5427	115	2208	15782	

Table 16. Summary of results of migration rates for four models listed in Table 15. Migration rates total 1 across rows for movement from either summer feeding areas to wintering areas or the return.

Model p(.)	⊕	S->W						Ψ	W->S					
		Asia	HI	Mx-Rev	Baja	MX-Main	CentAm		Russia-Kam	Al-Ber	GOA	SEAK-NBC	SBC-NWA	CA-OR
	Russia-Kam	0.990	0.010	0.000	0.000	0.000	0.000	Asia	0.963	0.037	0.000	0.000	0.000	0.000
	Al-Ber	0.013	0.169	0.032	0.765	0.020	0.000	HI	0.000	0.043	0.108	0.844	0.005	0.000
	GOA	0.002	0.169	0.734	0.070	0.026	0.000	Mx-Rev	0.000	0.026	0.950	0.022	0.002	0.000
	SEAK-NBC	0.000	0.973	0.015	0.007	0.005	0.000	Baja	0.000	0.695	0.108	0.027	0.013	0.157
	SBC-NWA	0.000	0.252	0.000	0.050	0.112	0.585	MX-Main	0.000	0.033	0.029	0.006	0.016	0.916
	CA-OR	0.000	0.000	0.000	0.149	0.708	0.143	CentAm	0.000	0.000	0.000	0.000	0.311	0.689

Model p(t)	⊕	S->W						Ψ	W->S					
		Asia	HI	Mx-Rev	Baja	MX-Main	CentAm		Russia-Kam	Al-Ber	GOA	SEAK-NBC	SBC-NWA	CA-OR
	Russia	0.989	0.011	0.000	0.000	0.000	0.000	Asia	0.960	0.040	0.000	0.000	0.000	0.000
	Al-Ber	0.013	0.165	0.035	0.766	0.021	0.000	HI	0.000	0.044	0.111	0.838	0.006	0.000
	GOA	0.002	0.162	0.748	0.063	0.025	0.000	Mx-Rev	0.000	0.027	0.948	0.022	0.003	0.000
	SEAK-NBC	0.000	0.972	0.017	0.007	0.005	0.000	Baja	0.000	0.698	0.107	0.026	0.013	0.156
	SBC-NWA	0.000	0.249	0.000	0.052	0.114	0.584	MX-Main	0.000	0.034	0.028	0.006	0.016	0.915
	CA-OR	0.000	0.000	0.000	0.148	0.713	0.138	CentAm	0.000	0.000	0.000	0.000	0.319	0.681

Model p(n)	⊕	S->W						Ψ	W->S					
		Asia	HI	Mx-Rev	Baja	MX-Main	CentAm		Russia-Kam	Al-Ber	GOA	SEAK-NBC	SBC-NWA	CA-OR
	Russia	0.989	0.011	0.000	0.000	0.000	0.000	Asia	0.959	0.041	0.000	0.000	0.000	0.000
	Al-Ber	0.013	0.168	0.036	0.764	0.019	0.000	HI	0.000	0.045	0.113	0.838	0.004	0.000
	GOA	0.002	0.165	0.747	0.062	0.024	0.000	Mx-Rev	0.000	0.027	0.950	0.021	0.002	0.000
	SEAK-NBC	0.000	0.973	0.016	0.007	0.004	0.000	Baja	0.000	0.706	0.108	0.026	0.007	0.153
	SBC-NWA	0.000	0.282	0.000	0.054	0.110	0.553	MX-Main	0.000	0.035	0.030	0.006	0.011	0.918
	CA-OR	0.000	0.000	0.000	0.165	0.682	0.153	CentAm	0.000	0.000	0.000	0.000	0.178	0.822

Model p(n) (2)	⊕	S->W						Ψ	W->S					
		Asia	HI	Mx-Rev	Baja	MX-Main	CentAm		Russia-Kam	Al-Ber	GOA	SEAK-NBC	SBC-NWA	CA-OR
	Russia	0.989	0.011	0.000	0.000	0.000	0.000	Asia	0.959	0.041	0.000	0.000	0.000	0.000
	Al-Ber	0.013	0.167	0.036	0.765	0.019	0.000	HI	0.000	0.045	0.113	0.839	0.003	0.000
	GOA	0.002	0.165	0.748	0.063	0.024	0.000	Mx-Rev	0.000	0.027	0.950	0.021	0.001	0.000
	SEAK-NBC	0.000	0.973	0.016	0.007	0.004	0.000	Baja	0.000	0.707	0.108	0.026	0.007	0.153
	SBC-NWA	0.000	0.276	0.000	0.051	0.104	0.569	MX-Main	0.000	0.036	0.030	0.006	0.010	0.918
	CA-OR	0.000	0.000	0.000	0.166	0.657	0.177	CentAm	0.000	0.000	0.000	0.000	0.162	0.838

small samples and abundance estimates for these two areas, because previous studies have shown a low interchange between these areas (Calambokidis *et al.* 2000, 2001).

Mark-recapture models based on the Hilborn model with non-Markovian movement are compared in Table 17 and key parameters summarized in Tables 18 and 19. These non-Markovian models fitted the data substantially better than the Markovian models, as indicated by the much lower AICc scores. The p(t) and p(n) models fitted the data better than the p(.) and p(S.,Wn) models. The p(n) model had the lowest AICc score, suggesting that capture probabilities were proportional to sample size in all areas.

Table 17. Comparison of key parameters for four Hilborn models with non-Markovian movement. Values in bold reflect lowest model score.

Parameter	p(.)	p(t)	p(n)	p(S.,Wn)
log likelihood	-489	-445	-465	-491
Observations	360	360	360	360
Parameters	156	168	156	156
Constraints	24	24	24	24
Unconstrained parameters	132	144	132	132
AIC	1,243	1,177	1,193	1,245
BIC	1,756	1,737	1,706	1,758
AICc	1,397	1,371	1,348	1,400

The non-Markovian models gave remarkably similar estimates of abundance in winter both by area and overall, but not in summer. Estimated total abundance ranged from 17,482 to 17,558 for the combined winter areas and from 18,568 to 21,225 for the combined summer feeding areas (Table 18). These abundance estimates are generally higher than for the Markovian models and closer to the pooled-area Petersen estimates. The p(n) model which provided the best overall fit to the data indicated an abundance of 17,558 for wintering areas and 19,056 for the feeding areas. The average of these two estimates (18,302) represents the best estimate of overall abundance of humpback whales in the North Pacific.

Among wintering areas, Hawaii was estimated at near 10,000 or about 57% of the population, the three Mexican areas totaled 6,000-7,000 (with Baja the largest at about 5,000 and Revillagigedo and Mainland Mexico at about 750), Asia was estimated at about 1,000 and Central America at about 500 whales. Among feeding areas, regional estimates differed greatly among models. Average estimates of abundance ranged from about 100-700 for Russia, 6,000-14,000 for the Bering Sea and Aleutians, 3,000-5,000 each for the Gulf of Alaska (W and N) and the combined SE Alaska and N British Columbia area, 200-400 for Washington/S British Columbia, and 1,400-1,700 for California-Oregon. Movement rates from summer to winter areas were similar among models (Table 19). Clear associations occurred between certain pairs of feeding and wintering areas (Russia/Asia, SE AK/Hawaii, CA-OR/Baja), but the Aleutians/Bering Sea, Gulf of Alaska, and S British Columbia/N Washington whales had more varied wintering destinations (Table 19).

Table 18. Modeling results based on non-Markov movement. The first scenario assumed that capture probabilities were constant [p(.)]. The second scenario assumed that capture probabilities varied over time [p(t)]. The third scenario assumed that capture probability was proportional to sample size each year [p(n)]. Effectively, model p(n) resulted in a single estimate of abundance for each area, since $N = n/p$ and the fourth scenario was examined for analytical purposes, in which winter capture probabilities were proportional to sample size and summer capture probabilities were constant [p(S., Wn)]

Capture probabilities and abundance																
Model p(.)		Capture probs p						Capture probs p								
Winter	Asia	HI	Mx-Rev	Baja	IX-Main	CentAm	Sum	Summer	ssia-Kam	Al-Ber	GOA	AK-NBC	IC-NWA	CA-OR	Sum	
All	0.234	0.095	0.279	0.024	0.346	0.109		All	0.044	0.048	0.142	0.214	0.290	0.166		
Estimated abundance							Estimated abundance									
2004	781	7332	1106	7298	608	166	17290	2004	565	5997	6516	5445	248	1498	20268	
2005	892	8853	689	6214	729	414	17791	<u>2005</u>	<u>859</u>	<u>6765</u>	<u>3728</u>	<u>3219</u>	<u>469</u>	<u>1830</u>	<u>16868</u>	
<u>2006</u>	<u>1225</u>	<u>10733</u>	<u>657</u>	<u>3420</u>	<u>917</u>	<u>414</u>	<u>17366</u>									
Average	966	8973	817	5644	751	331	17482	Average	712	6381	5122	4332	358	1664	18568	
			MX:	7213												
Model p(t)		Capture probs p						Capture probs p								
Winter	Asia	HI	Mx-Rev	Baja	IX-Main	CentAm	Sum	Summer	ssia-Kam	Al-Ber	GOA	AK-NBC	IC-NWA	CA-OR	Sum	
2004	Capture prob's in W04 cannot be estimated							2004	0.338	0.016	0.392	0.443	0.220	0.222		
2005	0.208	0.103	0.285	0.039	0.371	0.186		2005	0.386	0.034	0.158	0.220	0.299	0.175		
2006	0.249	0.090	0.271	0.014	0.333	0.089										
Estimated abundance							Estimated abundance									
2004	Abundance in W04 cannot be estimated							2004	74	17546	2356	2635	327	1119	24058	
2005	1004	8150	674	3798	679	242	14547	<u>2005</u>	<u>99</u>	<u>9642</u>	<u>3334</u>	<u>3132</u>	<u>455</u>	<u>1731</u>	<u>18393</u>	
<u>2006</u>	<u>1152</u>	<u>11346</u>	<u>674</u>	<u>5817</u>	<u>952</u>	<u>505</u>	<u>20445</u>									
Average	1078	9748	674	4807	816	373	17496	Average	86	13594	2845	2883	391	1425	21225	
			MX:	6297												
Model p(n)		Capture probs p						Capture probs p								
Winter	Asia	HI	Mx-Rev	Baja	IX-Main	CentAm	Sum	Summer	ssia-Kam	Al-Ber	GOA	AK-NBC	IC-NWA	CA-OR	Sum	
2004	0.165	0.069	0.452	0.039	0.270	0.043		2004	0.250	0.027	0.273	0.370	0.380	0.146		
2005	0.189	0.083	0.282	0.033	0.324	0.107		2005	0.381	0.031	0.156	0.219	0.718	0.178		
2006	0.259	0.101	0.269	0.018	0.408	0.107										
Estimated abundance							Estimated abundance									
2004	1107	10103	681	4471	777	420	17558	2004	100	10534	3375	3156	189	1702	19056	
2005	1107	10103	681	4471	777	420	17558	<u>2005</u>	<u>100</u>	<u>10534</u>	<u>3375</u>	<u>3156</u>	<u>189</u>	<u>1702</u>	<u>19056</u>	
<u>2006</u>	<u>1107</u>	<u>10103</u>	<u>681</u>	<u>4471</u>	<u>777</u>	<u>420</u>	<u>17558</u>									
Average	1107	10103	681	4471	777	420	17558	Average	100	10534	3375	3156	189	1702	19056	
			MX:	5928												
Model p(S., Wn)		Capture probs p						Capture probs p								
Winter	Asia	HI	Mx-Rev	Baja	IX-Main	CentAm	Sum	Summer	ssia-Kam	Al-Ber	GOA	AK-NBC	IC-NWA	CA-OR	Sum	
2004	0.165	0.069	0.452	0.039	0.270	0.043		2004	0.044	0.048	0.142	0.214	0.290	0.166		
2005	0.189	0.083	0.282	0.033	0.324	0.107		2005	0.044	0.048	0.142	0.214	0.290	0.166		
2006	0.259	0.101	0.269	0.018	0.408	0.107										
Estimated abundance							Estimated abundance									
2004	1107	10100	681	4470	777	420	17555	2004	565	5997	6516	5445	248	1498	20269	
2005	1107	10100	681	4470	777	420	17555	<u>2005</u>	<u>858</u>	<u>6765</u>	<u>3728</u>	<u>3219</u>	<u>469</u>	<u>1830</u>	<u>16870</u>	
<u>2006</u>	<u>1107</u>	<u>10100</u>	<u>681</u>	<u>4470</u>	<u>777</u>	<u>420</u>	<u>17555</u>									
Average	1107	10100	681	4470	777	420	17555	Average	712	6381	5122	4332	359	1664	18570	
			MX:	5928												

Table 19. Migration rates for four Non-Markov models described in Table 18.

Model p(.)	S->W							W->S							S->S							W->W							
	OSW	Asia	HI	Mx-Rev	Baja	MX-Mai	CentAm	OSW	Asia-Kam	Al-Ber	GOA	SEAK-N	SBC-NW	CA-OR	OSW	Russia	Al-Ber	GOA	SEAK-N	SBC-NW	CA-OR	OSW	Asia	HI	Mx-Rev	Baja	MX-Mai	CentAm	
Russia	1.000	0.000	0.000	0.000	0.000	0.000	0.000	Asia	0.724	0.276	0.000	0.000	0.000	0.000	Russia	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.972	0.028	0.000	0.000	0.000	0.000	0.000
Al-Ber	0.030	0.480	0.016	0.459	0.015	0.000	0.000	HI	0.000	0.316	0.295	0.377	0.012	0.000	Al-Ber	0.000	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.996	0.003	0.000	0.000	0.001	0.000
GOA	0.005	0.524	0.030	0.425	0.016	0.000	0.000	Mx-Rev	0.000	0.451	0.476	0.063	0.010	0.000	GOA	0.000	0.000	0.989	0.005	0.000	0.005	Mx-Rev	0.000	0.096	0.717	0.162	0.025	0.000	
SEAK-N	0.000	0.875	0.013	0.106	0.006	0.000	0.000	Baja	0.000	0.474	0.254	0.071	0.041	0.160	SEAK-N	0.000	0.000	0.023	0.965	0.011	0.000	Baja	0.000	0.022	0.017	0.860	0.088	0.013	
SBC-NW	0.000	0.509	0.000	0.299	0.152	0.040	0.000	MX-Mai	0.000	0.180	0.091	0.025	0.047	0.658	SBC-NW	0.000	0.000	0.000	0.000	1.000	0.000	MX-Mai	0.000	0.000	0.000	0.339	0.614	0.047	
CA-OR	0.000	0.000	0.000	0.588	0.173	0.239	0.000	CentAm	0.000	0.000	0.000	0.000	0.092	0.908	CA-OR	0.000	0.000	0.000	0.000	0.013	0.987	CentAm	0.000	0.000	0.000	0.000	0.076	0.924	

Model p(t)	S->W							W->S							S->S							W->W							
	OSW	Asia	HI	Mx-Rev	Baja	MX-Mai	CentAm	OSW	Asia-Kam	Al-Ber	GOA	SEAK-N	SBC-NW	CA-OR	OSW	Russia	Al-Ber	GOA	SEAK-N	SBC-NW	CA-OR	OSW	Asia	HI	Mx-Rev	Baja	MX-Mai	CentAm	
Russia	1.000	0.000	0.000	0.000	0.000	0.000	0.000	Asia	0.049	0.951	0.000	0.000	0.000	0.000	Russia	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.970	0.030	0.000	0.000	0.000	0.000	0.000
Al-Ber	0.029	0.476	0.016	0.464	0.015	0.000	0.000	HI	0.000	0.562	0.163	0.262	0.013	0.000	Al-Ber	0.000	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.996	0.003	0.000	0.000	0.001	0.000
GOA	0.005	0.526	0.030	0.424	0.015	0.000	0.000	Mx-Rev	0.000	0.717	0.232	0.040	0.011	0.000	GOA	0.000	0.000	0.989	0.006	0.000	0.005	Mx-Rev	0.000	0.095	0.718	0.161	0.025	0.000	
SEAK-N	0.000	0.876	0.013	0.105	0.006	0.000	0.000	Baja	0.000	0.669	0.122	0.045	0.042	0.122	SEAK-N	0.000	0.000	0.022	0.967	0.011	0.000	Baja	0.000	0.021	0.017	0.860	0.089	0.013	
SBC-NW	0.000	0.505	0.000	0.308	0.151	0.036	0.000	MX-Mai	0.000	0.330	0.051	0.017	0.050	0.552	SBC-NW	0.000	0.000	0.000	0.000	1.000	0.000	MX-Mai	0.000	0.000	0.000	0.342	0.613	0.044	
CA-OR	0.000	0.000	0.000	0.619	0.173	0.208	0.000	CentAm	0.000	0.000	0.000	0.000	0.103	0.897	CA-OR	0.000	0.000	0.000	0.000	0.013	0.987	CentAm	0.000	0.000	0.000	0.000	0.068	0.932	

Model p(n)	S->W							W->S							S->S							W->W							
	OSW	Asia	HI	Mx-Rev	Baja	MX-Mai	CentAm	OSW	Asia-Kam	Al-Ber	GOA	SEAK-N	SBC-NW	CA-OR	OSW	Russia	Al-Ber	GOA	SEAK-N	SBC-NW	CA-OR	OSW	Asia	HI	Mx-Rev	Baja	MX-Mai	CentAm	
Russia	1.000	0.000	0.000	0.000	0.000	0.000	0.000	Asia	0.054	0.946	0.000	0.000	0.000	0.000	Russia	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.970	0.030	0.000	0.000	0.000	0.000	0.000
Al-Ber	0.030	0.482	0.016	0.459	0.014	0.000	0.000	HI	0.000	0.505	0.200	0.288	0.007	0.000	Al-Ber	0.000	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.996	0.003	0.000	0.000	0.001	0.000
GOA	0.005	0.540	0.031	0.410	0.015	0.000	0.000	Mx-Rev	0.000	0.653	0.295	0.045	0.007	0.000	GOA	0.000	0.000	0.989	0.006	0.000	0.005	Mx-Rev	0.000	0.095	0.724	0.158	0.024	0.000	
SEAK-N	0.000	0.887	0.013	0.095	0.006	0.000	0.000	Baja	0.000	0.626	0.153	0.050	0.025	0.145	SEAK-N	0.000	0.000	0.022	0.973	0.005	0.000	Baja	0.000	0.019	0.018	0.873	0.077	0.013	
SBC-NW	0.000	0.507	0.000	0.313	0.140	0.040	0.000	MX-Mai	0.000	0.264	0.062	0.019	0.028	0.628	SBC-NW	0.000	0.573	0.000	0.000	0.427	0.000	MX-Mai	0.000	0.000	0.000	0.373	0.578	0.049	
CA-OR	0.000	0.000	0.000	0.597	0.162	0.241	0.000	CentAm	0.000	0.000	0.000	0.000	0.050	0.950	CA-OR	0.000	0.000	0.000	0.000	0.007	0.993	CentAm	0.000	0.000	0.000	0.000	0.063	0.937	

Model p(S,V)	S->W							W->S							S->S							W->W							
	OSW	Asia	HI	Mx-Rev	Baja	MX-Mai	CentAm	OSW	Asia-Kam	Al-Ber	GOA	SEAK-N	SBC-NW	CA-OR	OSW	Russia	Al-Ber	GOA	SEAK-N	SBC-NW	CA-OR	OSW	Asia	HI	Mx-Rev	Baja	MX-Mai	CentAm	
Russia	1.000	0.000	0.000	0.000	0.000	0.000	0.000	Asia	0.724	0.276	0.000	0.000	0.000	0.000	Russia	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.970	0.030	0.000	0.000	0.000	0.000	0.000
Al-Ber	0.030	0.482	0.016	0.459	0.014	0.000	0.000	HI	0.000	0.316	0.295	0.377	0.012	0.000	Al-Ber	0.000	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.996	0.003	0.000	0.000	0.001	0.000
GOA	0.005	0.540	0.031	0.410	0.015	0.000	0.000	Mx-Rev	0.000	0.451	0.476	0.063	0.010	0.000	GOA	0.000	0.000	0.990	0.005	0.000	0.005	Mx-Rev	0.000	0.095	0.724	0.158	0.024	0.000	
SEAK-N	0.000	0.887	0.013	0.095	0.006	0.000	0.000	Baja	0.000	0.474	0.254	0.071	0.041	0.160	SEAK-N	0.000	0.000	0.023	0.965	0.011	0.000	Baja	0.000	0.019	0.018	0.873	0.077	0.013	
SBC-NW	0.000	0.507	0.000	0.313	0.140	0.040	0.000	MX-Mai	0.000	0.180	0.091	0.025	0.047	0.657	SBC-NW	0.000	0.000	0.000	0.000	1.000	0.000	MX-Mai	0.000	0.000	0.000	0.373	0.578	0.049	
CA-OR	0.000	0.000	0.000	0.597	0.162	0.241	0.000	CentAm	0.000	0.000	0.000	0.000	0.092	0.908	CA-OR	0.000	0.000	0.000	0.000	0.013	0.987	CentAm	0.000	0.000	0.000	0.000	0.063	0.937	

Movement rates from winter to summer areas were not as similar among models but usually were within 10% of each other. In contrast with summer to winter movement, whales in most areas in the winter had multiple destinations in the summer. There was one important difference among models that greatly affected estimates of model parameters and abundance. Two destination areas (Russia, AI-Ber) were confounded in the Asia recaptures. For models p(.) and p(S.Wn), most Asia whales were estimated to move to Russia. However, for models p(t) and p(n), most Asia whales were estimated to move to the AI-Ber area. This caused the probability of capture in the AI-Ber area to drop and increased its estimate of abundance. Site fidelity from winter to winter and from summer to summer was generally near 1 for the same area (Table 19). The only exception was the set of three Mexico areas, which had an appreciable amount of interchange.

Movement rates from the non-Markovian models were often much different than for the Markovian models. This follows in part from the Aleutian-Bering Sea sampling problem described above, in that the higher abundance in that area requires movement from wintering areas. It appears that there are several combinations of movement parameters and capture probabilities that explain the data equally well. Additional modeling and analysis will be necessary to delineate area-specific knowledge.

The Petersen estimates using pooled-region winter and feeding areas yielded consistently higher estimates than the geographically stratified Hilborn models. The Petersen estimate could be biased upward if sampling of different feeding areas and different wintering areas occurred disproportionately in a manner that biased the number of recaptures downward. This appears to be the case in the western North Pacific, where we had a relatively thorough sample but where feeding areas were not well sampled (based on the low proportion of the animals on the Asian wintering areas that had been seen on any feeding area).

Previous estimates of abundance and determination of trends

A number of estimates have been made for humpback whale abundance in some of the areas covered by SPLASH. North Pacific humpback whale populations were estimated to be at about 15,000 prior to commercial exploitation in the twentieth century (Rice 1978), however, this estimate was based on whaling data that may have been inaccurate. Approximate numbers in the North Pacific after the end of commercial whaling were estimated at about 1,400 (Gambell, 1976) and 1,200 (Johnson and Wolman 1984). Barlow (1994b) suggested that abundance was greater than 3,000 whales in the early 1990s.

The only other basin-wide mark-recapture abundance estimate was the NPAC study made using post-hoc analysis from photo-identification photographs taken from 1990 to 1993 (Calambokidis *et al.* 1997). These estimates were determined using several geographically stratified capture-recapture models that yielded estimates of 6,000 (models using wintering areas only and subject to male bias) to 10,000 (models using wintering and feeding areas). The primary limitations of the NPAC study were the lack or limited coverage of many known wintering and feeding areas. This included no samples from feeding areas in the western North Pacific and only limited coverage of feeding areas in the central North Pacific. There was also no coverage of some critical wintering areas including the Philippines and Central America, and in

most years, the samples from Mexico were incomplete or only from one or two of the three main subareas.

Despite limitations in past estimates of humpback whales abundance in the North Pacific, the SPLASH results can selectively be compared to some of those estimates to examine rates of annual increase (Table 20). For the overall North Pacific, comparison of the most complete NPAC estimate using all feeding and wintering areas was 9,819. Over the 13-year span between these two studies, a 4.9% annual increase would be required to reach the best total estimate from SPLASH. Going back to the estimate of 1,400 whales at the end of whaling for humpbacks in 1966, a 6.8% annual increase over the 39-year period would be required to reach the current SPLASH abundance.

For several key sub-areas, a reasonable quantitative comparison could be made to SPLASH estimates. For Hawaii, three methods were used to compare estimates to determine trends based on NPAC and while the absolute abundance in these estimates had certain biases, the annual rates of increase were very similar and ranged from 5.5 to 6.0% (Table 20). Asia was more problematic because of differences in sampling but the winter-winter mark-recapture estimates between NPAC and SPLASH indicated a 6.7% rate of increase. This estimate could be biased upwards because of the more complete sampling effort in SPLASH (including sampling the Philippines).

Table 20. Estimates of annual increases in humpback whale abundance based on comparison to previous estimates and those with similar methods. Primary basis for 1991 to 1993 estimates is from NPAC study (Calambokidis et al. 1997, 2001) with recalculation of abundances to match samples described in table.

Region/basis	Previous		Current		Yr span	Annual incr.
	Yr	Estimate	Yr	Estimate		
Total N Pacific estimates						
Hilborn best NPAC to best SPLASH	1991-93	9,819	2004-06	18,307	13	4.9%
Rice to best SPLASH Hilborn	1966	1,400	2004-06	18,307	39	6.8%
Hawaii estimates						
Adj. year Petersen NPAC to SPLASH	1991-93	3,556	2004-06	7,120	13	5.5%
Hilborn – Wint/Feed NPAC-SPLASH	1991-93	3,760	2004-06	8,034	13	6.0%
Petersen using SEAK marks	1991-93	5,151	2004-06	10,425	13	5.6%
Asia						
Adj year Petersen NPAC to SPLASH	1991-93	405	2004-06	943	13	6.7%

SPLASH estimates will provide a much better basis for both the regional stock definitions and abundances currently used by NMFS in their Stock Assessment reports (Carreta et al. 2007). The minimum population estimate for humpback whales for the Eastern North Pacific stock of humpback whales is 1,158 whales (Carreta et al. 2007) using the capture-recapture estimates by Calambokidis et al. (2004a) with an increasing trend in numbers of 6-7%/year. Abundance for the Central North Pacific was 3,698 (Angliss and Outlaw 2007) using capture-recapture estimates from Calambokidis et al. (1997). Population trends for this region were estimated to increase at 10%/year (Mizroch et al. 2004) and 6.6%/year (Zerbini et al. 2006). Estimates for the Western North Pacific were 367 (Angliss and Outlaw 2007) using capture-recapture estimates from Calambokidis et al. (1997).

Comparison to past regional estimates of abundance

There were also earlier estimates of humpback whale abundance for Hawaii that can not be quantitatively compared to SPLASH because these involved even less complete sampling or different methods. In the early 1980s, abundance was estimated at about 900-1400 whales (Darling *et al.* 1983, Darling and Morowitz 1986, Baker and Herman 1987) using photo-identification data applied to discovery rates (Darling and Morowitz 1986) or a weighted Petersen mark-recapture model (Baker and Herman 1987). Later estimates were made using photo-identification techniques or aerial survey line-transect data that estimated about 3,000-5,000 whales in Hawaiian waters in the 1990s (Calambokidis *et al.* 1997, Cerchio 1998, Mobley *et al.*). Mobley *et al.* (1999, 2001) showed increasing numbers of whales and estimated a 7% increase between 1993 and 2000.

While a quantitative comparison of SPLASH estimates to past estimates could not be made for Mexico, two separate abundance estimates were made for the Mexican waters in the early 1990s. Using a modified model of the Jolly-Seber population model, Urbán *et al.* (1999) estimated that in 1991 there were 1813 (95% CI: 918-2505) whales in the coastal stock and 914 (CI: 590-1193) whales in the Revillagigedo Archipelago stock. Calambokidis *et al.* (1997) found disparate results using two different estimation models; 1,600 whales for all Mexican waters were estimated using the Darroch method and 4,200 whales using the Chapman/Petersen method, citing problems with uneven sampling between areas in Mexico and concluding that the true abundance of about 2,200-2,800 whales was likely between these two estimates, and more consistent with the estimate of Urbán *et al.* (1999). An increase from about 2,500 whales in the early 1990s to the SPLASH estimate of 5,928 would be consistent with a 6.9% rate of annual increase, but should be interpreted cautiously given the variability in the earlier estimates.

While other estimates prior to NPAC and SPLASH are not available for Asia, whaling data provides some indications that there was a much larger population present there than even the current SPLASH estimates. Whalers killed large numbers of humpback whales in these areas: 3,277 animals were killed between 1910 and 1965; 970 of these were killed off Okinawa primarily between 1958 and 1961, 817 were killed off Ogasawara between 1924 and 1944 (Nishiwaki 1959, Rice 1978).

Estimates of humpback whale abundance along the US West Coast from SPLASH (1,702 for California-Oregon from best Hilborn model) agree with other recent estimates of this region. Abundance has been estimated using aerial (Forney *et al.* 1995) and ship line-transect surveys (Barlow 1995, 2003, Barlow and Gerodette 1996, Calambokidis and Barlow 2004, Forney *et al.* 1995, Barlow and Forney 2007) and using photo-identification data to make capture-recapture estimates (Calambokidis *et al.* 1990, 2004a, Calambokidis and Barlow 2004). Abundances estimated off central California in the 1980s were about 300 animals (Dohl *et al.* 1993, Calambokidis *et al.* 1990). In the early 1990s, estimates were about 600 whales (Calambokidis *et al.* 1993, Barlow 1995) and increased at about 8% per year to just below 1000 whales by 1997 (Calambokidis and Barlow 2004). Since 1998, mark-recapture estimates for this area have been more variable with a dramatic drop in abundance in 1999-2001 followed by a rapid increase driven by an apparent influx of new animals that had not been seen in the area previously (Calambokidis *et al.* 2004a, 2005). Line-transect surveys estimated 1,769 (CV=0.16) animals for

surveys pooled between 1991 and 2005 between California and Washington (including northern Washington; Barlow and Forney 2007).

Humpback whale abundance estimates off the Washington coast have been made from vessel line-transect surveys and capture-recapture from photo-identification research and are generally consistent with the low estimates of fewer than 500 for the Washington- S British Columbia from SPLASH. Forney (2007) estimate the number of humpback whales off N Washington/S British Columbia at 208 (CV=0.28) in 2005. Vessel line-transect surveys just in the northern Washington area estimated about 100 whales between 1995 to 2000; however, the estimate in 2002 was substantially higher (562, CV=0.21), although this high estimate may be biased due to resighting animals multiple times (Calambokidis *et al.* 2004b). Capture-capture estimates showed the number of whales increasing from about 100 to 200 from 1995 to 2002 (Calambokidis *et al.* 2004b).

There have been a number of past estimates of humpback whales for SE Alaska but these did not include the wide geographic coverage of SPLASH and yielded much lower estimates than SPLASH. Early estimates of abundance for Southeast Alaska were about 300 whales from 1979 to 1983 (Baker *et al.* 1985). In 1986, Baker *et al.* (1992) estimated 547 whales (95% CL: 504-590). Straley (1994) estimated 404 humpback whales from 1985 through 1992. The most recent estimate was conducted in 2000 where abundance for northern Southeast Alaska was 961 (95% CI 657-1076) humpback whales (Straley *et al.* In Press).

Recent estimates of humpback whale abundance from other areas in Alaska have been even more limited. Vessel line-transect surveys long the Aleutian chain (west of St Matthew Island, north of the 200m bathymetric contour, south of the US/Russia boundary) estimated 1,175 (95% CI:197-7009) whales for this entire central Bering Sea region; however, sightings in this study were too clumped to provide a reliable estimate. Zerbini *et al.* (2006) estimated 1,652 whales (95% CI:1142-2398) along the Aleutian Islands and the Alaska Peninsula (from Kenai to Unimak Pass including Kodiak, the Shumagin Islands and north of Unimak Pass). Photo-identification studies have estimated 100-200 in Prince William Sound and Kenai Peninsula waters (Waite *et al.* 1999, von Ziegesar *et al.* 2000), 100-150 in the Barren Islands (G. Strong, pers. comm.), 300-500 in Kodiak waters (Waite *et al.* 1999), and 410 in the Shumagin Islands (Witteveen *et al.* 2004).

Although little is known regarding the distribution and numbers of humpback whales summering off the Russian Far East, historic whaling records and recent field observations suggest that this region serves as one of the migratory destinations for some portion of the North Pacific population (Doroshenko 2000, Melnikov 2000). In general, humpback whales are thought to occur in relatively coastal waters between the northern Chukotka Peninsula and the southern Kamchatka Peninsula. Recent records have noted small numbers of whales (< 30) off Chukotka, Olutorsky Cape, Karaginskiy Island and the Commander Islands.

CONCLUSION

The SPLASH project shows that humpback whales in the North Pacific have substantially recovered from whaling to a current population size of approximately 20,000. All areas with sufficient data show growing populations since the end of whaling in the 1960s and since the last substantial study of their abundance in the early 1990s. The SPLASH study also shows, for the first time, the detailed patterns of humpback whale migrations between their feeding areas in the North and their wintering areas in the South. Mysteries remain however, and SPLASH data reveals the likely existence of an undiscovered wintering area for many of the whales that feed in the Commander and Aleutian Islands and in the Bering Sea. This report is just the first of what is expected to be many publications to examine the SPLASH data in more detail.

ACKNOWLEDGMENTS

SPLASH was a collaborative effort of over 400 researchers and 50 research groups (see below) who made this project possible. Primary support for SPLASH came from a number of agencies and organizations including the National Marine Fisheries Service, the National Marine Sanctuary Program, National Fish and Wildlife Foundation, Pacific Life Foundation, Department of Fisheries and Oceans Canada, and Commission for Environmental Cooperation with additional support from a number of other organizations and governments for effort in specific regions. Matching of the SPLASH photographs primarily took place at Cascadia Research by Andrea Havron, Jessica Huggins, Dominique Camacho, Kiirsten Flynn, Andrea Bendlin, and Nora Moloney. Gretchen Steiger helped write and review the final report.

SPLASH participants grouped by primary affiliation.

Name	Primary affiliation for SPLASH data collection
Fred Sharpe	Alaska Whale Foundation
Alisa Schulman	American Cetacean Society
Bernardo Alps	American Cetacean Society
Andy Day	ANZO Cetos Research Organization
Ann Zoidis	ANZO Cetos Research Organization
Julie Rivers	ANZO Cetos Research Organization
Alessandro Ponzo	ASLC/Kamchatka Branch of Pacific Institute of Geography
Alexander Burdin	ASLC/Kamchatka Branch of Pacific Institute of Geography
Alexander Volkov	ASLC/Kamchatka Branch of Pacific Institute of Geography
Daniela Maldini	ASLC/Kamchatka Branch of Pacific Institute of Geography
Haruko Sato	ASLC/Kamchatka Branch of Pacific Institute of Geography
Juriy Artukhin	ASLC/Kamchatka Branch of Pacific Institute of Geography
Peter Nilsson	ASLC/Kamchatka Branch of Pacific Institute of Geography
Sergey Frolov	ASLC/Kamchatka Branch of Pacific Institute of Geography
Tim Ciosek	ASLC/Kamchatka Branch of Pacific Institute of Geography
Vladimir Burkanov	ASLC/Kamchatka Branch of Pacific Institute of Geography
Debbie Ferrari	CFWS Center for Whale Studies
Mark Ferrari	CFWS Center for Whale Studies
Barbara LaCorte	Channel Islands Naturalist Corps
Bill Weinerth	Channel Islands Naturalist Corps
Carolyn McCleskey	Channel Islands Naturalist Corps
Don Gillies	Channel Islands Naturalist Corps
Don Shubert	Channel Islands Naturalist Corps
Donna Peake	Channel Islands Naturalist Corps
Eileen Avery	Channel Islands Naturalist Corps
Geoff Grow	Channel Islands Naturalist Corps
Glenn Shuart	Channel Islands Naturalist Corps
John Kuizenga	Channel Islands Naturalist Corps
Josa Kaye-Carr	Channel Islands Naturalist Corps
Linda Flegal	Channel Islands Naturalist Corps
Lisa Angle	Channel Islands Naturalist Corps
Morgan Coffey	Channel Islands Naturalist Corps
Scott Leon	Channel Islands Naturalist Corps
Shari Smith	Channel Islands Naturalist Corps

Elliot Breslar	Channel Islands Naturalist Corps
Michael Carver	Cordell Bank National Marine Sanc.
Adrian Reich	Cornell University
Alberto Abad	Cornell University
Danielle Ameen	Cornell University
Danielle Cholewiak	Cornell University
Guilian Buhl	Cornell University
Jonathan Wood	Cornell University
Sal Cerchio	Cornell University
Tom Turowski	Cornell University
Veronica Iriarte	Cornell University
Eduardo Lugo Cabrera	COVISI Conexión con la Vida Silvestre
Gema Claudia Rivera Galicia	COVISI Conexión con la Vida Silvestre
María Eugenia Rodríguez Vázquez	COVISI Conexión con la Vida Silvestre
Zeida Cristina Foubert Corona	COVISI Conexión con la Vida Silvestre
Aaron Barna	CRC Cascadia Research Collective
Alexei Calambokidis	CRC Cascadia Research Collective
Amber Klimek	CRC Cascadia Research Collective
Andrea Havron	CRC Cascadia Research Collective
Andrea Montero	CRC Cascadia Research Collective
Annie Douglas	CRC Cascadia Research Collective
April Rabuck	CRC Cascadia Research Collective
Carlos Aguilera Caderon	CRC Cascadia Research Collective
Chantal Huijbers	CRC Cascadia Research Collective
Dominique Camacho	CRC Cascadia Research Collective
Erin Falcone	CRC Cascadia Research Collective
Frank Garita	CRC Cascadia Research Collective
Greg Schorr	CRC Cascadia Research Collective
Gretchen Steiger	CRC Cascadia Research Collective
Heidi Johnson	CRC Cascadia Research Collective
Jens Koblitz	CRC Cascadia Research Collective
Jessie Huggins	CRC Cascadia Research Collective
John Calambokidis	CRC Cascadia Research Collective
Kate Lomac-MacNair	CRC Cascadia Research Collective
Kate Thomas	CRC Cascadia Research Collective
Kathleen Bresslin	CRC Cascadia Research Collective
Kathryn Jeffs	CRC Cascadia Research Collective
Kevin Emerson	CRC Cascadia Research Collective
Kiirsten Flynn	CRC Cascadia Research Collective
Kristin Rasmussen	CRC Cascadia Research Collective
Lisa Schlender	CRC Cascadia Research Collective
Marco Saborio	CRC Cascadia Research Collective
Michael Smith	CRC Cascadia Research Collective
Nora Moloney	CRC Cascadia Research Collective
Norm Nickle	CRC Cascadia Research Collective
Paulina Godoy Aguilar	CRC Cascadia Research Collective
Robin Baird	CRC Cascadia Research Collective
Russell Sturm	CRC Cascadia Research Collective
Sherwin Cotler	CRC Cascadia Research Collective
Todd Chandler	CRC Cascadia Research Collective

Karel Beets	FIBB Ecotours de México
Octavia Jolley	FIBB Ecotours de México
Oscar Aranda	FIBB Ecotours de México
Oscar Guzon	FIBB Ecotours de México
Pablo Piña	FIBB Ecotours de México
Patricia Cerrillo	FIBB Ecotours de México
Raul Mejia	FIBB Ecotours de México
D. Martinez	Fundacion Keto
D. Palacios	Fundacion Keto
H. Echandi	Fundacion Keto
Janet Neilson	GBNP Glacier Bay National Park
Melissa Senac	GBNP Glacier Bay National Park
Chris Gabriele	GBNP Glacier Bay National Park, HMMC
Mark Deakos	HAMR The Hawaii Association for Marine Education and Research
Jonathan Whitney	HAMR The Hawaii Association for Marine Education and Research
Allan Ligon	HIWS Hawaiian Islands Humpback Whale National Marine Sanctuary
David Mattila	HIWS Hawaiian Islands Humpback Whale National Marine Sanctuary
Amanda Cummins	HIWS Hawaiian Islands Humpback Whale National Marine Sanctuary
Katie Jones	HIWS Hawaiian Islands Humpback Whale National Marine Sanctuary
Saeko Kumagai	HIWS Hawaiian Islands Humpback Whale National Marine Sanctuary
Ed Lyman	HIWS Hawaiian Islands Humpback Whale National Marine Sanctuary
Norm Garon	HIWS Hawaiian Islands Humpback Whale National Marine Sanctuary
Adam Frankel	HMMC Hawai'i Marine Mammal Consortium
Holly Sargeant-Green	HMMC Hawai'i Marine Mammal Consortium
Kim New	HMMC Hawai'i Marine Mammal Consortium
Michael Hoffhines	HMMC Hawai'i Marine Mammal Consortium
Susan Rickards	HMMC Hawai'i Marine Mammal Consortium
Suzanne Yin	HMMC Hawai'i Marine Mammal Consortium
Todd Buczyna	HMMC Hawai'i Marine Mammal Consortium
Andrea McManus	HSU Humboldt State University
Courtney Danley	HSU Humboldt State University
Jeff Jacobsen	HSU Humboldt State University
Joe Kaplan	HSU Humboldt State University
Melissa Crawford	HSU Humboldt State University
Talena Dehmel	HSU Humboldt State University
Adam Quinn	HWRF Hawai'i Whale Research Foundation
Dan Salden	HWRF Hawai'i Whale Research Foundation
David Glickman	HWRF Hawai'i Whale Research Foundation
Peggy Stap	HWRF Hawai'i Whale Research Foundation
Harrison A. Stubbs	HWRF Hawai'i Whale Research Foundation
Heather Ronek	HWRF Hawai'i Whale Research Foundation
Jeff Kalbach	HWRF Hawai'i Whale Research Foundation
Jill Mickelsen	HWRF Hawai'i Whale Research Foundation
Jim Walsh	HWRF Hawai'i Whale Research Foundation
Kevin Carter	HWRF Hawai'i Whale Research Foundation
Tanya Shober	HWRF Hawai'i Whale Research Foundation
Tif Thomas	HWRF Hawai'i Whale Research Foundation
Jim Greenough	J. Straley Investigations
John Straley	J. Straley Investigations
Kathy Peavey	J. Straley Investigations

Finn Straley	J. Straley Investigations
C. Scott Baker	Marine Mammal Institute, Oregon State University
Debbie Steel	Marine Mammal Institute, Oregon State University
Al Ness	MMRC Marine Mammal Research Consultants Ltd.
Joe Mobley	MMRC Marine Mammal Research Consultants Ltd.
Nancy Black	Monterey Bay Whale Watch
Joelle Sweeney	Moss Landing Marine Labs
Craig Matkin	NGOS - North Gulf Oceanic Society
Dave Ellifrit	NGOS - North Gulf Oceanic Society
David Grimes	NGOS - North Gulf Oceanic Society
Eva Saulitus	NGOS - North Gulf Oceanic Society
John Maniscalco	NGOS - North Gulf Oceanic Society
Kathleen Leonard	NGOS - North Gulf Oceanic Society
Laura Patty	NGOS - North Gulf Oceanic Society
Lisa Pytka	NGOS - North Gulf Oceanic Society
Lori Mazzuca	NGOS - North Gulf Oceanic Society
Mike Britain	NGOS - North Gulf Oceanic Society
Sarah Gibbons	NGOS - North Gulf Oceanic Society
Tara Mauro	NGOS - North Gulf Oceanic Society
Tim Markowitz	NGOS - North Gulf Oceanic Society
Tony Sarabio	NGOS - North Gulf Oceanic Society
Olga von Ziegesar	NGOS Eye of the Whale - Opportunistic
Sandy Cronland	NGOS Eye of the Whale - Opportunistic
Shelley Gill	NGOS Eye of the Whale - Opportunistic
Stephaine Zuniga	NGOS Eye of the Whale - Opportunistic
DeWaine Tellefsrud	NGOS Rainbow Connection - Opportunistic
Ginger Strong	NGOS Rainbow Connection - Opportunistic
Jack Montgomery	NGOS Rainbow Connection - Opportunistic
Monica Chase	NGOS Rainbow Connection - Opportunistic
Jerry Sanger	NGOS Sound Access - Opportunistic
Brenda Rone	NMML NMFS Alaska Fisheries Science Center
Candice Emmons	NMML NMFS Alaska Fisheries Science Center
Christy Sims	NMML NMFS Alaska Fisheries Science Center
Elizabeth Mitchell	NMML NMFS Alaska Fisheries Science Center
Janice Waite	NMML NMFS Alaska Fisheries Science Center
John Durban	NMML NMFS Alaska Fisheries Science Center
Kate Sardi	NMML NMFS Alaska Fisheries Science Center
Kathy Hough	NMML NMFS Alaska Fisheries Science Center
Laura Morse	NMML NMFS Alaska Fisheries Science Center
Marilyn Dahlheim	NMML NMFS Alaska Fisheries Science Center
Nancy Friday	NMML NMFS Alaska Fisheries Science Center
Nelson	NMML NMFS Alaska Fisheries Science Center
Paul Wade	NMML NMFS Alaska Fisheries Science Center
Phil Clapham	NMML NMFS Alaska Fisheries Science Center
Yulia Ivashchenko	NMML NMFS Alaska Fisheries Science Center
Kim Parsons	NMML NMFS Northwest Fisheries Science Center
Hermann Meuter	North Coast Cetacean Survey
Janie Wray	North Coast Cetacean Survey
Yuji Sano	OCA Nature Land Sea Kayak
Aiko Sakuraba	OCA Okinawa Churaumi Aquarium

Hidehiko Ono	OCA Okinawa Churaumi Aquarium
Haruna Okabe	OCA Okinawa Churaumi Aquarium
Hisao Miyahira	OCA Okinawa Churaumi Aquarium
Kana Tomiyama	OCA Okinawa Churaumi Aquarium
Kei Kajiwara	OCA Okinawa Churaumi Aquarium
Kiyoteru Tooyama	OCA Okinawa Churaumi Aquarium
Mami Nomura	OCA Okinawa Churaumi Aquarium
Masaya Koami	OCA Okinawa Churaumi Aquarium
Naoto Higashi	OCA Okinawa Churaumi Aquarium
Noriko Minei	OCA Okinawa Churaumi Aquarium
Sae Ishihara	OCA Okinawa Churaumi Aquarium
Shigeo Saino	OCA Okinawa Churaumi Aquarium
Shinji Higa	OCA Okinawa Churaumi Aquarium
Takayuki Sato	OCA Okinawa Churaumi Aquarium
Tomomi Kasagi	OCA Okinawa Churaumi Aquarium
Yukinobu Taira	OCA Okinawa Churaumi Aquarium
Barbara Blackie	Olympic Coast National Marine Sanctuary
Hiroko Yamada	OMC Chiba University
Hiroyuki Suganuma	OMC Everlasting Nature of Asia
Koji Narushima	OMC Everlasting Nature of Asia
Manami Yamaguchi	OMC Ogasawara Marine Center
Yuki Horita	OMC Ogasawara Marine Center
Ai Hamamoto	OMC Ogasawara Marine Center - research volunteer
Ayano Asakura	OMC Ogasawara Marine Center - research volunteer
Chisa Okura	OMC Ogasawara Marine Center - research volunteer
Daiki Inamori	OMC Ogasawara Marine Center - research volunteer
Haruka Sato	OMC Ogasawara Marine Center - research volunteer
Haruna Okabe	OMC Ogasawara Marine Center - research volunteer
Hiromi Kodama	OMC Ogasawara Marine Center - research volunteer
Hitomi Morishita	OMC Ogasawara Marine Center - research volunteer
Miho Jiomaru	OMC Ogasawara Marine Center - research volunteer
Kanae Matsushima	OMC Ogasawara Marine Center - research volunteer
Kaoru Ito	OMC Ogasawara Marine Center - research volunteer
Katsuko Miyoshi	OMC Ogasawara Marine Center - research volunteer
Keiko Kato	OMC Ogasawara Marine Center - research volunteer
Kenji Kitano	OMC Ogasawara Marine Center - research volunteer
Kiichiro Saito	OMC Ogasawara Marine Center - research volunteer
Kotomi Aoki	OMC Ogasawara Marine Center - research volunteer
Kozue Tomomatsu	OMC Ogasawara Marine Center - research volunteer
Madoka Nagai	OMC Ogasawara Marine Center - research volunteer
Mahoko Osanaga	OMC Ogasawara Marine Center - research volunteer
Makoto Ogasawara	OMC Ogasawara Marine Center - research volunteer
Mariko Nakayama	OMC Ogasawara Marine Center - research volunteer
Mariou Uchimiya	OMC Ogasawara Marine Center - research volunteer
Minoru Tamura	OMC Ogasawara Marine Center - research volunteer
Nagisa Sakamoto	OMC Ogasawara Marine Center - research volunteer
Noriko Ishima	OMC Ogasawara Marine Center - research volunteer
Noriko Minei	OMC Ogasawara Marine Center - research volunteer
Norimitsu Sasaki	OMC Ogasawara Marine Center - research volunteer
Nozomi Kobayashi	OMC Ogasawara Marine Center - research volunteer

Ryosuke Aruga	OMC Ogasawara Marine Center - research volunteer
Saori Oda	OMC Ogasawara Marine Center - research volunteer
Sawako Takekura	OMC Ogasawara Marine Center - research volunteer
Sayaka Shiba	OMC Ogasawara Marine Center - research volunteer
Shoko Abe	OMC Ogasawara Marine Center - research volunteer
Shoko Hagiwara	OMC Ogasawara Marine Center - research volunteer
Shota Yoshino	OMC Ogasawara Marine Center - research volunteer
Shunsuke Hata	OMC Ogasawara Marine Center - research volunteer
Sumiko Nakamura	OMC Ogasawara Marine Center - research volunteer
Takayuki Sato	OMC Ogasawara Marine Center - research volunteer
Teruaki Yuta	OMC Ogasawara Marine Center - research volunteer
Tomomi Kasagi	OMC Ogasawara Marine Center - research volunteer
Tomomi Ogura	OMC Ogasawara Marine Center - research volunteer
Tsutomu Iwasaki	OMC Ogasawara Marine Center - research volunteer
Yasumasa Hase	OMC Ogasawara Marine Center - research volunteer
Yasutoki Shibata	OMC Ogasawara Marine Center - research volunteer
Yoko Maruta	OMC Ogasawara Marine Center - research volunteer
Yuki Kawasaki	OMC Ogasawara Marine Center - research volunteer
Yuki Nagao	OMC Ogasawara Marine Center - research volunteer
Yukiko Sakawa	OMC Ogasawara Marine Center - research volunteer
Yuko Sueyoshi	OMC Ogasawara Marine Center - research volunteer
Diane Blair	Opportunistic contributor
Don Doolittle	Opportunistic contributor
Matthew Pike	Opportunistic contributor
Melissa Gong	Opportunistic contributor
Pam Parker	Opportunistic contributor
Roger Deffendall	Opportunistic contributor
Alison Stimpert	OWSI Oceanwide Science Institute
Aran Mooney	OWSI Oceanwide Science Institute
Marc Lammers	OWSI Oceanwide Science Institute
Tom Kieckhefer	Pacific Cetacean Group
Bob Haskel	Pacific Life Foundation (CRC effort)
Ola Svensson	PALCO Charters
Pat Hughes	PALCO Charters
Bill Stehl	Petersburg Marine Mammal Center
Dennis Rogers	Petersburg Marine Mammal Center
Jeff Reynolds	Petersburg Marine Mammal Center
Jim Nahmens	Petersburg Marine Mammal Center
Charlie Smith	S/V Ragland
Claud Renalt	S/V Ragland
Greg Kaolian	S/V Ragland
Stephen Dutton	Sanctuary Cruises
Erin Oleson	Scripps Institution of Oceanography
Josh Jones	Scripps Institution of Oceanography
Amanda Cummins	SWFSC Southwest Fisheries Science Center
Annette Henry	SWFSC Southwest Fisheries Science Center
Barbara Taylor	SWFSC Southwest Fisheries Science Center
Beth Goodwin	SWFSC Southwest Fisheries Science Center
Bob Pitman	SWFSC Southwest Fisheries Science Center
Brittany Hancock	SWFSC Southwest Fisheries Science Center

Carrie LeDuc	SWFSC Southwest Fisheries Science Center
Chris Cutler	SWFSC Southwest Fisheries Science Center
Cornelia Oedekoven	SWFSC Southwest Fisheries Science Center
David Weller	SWFSC Southwest Fisheries Science Center
Gabriela Serra-Valente	SWFSC Southwest Fisheries Science Center
Gary Friedrichsen	SWFSC Southwest Fisheries Science Center
Holly Fearnbach	SWFSC Southwest Fisheries Science Center
Jason Appler	SWFSC Southwest Fisheries Science Center
Jay Barlow	SWFSC Southwest Fisheries Science Center
Jessica Redfern	SWFSC Southwest Fisheries Science Center
Jim Cotton	SWFSC Southwest Fisheries Science Center
Juan Carlos Salinas	SWFSC Southwest Fisheries Science Center
Julie Oswald	SWFSC Southwest Fisheries Science Center
Karin Forney	SWFSC Southwest Fisheries Science Center
Kate Stafford	SWFSC Southwest Fisheries Science Center
Katie Roberts	SWFSC Southwest Fisheries Science Center
Kelly Robertson	SWFSC Southwest Fisheries Science Center
Leigh Torres	SWFSC Southwest Fisheries Science Center
Lilian Carswell	SWFSC Southwest Fisheries Science Center
Linda Hoffman	SWFSC Southwest Fisheries Science Center
Lisa Ballance	SWFSC Southwest Fisheries Science Center
Lisa Munger	SWFSC Southwest Fisheries Science Center
Liz Zele	SWFSC Southwest Fisheries Science Center
Megan Ferguson	SWFSC Southwest Fisheries Science Center
Michael Richlen	SWFSC Southwest Fisheries Science Center
Nicole Hedrick	SWFSC Southwest Fisheries Science Center
Peter Pyle	SWFSC Southwest Fisheries Science Center
Richard Rowlett	SWFSC Southwest Fisheries Science Center
Rick LeDuc	SWFSC Southwest Fisheries Science Center
Robert Holland	SWFSC Southwest Fisheries Science Center
Robert Pitman	SWFSC Southwest Fisheries Science Center
Sage Tezak	SWFSC Southwest Fisheries Science Center
Shannon Rankin	SWFSC Southwest Fisheries Science Center
Siri Hakala	SWFSC Southwest Fisheries Science Center
Sophie Webb	SWFSC Southwest Fisheries Science Center
Suzanne Yin	SWFSC Southwest Fisheries Science Center, HMMC
Adam Pack	TDI The Dolphin Institute
Andrea Bendlin	TDI The Dolphin Institute, Cascadia Research
Anita Basudev	TDI The Dolphin Institute
Elia Herman	TDI The Dolphin Institute
Gemma Clay	TDI The Dolphin Institute
Julien Delarue	TDI The Dolphin Institute
Kathleen Mohning	TDI The Dolphin Institute
Katie Kuker	TDI The Dolphin Institute
Kira Goetschius	TDI The Dolphin Institute
Lou Herman	TDI The Dolphin Institute
Nina Hamacher	TDI The Dolphin Institute
Scott Terna	TDI The Dolphin Institute
Aliza Milette	TDI The Dolphin Institute
Jamie Gibbon	TDI The Dolphin Institute

Laura McCue
Soledad Esnaola
Marianne Rasmussen
Adalberto Herrera
Alejandro Gómez Gallardo
Benjamín Troyo Vega
Carlos Navarro
Christian Ramp
Clara Elena Peréz
Clara Pérez Sánchez
Claudia Díaz Guzmán
Diane Gendron
Esther Jiménez López
Gabriel Aguirre Fernández
Gabriela Díaz Erasles
Gustavo Cárdenas Hinojosa
Jorge Urbán Ramírez
Kirsten Krans
Lourdes García Martínez
Mario Salinas Zacarías
Mauricio Najera
Mike Greenfelder
Renske de Jonge
Sergio Nigenda Morales
Steven Zeff
Sylviane Jaume Schinkel
Urmás Kaldveer
Ursula González Peral
Betsy Wilson
Bree Witteveen
Casey Clark
Jordy Thomson
Kate Wynne
Petra Reiman
Heather Vukelic
Jan Straley
Jennifer Cedarleaf
Rachel Myron
Steve Lewis
Steve Weissberg
Kelly Newton
Axayacatl Brambila Villaseñor
Carlos Alberto López Montalvo
Gloria Eunice Panecatl Urquiza
Hiram Rosales Nanduca
Ivette Ruíz Boijseaneau
Juan Manuel Sánchez Parra
Lorena Viloria Gómora
Luis Medrano González
María de Jesús Vázquez Cuevas

TDI The Dolphin Institute
TDI The Dolphin Institute
TDI The Dolphin Institute
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UABCS Universidad Autónoma de Baja California Sur
UAFB University of Alaska Fairbanks
UAFK - University of Alaska Fairbanks Kodiak Campus
UAFK - University of Alaska Fairbanks Kodiak Campus
UAFK - University of Alaska Fairbanks Kodiak Campus
UAFK - University of Alaska Fairbanks Kodiak Campus
UAFK - University of Alaska Fairbanks Kodiak Campus
UASE University of Alaska Southeast
UASE University of Alaska Southeast
UASE University of Alaska Southeast
UASE University of Alaska Southeast
UASE University of Alaska Southeast
UCSC
Universidad Nacional Autónoma de México
Universidad Nacional Autónoma de México
Universidad Nacional Autónoma de México
Universidad Nacional Autónoma de México
Universidad Nacional Autónoma de México
Universidad Nacional Autónoma de México
Universidad Nacional Autónoma de México
Universidad Nacional Autónoma de México
Universidad Nacional Autónoma de México
Universidad Nacional Autónoma de México

María José Villanueva Noriega	Universidad Nacional Autónoma de México
María Violeta Piña González	Universidad Nacional Autónoma de México
Ricardo Axayácatl Juárez Salas	Universidad Nacional Autónoma de México
Sandra Elizabeth Smith Aguilar	Universidad Nacional Autónoma de México
Sandra Pompa Mansilla	Universidad Nacional Autónoma de México
Sergio Martínez Aguilar	Universidad Nacional Autónoma de México
Eric Ward	University of Washington
John Brandon	University of Washington
Greg Killinger	US Forest Service, Craig, AK
Tyania Diffin	Vallarta Adventures
Lance Barrett-Lennard	Vancouver Aquarium
Flip Nicklin	Whale Trust
Jim Darling	Whale Trust
Meagan Jones	Whale Trust
Alexander Aloy	WWF Philippines
Elson Aca	WWF Philippines
Freya Adamczyk	WWF Philippines - research volunteers
Hussein Macarambon	WWF Philippines - research volunteers
James Ballesteros	WWF Philippines
Jo Marie Acebes	WWF Philippines
Jose Mari Daclan	WWF Philippines
Joel Palma	WWF Philippines
A.G. Sano	WWF Philippines - research volunteers
Alden Tagarino	WWF Philippines - research volunteers
Carl Oliveros	WWF Philippines - research volunteers
Gabriel Timoteo	WWF Philippines - research volunteers
Jamie Angelika Placides	WWF Philippines - research volunteers
Jun Miramonte	WWF Philippines - research volunteers
Kenneth Liwanag	WWF Philippines - research volunteers
Leonard Soriano	WWF Philippines - research volunteers
Mark Anthony Reyes	WWF Philippines - research volunteers
Mark Jason Villa	WWF Philippines - research volunteers
Razcel Salvarita	WWF Philippines - research volunteers
Rex Barrer	WWF Philippines - research volunteers
Rolando Magpayo	WWF Philippines - research volunteers
Zenon Villongco	WWF Philippines - research volunteers
Angeline Telan	WWF Philippines - research volunteers
Cynthia Layusa	WWF Philippines - research volunteers
Grazen Acerit	WWF Philippines - research volunteers
Crissy Canlas	WWF Philippines - research volunteers
Patrick Pattugalan	WWFP 3rd yr. CSU Student BS Marine Science
Cyrel Arquero	WWFP 4th yr. CSU Student BS Fisheries
Edgar Castillo	WWFP 4th yr. CSU Student BS Fisheries
Renier Umengan	WWFP 4rd yr. CSU Student BS Marine Science
Evan Wright	
Jimmer McDonald	
Ski Laniewicz	
Steve Sinelli	

REFERENCES

- Acebes, J.M. 2001. Photographic identification of humpback whales (*Megaptera novaeangliae*) in the Babuyan Islands, Northern Luzon, Philippines. Abstract. Fourteenth Biennial Conference on the Biology of Marine Mammals. November 28-December 3 2002. Vancouver B.C., Canada.
- Acebes, J.M.V., J.D. Darling and M. Yamaguchi. 2007. Status and distribution of humpback whales (*Megaptera novaeangliae*) in northern Luzon, Philippines. *Journal of Cetacean Research and Management* 9:37-43.
- Angliss, R.P. and R.B. Outlaw. 2007. Alaska Marine Mammal Stock Assessments, 2006. NOAA Technical Memorandum NOAA-TM-AFSC-168. 244p.
- Baker, C.S. and L.M. Herman. 1987. Alternative population estimates of humpback whales (*Megaptera novaeangliae*) in Hawaiian waters. *Canadian Journal of Zoology* 65:2818-2821.
- Baker, C.S. L.M. Herman, A. Perry, W.S. Lawton, J.M. Straley and J.H. Straley. 1985. Population characteristics and migration of summer and late-season humpback whales (*Megaptera novaeangliae*) in southeastern Alaska. *Marine Mammal Science* 1:304-323.
- Baker, C. S., L. Herman, A. Perry, W. Lawton, J. Straley, A. Wolman, H. Winn, J. Hall, G. Kaufman, J. Reinke and J. Ostman. 1986. The migratory movement and population structure of humpback whales (*Megaptera novaeangliae*) in the central and eastern North Pacific. *Marine Ecology Progress Series* 31:105-119.
- Baker, C.S., J.M. Straley and A. Perry. 1992. Population characteristics of individually identified humpback whales in southeastern Alaska: summer and fall 1986. *Fishery Bulletin* 90:429-437.
- Baker, C.S., A. Perry, J.L. Bannister, M.T. Weinrich, R.B. Abernethy, J. Calambokidis, J. Lien, R.H. Lambertson, J. Urban Ramirez, O. Vasquez, P.J. Clapham, A. Ailing, S.J. O'Brien, and S.R. Palumbi. 1993. Abundant mitochondrial DNA variation and world-wide population structure in humpback whales. *Proceedings of the National Academy of Science* 90:8239-8243.
- Baker, C.S., R.W. Slade, J.L. Bannister, R.B. Abernathy, M.T. Weinrich, J. Lien, J. Urbán, P. Cockeron, J. Calambokidis, O. Vasquez, and S.R. Palumbi. 1994. Hierarchical structure of mitochondrial DNA gene flow among humpback whales *Megaptera novaeangliae*, world-wide. *Molecular Ecology* 3:313-327.
- Baker, C..S., L. Medrano-Gonzalez, J. Calambokidis, A. Perry, F. Pichler, H. Rosenbaum, J. M. Straley, J. Urban-Ramirez, M. Yamaguchi, and O. von Ziegeler. 1998. Population structure of nuclear and mitochondrial DNA variation among humpback whales in the North Pacific. *Molecular Ecology* 6:695-707.

- Barlow, J. 1994a. Abundance of large whales in California coastal waters: a comparison of ship surveys in 1979/80 and in 1991. Reports of the International Whaling Commission 44:399-406.
- Barlow, J. 1994b. Recent information on the status of large whales in California waters. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-203. 27pp.
- Barlow, J. 1995. The abundance of cetaceans in California waters. Part I: Ship surveys in summer and fall of 1991. Fisheries Bulletin 93:1-14.
- Barlow, J. 2003. Preliminary estimates of the abundance of cetaceans along the U.S. west coast: 1991_2001. Southwest Fisheries Science Center Administrative Report LJ-03-03. Available from SWFSC, 8604 La Jolla Shores Dr., La Jolla CA 92037. 31p.
- Barlow, J. and K.A. Forney. 2007. Abundance and population density of cetaceans in the California Current ecosystem. Fisheries Bulletin 105:509-526.
- Brown, M.R., P.J. Corkeron, P.T. Hale, K.W. Shultz, and M.M. Bryden. 1995. Evidence for a sex-segregated migration in the humpback whale (*Megaptera novaeangliae*). Proceeding of the Royal Society of London B Biological Sciences 259:229–234.
- Burnham, K.P. & Anderson, D.R. 2002. *Model Selection and Inference*. Springer-Verlag, New York.
- Calambokidis, J. and J. Barlow. 2004. Abundance of blue and humpback whales in the eastern North Pacific estimated by capture-recapture and line-transect methods. Marine Mammal Science 20:63-85.
- Calambokidis, J., J. C. Cabbage, G.H. Steiger, K. C. Balcomb, and P. Bloedel. 1990. Population estimates of humpback whales in the Gulf of the Farallones, California. Report to the International Whaling Commission (special issue 12):325-333.
- Calambokidis, J., G. H. Steiger, J. R. Evenson, K. R. Flynn, K. C. Balcomb, D. E. Claridge, P. Bloedel, J. M. Straley, C. S. Baker, O. von Ziegesar, M. E. Dahlheim, J. M. Waite, J. D. Darling, G. Ellis and G. A. Green. 1996. Interchange and isolation of humpback whales off California and other North Pacific feeding grounds. Marine Mammal Science 12:215-226.
- Calambokidis, J. G.H. Steiger, J.M. Straley, T. Quinn, L.M. Herman, S. Cerchio, D.R. Salden, M. Yamaguchi, F. Sato, J.R. Urban, J. Jacobsen, O. VonZeigesar, K.C. Balcomb, C.M. Gabriele, M.E. Dahlheim, N. Higashi, S. Uchida, J.K.B. Ford, Y. Miyamura, P. Ladrón de Guevara, S.A. Mizroch, L. Schlender and K. Rasmussen. 1997. Abundance and population structure of humpback whales in the North Pacific basin. Final Contract Report 50ABNF500113 to Southwest Fisheries Science Center, P.O. Box 271, La Jolla, CA 92038 72pp.
- Calambokidis, J., G. H. Steiger, K. Rasmussen, J. Urbán R., K. C. Balcomb, P. Ladrón de Guevara, M. Salinas Z., J. K. Jacobsen, C. S. Baker, L. M. Herman, S. Cerchio and J. D.

- Darling. 2000. Migratory destinations of humpback whales from the California, Oregon and Washington feeding ground. *Marine Ecology Progress Series* 192:295-304.
- Calambokidis, J., G.H Steiger, J.M Straley, L.M. Herman, S. Cerchio, D.R. Salden, J. Urbán R., J.K. Jacobsen, O. von Ziegesar, K.C. Balcomb, C.M. Gabriele, M.E. Dahlheim, S. Uchida, G. Ellis, Y. Miyamura, P. Ladrón de Guevara P., M. Yamaguchi, F. Sato, S.A. Mizroch, L. Schlender, K. Rasmussen, J. Barlow and T.J. Quinn II. 2001. Movements and population structure of humpback whales in the North Pacific. *Marine Mammal Science* 17:769-794.
- Calambokidis, J., T. Chandler, L. Schlender, K. Rasmussen, and G.H. Steiger. 2002. Research on humpback and blue whales off California, Oregon, and Washington in 2001. Final report to Southwest Fisheries Science Center, La Jolla, CA by Cascadia Research, 218½ W Fourth Ave., Olympia, WA 98501. 50pp.
- Calambokidis, J., T. Chandler, L. Schlender, G.H. Steiger and A. Douglas. 2003. Research on humpback and blue whales off California, Oregon and Washington 2002. Final report to Southwest Fisheries Science Center, La Jolla, California. 47pp. Available at: <http://www.cascadiaresearch.org/reports/rep-MN02.pdf>
- Calambokidis, J., T. Chandler, E. Falcone and A. Douglas. 2004a. Research on large whales off California, Oregon and Washington in 2003. Report to Southwest Fisheries Science Center for Contract #50ABNF100065.
- Calambokidis, J., G.H. Steiger, D.K. Ellifrit, B.L. Troutman and C.E. Bowlby. 2004b. Distribution and abundance of humpback whales (*Megaptera novaeangliae*) and other marine mammals off the northern Washington coast. *Fisheries Bulletin* 102:563-580.
- Calambokidis, J., J. Barlow, E.A. Falcone, L. Schlender, A.B. Douglas, G.H. Steiger, and K.B. Ford. 2005. Changes in abundance of humpback whales off the west coast of the U.S. Abstract (Proceedings) 16th Biennial Conference on the Biology of Marine Mammals, San Diego, CA, December 12-16, 2005.
- Carretta, J.V., K.A. Forney, M.M. Muto, J. Barlow, J. Baker, B. Hanson and M.S. Lowry. 2007. U.S. Pacific Marine Mammal Stock Assessments: 2006. NOAA Technical Memorandum NMFS-SWFSC-398. 312p.
- Cerchio, S. 1998. Estimates of humpback whale abundance off Kauai, Hawaii, 1989 to 1993: evaluating biases associated with sampling the Hawaiian Islands wintering assemblage. *Marine Ecology Progress Series* 175:23-34.
- Clapham, P.J., M. Bérubé, D.K. Mattila. 1995. Sex ratio of the Gulf of Maine humpback whale population. *Marine Mammal Science* 11:227–231.
- Craig A.S., and L.M. Herman. 1997. Sex differences in site fidelity and migration of humpback whales (*Megaptera novaeangliae*) to the Hawaiian waters. *Canadian Journal of Zoology* 75:1923–1933.

- Craig, A.S., and L.M. Herman. 2000. Habitat preferences of female humpback whales *Megaptera novaeangliae* in the Hawaiian Islands are associated with reproductive status. *Marine Ecology Progress Series* 193:209–216.
- Craig, A.S., L.M. Herman, C.M. Gabriele and A.A. Pack. 2003. Migratory timing of humpback whales (*Megaptera novaeangliae*) in the Central North Pacific varies with age, sex and reproductive condition. *Behaviour* 140:981–1001.
- Darling, J. D. and Mori, K. 1993. Recent observations of humpback whales (*Megaptera novaeangliae*) in Japanese waters off Ogasawara and Okinawa. *Can. J. Zool.* 71(2):325-33.
- Darling, J.D. and H. Morowitz. 1986. Census of “Hawaii” humpback whales (*Megaptera novaeangliae*) by individual identification. *Canadian Journal of Zoology* 64:105-111.
- Darling, J.D., K.M. Gibson and G.K. Silber. 1983. Observations on the abundance and behavior of humpback whales (*Megaptera novaeangliae*) off West Maui, Hawaii 1977-1979. Pp. 201-222 in *Communication and behavior of whales*. R. Payne (ed). Westview Press, Boulder, Colorado.
- Darling, J.D., M. Moncur and J. Cleland. 2005. Clayoquot Sound humpback whale fluke identification catalog 1995-2004. West Coast Whale Project. Pacific Wildlife Foundation, Box 384, Tofino, B.C. Canada V0R 2Z0. 20pp.
- Darroch, J.N. 1961. The two-sample capture-recapture census when tagging and sampling are stratified. *Biometrika* 48: 241--260.
- Doroshenko, N V. 2000. Soviet catches of humpback whales (*Megaptera novaeangliae*) in the North Pacific. pp. 48-95. *In: A. V. Yablokov and V. A. Zemsky (eds.) Soviet Whaling Data (1949-1979)*. Center for Russian Environmental Policy, Moscow. 408pp.
- Forney, K.A. 2007. Preliminary estimates of cetacean abundance along the U.S. West Coast and within four National Marine Sanctuaries during 2005. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-406. 27pp.
- Forney, K.A., J. Barlow and J.V. Carretta. 1995. The abundance of cetaceans in California waters. Part II: Aerial surveys in winter and spring of 1991 and 1992. *Fishery Bulletin* 93:15-26.
- Gambell, R. 1976. World whale stocks. *Mammal Review* 6:41-53.
- Hilborn, R. 1990. Determination of fish movement patterns from tag recoveries using maximum likelihood estimators. *Canadian Journal of Fisheries and Aquatic Sciences* 47: 635--643.
- Johnson, J. H., and A. A. Wolman. 1984. The humpback whale, *Megaptera novaeangliae*. *Mar. Fish. Rev.* 46(4):30-37.

- Katona, S., Baxter, B., Brazier, O., Kraus, S., Perkins, J. and Whitehead, H. 1979. Identification of humpback whales by fluke photographs. pp. 33-44. *In: H.E. Winn and B. L. Olla (eds.) Behavior of Marine Animals. Vol. 3. Cetaceans.* Plenum Press, New York and London. i-xix + 438pp.
- Melnikov, V.V. 2000. Humpback whales *Megaptera novaeangliae* off Chukchi Peninsula. *Russian Journal of Oceanology* 4:844-849.
- Medrano-González, L., A. Aguayo-Lobo, J. Urbán-Ramirez and C. S. Baker. 1995. Diversity and distribution of mitochondrial DNA lineages among humpback whales, *Megaptera novaeangliae*, in the Mexican Pacific Ocean. *Canadian Journal of Zoology* 73: 1735-1743.
- Mizroch, S.A., L.M. Herman, J.M. Straley, D.A. Glockner-Ferrari, C. Jurasz, J. Darling, S. Cerchio, C.M. Gabriele, D.R. Salden and O. von Ziegesar. 2004. Estimating the adult survival rate of central North Pacific humpback whales (*Megaptera novaeangliae*). *Journal of Mammalogy* 85:963-972.
- Mobley, Jr., J.R., G.B. Bauer and L.M. Herman. 1999. Changes over a ten-year interval in the distribution and relative abundance of humpback whales (*Megaptera novaeangliae*) wintering in Hawaiian waters. *Aquatic Mammals* 25:63-72.
- Mobley Jr., J.R., S. Spitz, and R. Grotefendt. 2001. Abundance of humpback whales in Hawaiian waters: results of 1993-2000 aerial surveys. Report for Hawaiian Islands Humpback Whale National Marine Sanctuary.
- Mori, K. 1994. Distribution, migration and local movements of humpback whales (*Megaptera novaeangliae*) in the adjacent waters of the Ogasawara (Bonin) Islands, Japan. (in Japanese) Tokai University. Doctoral thesis. 129pp.
- Nishiwaki, M. 1959. Humpback whales in Ryukyuan waters. *Scientific results of the Whales Research Institute Tokyo* 14:49-87.
- Ogasawara Marine Center 2000. Fluke ID Catalog (1987-1994)-Humpback whales in Ogasawara and Okinawa. Marine Environmental Association of Tokyo. 139pp.
- Quinn, T.J., II, and Deriso, R.B. 1999. *Quantitative Fish Dynamics.* Oxford University Press, New York.
- Rasmussen, K., J. Calambokidis and G.H. Steiger. 2003. Humpback whales and other marine mammals off Costa Rica and surrounding waters, 1996-2003. Report by Cascadia Research to the Oceanic Society Expeditions, San Francisco, California. Available at: <http://www.cascadiaresearch.org/reports/REP-OSE-03.pdf>
- Rice, D. W. 1978. The humpback whale in the North Pacific: distribution, exploitation, and numbers. *In: K. S. Norris and R. Reeves (eds.) Report on a Workshop on Problems Related to Humpback Whales (Megaptera novaeangliae) in Hawaii.* Report to the Marine Mammal Commission, July 1977, Washington, DC. 21pp.

- Robbins, J. 2007. Structure and dynamics of the Gulf of Maine humpback whale population. PhD thesis, University of St Andrews, Scotland.
- Seber, G. A. F. 1982. The Estimation of Animal Abundance, Second Ed., Griffin, London.
- Smith, T. D., J. Allen, P. J. Clapham, P. S. Hammond, S. Katona, F. Larsen, J. Lien, D. Mattila, P. J. Palsboll, J. Sigurjónsson, P. T. Stevick and N. Oien. 1999. An ocean-wide mark-recapture study of the North Atlantic humpback whale (*Megaptera novaeangliae*). Marine Mammal Science 15:1-32.
- Steiger, G. H., J. Calambokidis, R. Sears, K. C. Balcomb and J. C. Cabbage. 1991. Movement of humpback whales between California and Costa Rica. Marine Mammal Science 7:306-310.
- Stevick, P. T., P. J. Palsboll, T. D. Smith, M. V. Bravington and P. S. Hammond. 2001. Errors in identification using natural markings: rates, sources, and effects on capture-recapture estimates of abundance. Canadian Journal of Fisheries and Aquatic Sciences 58:1861-1970.
- Straley, J.M. 1994. Seasonal characteristics of humpback whales (*Megaptera novaeangliae*) in southeastern Alaska. M.S. thesis, University of Alaska, Fairbanks. 121pp.
- Straley, J.M., T.J. Quinn II and C.M. Gabriele. In Press. Assessment of mark recapture models to estimate the abundance of a humpback whale feeding aggregation in Southeast Alaska. Journal of Biogeography Special Issue: Southeast Alaska XX:XX.
- Urbán, J. and A. Aguayo. 1987. Spatial and seasonal distribution of the humpback whale, *Megaptera novaeangliae*, in the Mexican Pacific. Marine Mammal Science 3:333-344.
- Urbán, J., C. Alvarez F., M. Salinas Z., J. Jacobsen, K. C. Balcomb III, A. Jaramillo L., P. Ladrón de Guevara P., and A. Aguayo L. 1999. Population size of the humpback whale (*Megaptera novaeangliae*) in waters off the Pacific Coast of Mexico. Fishery Bulletin 97:1017-1024.
- Urbán R., J., A. Jarmillo L., A. Aguayo L., P. Ladrón de Guevara P., M Salinas Z., C. Alvarez F., L. Medrano G., J. K. Jacobsen, K. C. Balcomb., D. E. Claridge, J. Calambokidis, G. H. Steiger, J. M. Straley, O. von Ziegesar, J. M. Waite, S. Mizroch, M. E. Dahlheim, J. D. Darling and C. S. Baker. 2000. Migratory destinations of humpback whales wintering in the Mexican Pacific. Journal of Cetacean Research and Management 2:101-110.
- Von Ziegesar, O., B. Goodwin, and R. Devito. 2000. A catalog of humpback whales in Prince William Sound Alaska (1977-2000). Unpubl Rept., Eye of the Whale Research, Homer AK 99603
- Waite, J. M., M. E. Dahlheim, R. C. Hobbs, S. A. Mizroch, O. von Ziegesar-Matkin, L. M. Herman and J. Jacobsen. 1999. Evidence of a feeding aggregation of humpback whales (*Megaptera novaeangliae*) around Kodiak Island, Alaska. Marine Mammal Science 15:210-220.

- White, G.C. and K.P. Burnham. 1999. Program MARK: Survival estimation from populations of marked animals. *Bird Study Supplement* 46:120-138.
- Witteveen, B.H. 2003. Abundance and feeding ecology of humpback whales (*Megaptera novaeangliae*) in Kodiak Alaska. Master's thesis, University of Alaska Fairbanks. 109pp.
- Witteveen, B. and J. Straley. 2002. Sandpoint, Alaska Humpback whale photo-identification study. Final contract rept TO3324686 to Marine Mammal Commission, 4340 East-West Hwy, Bethesda, MD 29814. 18pp.
- Witteveen, B.H., J.M. Straley, O. von Ziegesar, D. Steel, and C.S. Baker. 2004. Abundance and mtDNA differentiation of humpback whales (*Megaptera novaeangliae*) in the Shumagin Islands, Alaska. *Canadian Journal of Zoology* 82:1352-1359.
- Yamaguchi, M., Suganuma, H., Miyamura, Y., Sato, F. and Mori, K. 1995. The interchanges of humpback whales between wintering grounds in the western North Pacific: the Ogasawara Islands and the Kerama Islands, Ryukyu Archipelago. In: Eleventh Biennial Conference on the Biology of Marine Mammals Abstracts. 124pp.
- Yamaguchi, M., Acebes, J.M. and Miyamura, Y. 2002. The breeding ground distribution of the Humpback whales, *Megaptera novaeangliae*, in the western North Pacific and their trans-movements among the Ogasawara Islands, the Ryukyu Islands and the Philippines. Abstract. Second Conference on Marine Mammals of Southeast Asia. July 2002. Dumaguete, Philippines.
- Zerbini, A.M., J.M. Waite, J.L. Laake and P.R. Wade. 2006. Abundance, trends and distribution of baleen whales off Western Alaska and the central Aleutian Islands. *Deep-Sea Research I* 53:1772-1790.