

# JUNDERS 3 Little Patch

The Sounders visit the Northern Puget Sound region each year typically between March-May to feed on ghost shrimp beds.



Little Patch has been known to arrive earlier than other Sounders on some years to begin feeding in the winter months and has been sighted as early as December!



Scan the QR code to learn more about the Sounders, or visit www.cascadiaresearch.org twitter: @CascadiaRes, Instagram: @cascadiaresearch or facebook: @CascadiaResearch

# Earhart

It is suspected that female Sounders do not enter the Puget Sound when they have a calf as their feeding strategy is risky



In 2017, Earhart was struck by a vessel as she surfaced. Thankfully her injuries were minor! When boating, it is important to be aware you are in whale territory and to proceed with caution.



Scan the QR code to learn more about the Sounders, or visit www.cascadiare

# UUNDERS 27 Shackleton

Shackleton has a special relationship with Earhart. They were first documented together in 1990 utilizing the North Puget Sound area to feed. They have been frequently sighted together ever since!



Shackleton sustained an injury to his tailstock at some point, possibly a result of entanglement, which may impede his ability to raise his fluke. It is extremely rare to see him lift his tail!



The Sounders do not appear to be part of a larger group of whales who feed along the West Coast called the Pacific Group Feeding Group, also known as the PCFG.



Stardust has a reputation for avoiding boats, which can make him difficult to track, but he has returned to the Puget Sound every year since 2003!



The WA DNR and Cascadia Research conducted surveys which changed the regulations to suspend ghost shrimp harvest during the Sounders' feeding months



Patch was harassed by a group of transient killer whales in 2010 in Saratoga Passage, he was observed rolling over during the encounter before the killer whales left the scene.



# Dubnuck

The Sounders create large pits in the sediment when foraging. These pits can be seen from the shore at low tide, and are big enough to be seen via satellite images!



Dubnuck was first sighted wandering Southern Puget Sound in 1991, and made it as far south as Budd Inlet. By April 25 1991, he was seen in Port Susan, near the regular feeding grounds.



The Sounders have been thought to have discovered Puget Sound when looking for extra food sources during years of unusual mortality events.



Cascade has visited Puget Sound every year since 2003! He is notorious for fluking nearly every time he takes a deep dive and seems to have a gregarious nature as he is frequently sighted with other whales.



# 356 Carson

Carson does not return to the Puget Sound every year, sometimes skipping the 100 mile journey off the migration route multiple years in a row.



Carson arrived in the secondary wave of animals joining the group in 1999-2000, coinciding with an Unusual Mortality Event for gray whales, where they were potentially searching for new food sources.



# Gisborne


Gisborne made his way into the North Puget Sound area in March of 2018 and has returned every year since.




Gisborne is part of the PCFG group of gray whales, which consists of ~200 individuals that feed from Northern California to British Columbia each spring, summer and fall, instead of migrating further north with the Eastern North Pacific population.




can the QR code to learn more about the Sounders, or visit www.cascadiaresearch. witter: @CascadiaRes, Instagram: @cascadiaresearch or facebook: @CascadiaRese:





















## UNDERS

Gray whales are filter feeders. They have a narrow head with around 300 plates of yellowish-white baleen that makes up



CRCID 2246 has very distinctive knuckles (ridges along the animal's back). This varies among individuals, and researchers use the shape and spacing of the knuckles to make matches between individuals



The Sounders are spotted feeding around Whidbey and Camano Islands, Saratoga Passage, Port Susan, Gedney/Hat Island, and the Snohomish Delta in Washington's Salish Sea.



Gray whales are primarily benthic feeders, which means they feed off the ocean floor. Gray whales turn on their sides in shallow water and take in sediment and prey, then use their baleen to filter the water out retaining the prey.



# 31 Gretchen

Gray whales are considered mid-size baleen whales that can grow up to 45 feet long and weigh as much as 30-40 tons!



It is presumed Gretchen had calves in 2012 and 2015 when she was absent from North Puget Sound during the typical feeding months.



Scan the QR code to learn more about the Sounders, or visit www.cascadiaresearch. twitter: @CascadiaRes, Instagram: @cascadiaresearch or facebook: @CascadiaRese.

## INDERS

Other names for the Sounders include "Puget Sound Regulars" and "Saratoga Grays."



CRCID 2259's large depigmentation patches on their left side make this individual easy to spot and match between encounters.



in the QR code to learn more about the Sounders, or visit www.cascadiaresearch.org tter: @CascadiaRes, Instagram: @cascadiaresearch or facebook: @CascadiaResearch

Gray whales are currently threatened by ship strikes, pollution, climate change, habitat destruction and entanglement from fishing gear.



Most of the Eastern North Pacific gray whales complete a 10,000 to 12,000 mile round trip migration each year, one of the longest of any mammal! It is thought the Sounders continue north after they leave Puget Sound.



Scan the QR code to learn more about the Sounders, or visit www.cascadiaresear twitter: @CascadiaRes, Instagram: @cascadiaresearch or facebook: @CascadiaRe

# SOUNDERS #2249 Hattie

SOUNDERS FACT:
Gray whales have distinct coloration and markings that can be used to identify them and track their movements over time.



Gray whales adjust their feeding strategy to the available prey, having been known to surface skim to take in prey in the water column as well as feed on ghost shrimp in shallow intertidal areas.



## SOUNDERS

2362Thidwick

You can help protect gray whales by reporting sightings to public sighting platforms like Orca Network, supporting research



The underside of the fluke can also be used to tell individual whales apart, each animal has a unique shape, ridging, and markings. Thidwick has quite the unique fluke!



# #2356 Stalwart

SOUNDERS FACT: The Unusual Mortality Event has triggered increased research on gray whales to assess their body condition and use of the North Puget Sound area across years.



Stalwart has been observed spy hopping, which is when a whale lifts its head above the water, and breaching, which is when a whale throws its body out of the water.



Scan the QR code to learn more about the Sounders, or visit www.cascadiares twitter: @CascadiaRes, Instagram: @cascadiaresearch or facebook: @Cascadi


Gray whales have been observed closely associating with one another, camera tags deployed by Cascadia have shown the whales bump and rub against each other under the surface.



Cascadia has used photos taken during field efforts and contributed by the public, as well as satellite imagery, camera tag deployments, and biopsies for genetic analysis to understand gray whale feeding, health and use of the area.



Scan the QR code to learn more about the Sounders, or visit www.cascadiaresearch.org twitter: @CascadiaRes, Instagram: @cascadiaresearch or facebook: @CascadiaResearch





SOUNDERS FACT:
Tahoma does not appear to have left the Salish Sea since their arrival in January 2022 - they appear to have forgone normal migration to stay and feed.



Tahoma caused quite the uproar in 2022 when someone mistook 2440's white patches for buoys and thought the whale was entangled.



Scan the QR code to learn more about the Sounders, or visit www.cascadiaresearch.org of twitter: @CascadiaRes. Instagram: @cascadiaresearch or facebook: @CascadiaResearch